

How Many People Live Near Protected Areas in Developing Countries? Estimates from Gridded Population Data (2000–2020)

Abstract Impact evaluations of protected areas routinely define “local populations” through spatial proximity, yet the demographic scale implied by these definitions is almost never quantified. We estimate population counts inside protected areas and within 10 km of their boundaries for 75 low- and lower-middle-income countries (LMICs) in 2000 and 2020, using the World Database on Protected Areas and globally harmonized gridded population data (GHSL, with WorldPop as robustness check). India, which represents about 36% of the LMIC population, is excluded because its national protected areas are not publicly available in the WDPA. Results are decomposed by IUCN management category (strict and non-strict). In 2020, out of a combined population of about 2.48 billion people, approximately 61 million (2.5%) lived inside protected areas and 732 million (29.5%) lived inside or within 10 km of their boundaries; the corresponding figures for 2000 are 19 million (1.1%) and 258 million (15.4%). Although the IUCN management category is not recorded for 59% of protected areas, among those for which it is documented PA-adjacent populations are located mostly near non-strict protected areas (IUCN categories IV to VI) rather than strict protection regimes (IUCN categories I to III), largely because non-strict PAs are more numerous and cover more land. These population orders of magnitude are relevant for interpreting the scope of PA impact evaluations but do not themselves constitute impact estimates.

1 Introduction

Protected areas (PAs) were primarily established to achieve ecological purposes (Maxwell et al. 2020). Their implications for human populations have nevertheless been a persistent concern in conservation research (Adams et al. 2004), particularly in low and lower middle income countries where rural households account for a large share of the population and global poverty (Castañeda et al. 2018; Byerlee and Janvry 2008). The expansion of PAs makes this issue increasingly salient: from 2000 to 2023, global terrestrial PA coverage expanded by 62%, reaching 17.6% of land area (UNEP-WCMC and IUCN 2024). This growth is set to accelerate with the COP15 commitment to protect 30% of terrestrial land by 2030 (2022).

In the literature assessing the social and economic implications of protected areas, the effects of conservation are consistently framed as local in nature. Yet what constitutes a “local population” is rarely made explicit in demographic terms, even though these definitions implicitly determine the scale to which empirical results apply and can be compared. Two dominant notions of locality

recur. One focuses on populations living inside protected areas, directly subject to restrictions and management regimes. The other extends locality to populations living outside but near protected areas, on the premise that social effects attenuate with distance, reflecting spatial limits to resource use and interaction documented in earlier work (West et al. 2006).

A 10 km buffer around PA boundaries has emerged as a frequently used threshold. It was the inclusion criteria in the widely cited systematic review on of conservation outcomes by Oldekop et al. (2016). It is also pervasive in individual empirical studies. In one of the few global-scale assessments, Naidoo et al. (2019) analyzed conservation wellbeing outcomes across 34 low- and middle-income countries with Demographic and Health Surveys, defining exposure to protected areas as residing at 10 km or less from PAs. Fisher et al. (2024) used Afrobarometer data to assess social wellbeing, explicitly classifying households based on their distance to protected area boundaries, including a 10 km threshold. Among smaller-scale impact evaluations, spatial definitions vary but converge around similar thresholds. Kandel et al. (Kandel et al. 2022) meta-analyze 30 counterfactual studies. Eleven define exposure through in-out comparisons between populations residing inside protected areas and those outside; two of these explicitly exclude populations living within 10 km and 20 km, respectively, from the control group because they may be affected by spillover effects. Ten studies use distance-based or buffer definitions, five of which refer to a 10 km threshold. Nine rely on administrative units that overlap protected areas to varying degrees, reflecting the spatial resolution of available socioeconomic data. Across all these designs, the population sizes implied by exposure definitions are not reported.

This paper addresses this gap by quantifying the demographic scale of conservation exposure definitions. Rather than assessing conservation impacts or making claims about them, it asks a simpler and different question: when studies of PA social impacts define “local populations” spatially, how many people does this actually correspond to, and how has this magnitude evolved since 2000, as protected area systems have expanded? Using globally harmonized PA boundaries and gridded population data, the paper quantifies population counts inside PAs and within a reference buffer commonly used in the literature, and documents how these numbers vary across countries, protected area categories, and time. Providing population orders of magnitude is necessary for interpreting the scope and external validity of estimated conservation social impacts.

2 Methods

2.1 Study Area

The analysis focuses on countries classified by the World Bank as low-income or lower-middle-income economies in 2020 (Vaggi 2017). South Sudan and Timor-Leste did not exist as sovereign states at the beginning of our analysis period (2000), so we did not include them in the analysis.

India is excluded because data on its nationally designated protected areas are restricted from public access in the WDPA. The WDPA records approximately 990 protected areas for India, but only 90 internationally designated sites (Ramsar wetlands, World Heritage sites, UNESCO-MAB Biosphere Reserves) are publicly available, covering approximately 0.5% of India’s land area; data on the remaining 900 nationally designated sites cannot be viewed or downloaded (UNEP-WCMC

2026). Including these restricted sites, India’s terrestrial PA coverage reaches approximately 7.7%. India has stated that the process of making its data publicly accessible is in progress (Wani et al. 2025), and this data gap has been noted in global assessments (Zhang et al. 2025). Because India accounts for approximately 36% of the total LMIC population, including it with incomplete spatial data would mechanically depress aggregate PA coverage and exposure statistics. The analysis therefore covers 75 LMICs.

The population of interest is defined following spatial exposure concepts commonly used in the literature evaluating the impacts of protected areas. Two reference perimeters are considered: residence inside protected areas, and residence within a 10 km buffer around protected area boundaries. These perimeters are used as illustrative thresholds to quantify the population sizes implied by commonly used definitions of “local populations” in conservation impact studies, not as claims about the spatial extent or magnitude of actual conservation impacts.

2.2 Data Sources

National boundaries were drawn from geoBoundaries version 6.0.0, a standardized and openly available global dataset designed for cross-country comparative analysis (Runfola et al. 2020).

Protected area boundaries, status, and designation year are obtained from the World Database on Protected Areas (WDPA), compiled by UNEP-WCMC in collaboration with IUCN (Bingham et al. 2019; UNEP-WCMC and IUCN 2023). We use the May 2021 WDPA release, which underpinned the final assessment of Aichi Target 11 under the Convention on Biological Diversity (CBD) Strategic Plan for Biodiversity 2011–2020 (UNEP-WCMC and IUCN 2021). This snapshot reflects the culmination of reporting efforts by CBD member countries for the 2020 milestones, and is therefore more likely to be complete, while limiting the inclusion of post-2020 designations recorded with missing status years.

Population estimates are drawn from two globally harmonized gridded datasets. The Global Human Settlement Layer (GHSL) combines census-based population counts with remote sensing and ancillary data to produce gridded estimates at 250 m resolution (Freire et al. 2016). WorldPop applies a machine-learning approach trained on census data and spatial covariates to generate population estimates at 100 m resolution (Stevens et al. 2015). Both datasets have been evaluated against geolocated census data in contexts where such data are available (Leyk et al. 2019; Chen et al. 2020; Thomson et al. 2022). GHSL is used as the main reference dataset, while WorldPop is used to assess the robustness of population orders of magnitude.

2.3 Analytical approach

We processed high-resolution population estimates within and around protected areas across the 75 countries in 2000 and 2020 entirely in R (R Core Team 2023), using the terra (Hijmans 2025) and exactextractr (Baston 2023) packages for raster extraction and sf (Pebesma 2018) for vector operations. This pipeline reimplements in R an earlier version that relied on Google Earth Engine (Gorelick et al. 2017), both to improve reproducibility and to correct two issues identified during revision: a double counting of overlapping buffer zones and a cross-border attribution of protected areas (see below). Given the volume of the underlying rasters, the computation was run on the SSP Cloud, an open cloud-based data-science platform built on the Onyxia project (Avouac et al.

2025). All geometries were processed in an equal-area projection (EPSG:6933), and protected areas were filtered to each country's own boundaries by ISO3 code, so that populations near protected areas located in neighbouring countries are not attributed across borders.

Protected areas were filtered to retain only designated, established, or inscribed sites, excluding UNESCO-MAB Biosphere Reserves – as is common practice in conservation analyses due to their often limited legal protection (Hanson 2022; Coetzer et al. 2014) – and purely marine protected areas, since this study focuses on terrestrial populations. Protected areas were classified into three groups based on reported IUCN management categories, following Lebergers et al. (2020): strict protected areas (status I to III), non-strict protected areas (status IV to VI), and protected areas whose IUCN category is not recorded in the WDPA. Protected areas were considered as existing by the end of the study period if they had a reported designation year less than or equal to 2020 or if the designation year was missing in the May 2021 WDPA release. For analyses of change over time, protected areas with missing designation year were excluded to avoid ambiguity in temporal classification.

For each designation-period class and protected area category, protected area polygons were dissolved and expanded using a 10 km buffer to represent proximity-based reference perimeters. To avoid double counting, the three IUCN category groups were sliced hierarchically (strict, then non-strict, then unknown), and each category's 10 km buffer ring was defined to exclude any area already attributed to a protected area interior or to a higher-priority category; population and area were then summed over these mutually exclusive zones. This exclusive construction, together with a single joint treatment of protected areas with and without a recorded designation year, avoids double counting of overlapping buffer zones. Proximity buffers around protected areas are truncated at international borders.

The data outputs underlying this study are available at https://github.com/BETSAKA/population_near_protected_areas/tree/main/data. The full replication package, including the R code and additional materials, is available at https://github.com/BETSAKA/population_near_protected_areas and archived on Software Heritage (swh:1:dir:2c8237e16e74d94043e37899aed0de6f262ed420).

3 Results

3.1 Population magnitudes

Restricting to populations living inside protected areas refers to a relatively small population: 61 million people, or 2.5% of the combined population of these 75 countries. Extending the exposure definition to include people living within 10 km of PA boundaries increases the encompassed population roughly twelvefold, to 732 million (29.5%). For comparison, in 2000 (restricted to PAs with recorded designation years), 18.8 million people (1.1%) lived inside PAs and 257.6 million inside or within 10 km (15.4% of the population).

The decomposition by PA category (Figure 1) shows that populations adjacent to non-strict PAs (IUCN categories IV to VI) exceed those adjacent to strict PAs (IUCN categories I to III): 207 million inside or within 10 km of non-strict PAs, compared with 159 million near strict PAs. The largest

share, however, is associated with PAs whose IUCN category is not recorded in the WDPa: 366 million people.

Standardizing by PA area and by total boundary length clarifies the source of this difference. Non-strict PAs are both more numerous than strict PAs (2,564 versus 1,160 sites) and slightly more extensive (1.39 versus 1.35 million km²). Their combined boundary length is also larger: 206,921 km for non-strict PAs versus 178,109 km for strict PAs. After dividing nearby population by PA area, the gap narrows but does not disappear: non-strict PAs are associated with 149 people inside or within 10 km per km² of PA, compared with 118 for strict PAs. The same conclusion holds when standardizing by total boundary length: non-strict PAs are associated with 999 people inside or within 10 km per km of boundary, compared with 891 for strict PAs. The larger raw population near non-strict PAs therefore reflects primarily their greater number, total area, and total boundary length, with only a modest additional contribution from higher nearby population density.

To connect these magnitudes to the impact evaluation literature, Table 1 translates common spatial exposure definitions into population orders of magnitude.

Table 1: Implied population magnitudes by exposure definition

All PAs in 2020 (including missing designation year), 75 LMICs excluding India, GHSL population estimates

Exposure definition	Population (millions)	% of 75-country GHSL population
Inside PAs only, all categories	60.8	2.5
Inside strict PAs only	10.4	0.4
Inside non-strict PAs only	19.0	0.8
Inside unknown-category PAs only	31.5	1.3
Inside PAs or within 10 km, all categories	731.5	29.5
Inside or within 10 km of strict PAs	158.7	6.4
Inside or within 10 km of non-strict PAs	206.7	8.3
Inside or within 10 km of unknown-category PAs	366.2	14.8

Source: WDPa May 2021 and GHSL 2020. 'Inside or within 10 km' includes populations inside PAs plus the exclusive 10 km buffer ring.

An in-out comparison – where “treated” populations are those residing inside PAs – implies a study population of approximately 61 million people (2.5% of the population of these 75 countries). The same definition applied only to strict PAs would concern 10 million people, compared with 19 million for non-strict PAs and 31 million for PAs with unrecorded IUCN category. When the spatial definition extends to populations within 10 km, the implied population rises to 732 million (29.5%), of whom 159 million are near strict PAs, 207 million near non-strict PAs, and 366 million near PAs with unrecorded IUCN category. These numbers do not appear in the studies reviewed by Kandel et al. (2022), yet they define the implicit scope to which estimated impacts apply.

3.2 Change over time by PA category

Comparing 2020 with 2000 is subject to an important caveat: designation years are missing for 2,912 out of 9,170 PAs in the WDPA (31.8% across the 75 LMICs studied), so the set of PAs that can be confirmed as existing before 2001 is a subset of those known to exist by 2020. The 2000-2020 comparison is therefore restricted to PAs with recorded designation years and should be interpreted as indicative of broad trends rather than a precise measure of change.

With this caveat in mind, Figure 1 shows how PA-adjacent populations evolved between 2000 and 2020 across PA categories, separately for populations inside PAs and within the 10 km buffer.




Figure 1: Population inside protected areas and within 10 km of protected area boundaries by IUCN category, 2000 versus 2020. Estimates are for 75 low- and lower-middle-income countries excluding India, using GHSL and restricted to protected areas with recorded designation year.

Growth in exposed populations occurred across all three PA categories. The net change (2020 minus 2000) for the 75 LMICs combined (Figure S1, Supplementary Materials) shows that the population inside or within 10 km of non-strict PAs grew by approximately 94 million, that inside or within 10 km of PAs with unrecorded IUCN category by 101 million, and that near strict PAs by 63 million.

This overall increase reflects two distinct mechanisms: population growth near PAs that already existed by 2000, and the designation of new PAs between 2000 and 2020. Using national population growth rates to approximate what 2000-era PA footprints would have contained under 2020 population levels, we estimate that roughly 51% of the increase is attributable to demographic growth around pre-existing PAs, and 49% to PA network expansion. The two mechanisms contribute in

approximately equal measure at the global LMIC level, though their relative importance varies substantially across countries.

3.3 Uncertainty from missing designation years

The WDPA records a designation year for each protected area, but this field is not reported for 2,912 of the 9,170 PAs in our 75 LMICs (31.8%; see Supplementary Materials, Section A). These PAs are known to exist, since they appear in the May 2021 WDPA release, but they cannot be unambiguously assigned to a specific designation period. Figure S2 (Supplementary Materials) shows, for each country, the gap between two counts: one restricted to PAs whose designation year is explicitly recorded as falling on or before 2020, and one that also includes PAs with missing designation year. The gap represents the uncertainty attributable to incomplete temporal metadata in the WDPA.

For 60 of the 75 countries, the gap is below 5 percentage points, indicating that the large majority of PAs have recorded designation years. However, in several countries – notably those where IUCN categories are missing for large protected areas – the gap exceeds 10 percentage points. For these countries, population figures are sensitive to how missing designation years are treated.

3.4 Cross-country variation

Figure 2 displays, for each country, the percentage of the national population residing inside PAs or within 10 km, in 2000 and 2020. The wide cross-country variation – from less than 1% to over 90% – underscores that aggregate statistics mask considerable heterogeneity. In 62 of the 75 countries, the share strictly increased between 2000 and 2020 (blue segments), consistent with PA expansion. Seven countries show no change over this comparison: Kiribati, North Korea, Somalia, and Syria have no protected areas in the WDPA extract, while in Micronesia, Moldova, and Palestine every protected area lacks a recorded designation year and is therefore excluded from the confirmed-year comparison shown here (their populations do appear in the 2020 all-PA figures). Six countries experienced marginal declines (all below 2.5 percentage points), most notably Eritrea, Zambia, and Malawi, where stable PA coverage combined with population growth in non-PA areas slightly reduced the relative share of PA-adjacent populations. Full country-level detail is provided in Table S1 (Supplementary Materials).




Figure 2: Share of the national population living inside protected areas or within 10 km of their boundaries, by country, in 2000 and 2020. Estimates are for 75 low- and lower-middle-income countries excluding India, using GHSL and restricted to protected areas with recorded designation year.

Figure 3 compares GHSL and WorldPop estimates of the percentage of national population inside PAs and within 10 km, for all PAs present in the 2020 WDPA release. Agreement is close for the broader 10 km perimeter, whereas the inside-PA panel shows a systematic tendency for WorldPop estimates to exceed GHSL.




Figure 3: Comparison of GHSL and WorldPop estimates of the share of national population living inside protected areas and inside or within 10 km of protected areas in 2020. Estimates are for 75 low- and lower-middle-income countries excluding India. The dashed line indicates 1:1 agreement. The inside-PA panel is systematically shifted upward relative to that line, whereas the broader 10 km perimeter is closer to parity.

For populations inside PAs, 64 of 75 country estimates lie above the 1:1 line, with a median WorldPop-to-GHSL ratio of 1.87 and a median difference of 0.77 percentage points. For the broader 10 km buffer perimeter, the asymmetry is weaker: 43 of 75 estimates lie above the line, the median ratio is 1.01, and the median difference is 0.2 percentage points. The likely reason is that the very small absolute population counts inside PAs amplify relative differences between the two datasets' spatial allocation models. Table S2 (Supplementary Materials) lists the countries where the absolute difference between GHSL and WorldPop exceeds 5 percentage points and the relative difference (computed as the absolute difference divided by the GHSL estimate) exceeds 10%. These discrepancies do not alter the order-of-magnitude conclusions: both datasets indicate hundreds of millions of people residing near protected areas in LMICs.

4 Discussion

4.1 Demographic scale and the impact evaluation literature

The central finding of this paper is that the populations encompassed by standard spatial exposure definitions in PA impact evaluations are large – on the order of several hundred million people.

Among PAs with recorded designation years, these populations are substantially larger in 2020 than in 2000, driven both by PA expansion itself and by population increase in areas where PAs were already present.

These population orders of magnitude are absent from the impact evaluation literature – whether in meta-analyses such as Kandel et al. (2022) or in large-scale assessments such as Naidoo et al. (2019) and Fisher et al. (2024) – yet they define the scope to which estimated impacts implicitly apply. When a study estimates the welfare effect of living within 10 km of a PA in a given country, and that effect is used to inform policy discussions about PA expansion, it is relevant to know that the population meeting this exposure criterion across LMICs numbers in the hundreds of millions - and that this population is growing.

This matters for at least three reasons. First, it sets a demographic frame for external validity: a treatment effect estimated on a sample of a few thousand households in a handful of countries implicitly speaks to a reference population that is orders of magnitude larger. Second, it reveals compositional heterogeneity: the “local population” implied by a 10 km buffer near a strict national park in a sparsely populated area is demographically very different from that implied by the same buffer near a densely inhabited community conservation area. Third, the heterogeneity of findings in the impact evaluation literature may itself partly reflect the heterogeneity of the populations being studied – populations that differ not only across countries but across PA management categories, as our decomposition shows.

4.2 PA categories matter for interpreting impacts

The decomposition by IUCN management category reveals that populations adjacent to non-strict PAs (IUCN IV-VI) exceed those adjacent to strict protection regimes (IUCN I-III). This pattern held in both 2000 and 2020.

This has direct relevance for interpreting impact evaluation results. The term “protected area” encompasses management regimes with vastly different implications for local populations – from national parks that prohibit entry to community conservation areas that allow resource extraction – yet comparisons of impacts across management categories remain rare (Kandel et al. 2022; Oldekop et al. 2016). Our results show that the demographic weight of non-strict PAs exceeds that of strictly protected areas in LMICs. This is mainly because non-strict PAs are more numerous and cover more land, though nearby population intensity per km² of PA is also somewhat higher around non-strict than around strict PAs. Part of the difference in populations *inside* PAs also reflects management rules: strict categories (IUCN I-III) are frequently designated as areas where human habitation and resource extraction are prohibited or tightly limited, whereas non-strict categories (IUCN IV-VI) explicitly accommodate settlement and sustainable use, so the near-absence of residents inside strict PAs is partly by design rather than an incidental settlement pattern. These functional differences between management regimes are not confined to human populations: in ecological studies for instance, habitat suitability for large fauna can be substantially improved by proximity to strictly protected areas but remain unaffected by less strictly protected areas where entry and extraction are permitted (Kittle et al. 2018). Studies that treat all PAs as a homogeneous treatment condition may therefore conflate very different exposure

regimes. PAs whose IUCN category is not recorded in the WDPA represent an additional source of ambiguity: for these areas, even the management regime is unknown.

Assessing what these different management regimes imply for local welfare is the domain of impact evaluation studies, not the purpose of this paper. What we document is that the population living near PAs is larger around non-strict protection regimes.

4.3 Data limitations

Our estimates rely on globally harmonized gridded population datasets. These datasets have known limitations. GHSL and WorldPop are constructed from census data combined with remote sensing and machine learning, and their accuracy varies across settings. Chen et al. (2020) evaluated four global gridded datasets and found that GHSL exhibited the closest correspondence to administrative census totals, while WorldPop showed larger deviations from census totals but high spatial consistency with other datasets. We use GHSL as our primary dataset and WorldPop as a robustness check. Figure 3 shows that the two datasets remain close for the broader 10 km zone, but that WorldPop tends to produce higher population shares inside PAs than GHSL. We cannot identify the precise cause from our data alone, but this pattern is consistent with documented differences in the products' spatial-allocation methods and in their treatment of sparse populations (Leyk et al. 2019; Thomson et al. 2022). This asymmetry does not alter the order-of-magnitude conclusion that protected areas in LMICs are adjacent to populations numbering in the hundreds of millions, even if point estimates for individual countries may differ.

The exclusion of India deserves further comment. India accounts for approximately 36% of the total LMIC population, and its actual PA network – 900 nationally designated sites, for a terrestrial coverage of approximately 7.7% (UNEP-WCMC 2026) – is substantial. However, India has restricted public access to these data in the WDPA, making only internationally designated sites (Ramsar wetlands, World Heritage sites) available, covering less than 1% of Indian territory. This data restriction has been documented: Wani et al. (2025) note that India has stated the process of making its data publicly accessible is in progress, and Zhang et al. (2025) acknowledge that the absence of publicly available Indian PA data may underestimate conservation extent in global analyses. Including India with incomplete spatial data would mechanically depress aggregate exposure statistics, producing artefactual rather than substantive results. We therefore exclude India entirely. This means our estimates do not cover the world's most populous lower-middle-income country, which is a significant limitation. If India's PA data were publicly available, the total number of people estimated to live near protected areas would increase substantially. At the same time, the share of the population concerned would likely decrease, given India's comparatively low protected area coverage.

The treatment of missing designation years introduces temporal ambiguity (see Appendix A). In our 2000-2020 comparison, we restrict attention to PAs with a recorded designation year, which means we are comparing a subset of PAs known to exist by 2000 with a different subset known to exist by 2020. This comparison is informative about orders of magnitude but should not be read as a precise estimate of temporal change, since some PAs with missing designation years may in fact have been established before 2000. Figure S2 (Supplementary Materials) makes the resulting uncertainty band visible at the country level.

4.4 Scope and interpretive boundaries

This paper quantifies population magnitudes. It does not estimate socio-economic impacts, welfare effects, or causal relationships. No claim is made - explicitly or implicitly - about whether living near a protected area makes people better or worse off. Such questions require impact evaluation designs with appropriate identification strategies, which are beyond the scope of this work.

We also do not rank conservation policies or PA categories in terms of desirability. The decomposition by management category is intended to describe the demographic composition of PA-adjacent populations, not to evaluate the relative merits of different protection regimes.

The paper does not claim that spatial proximity constitutes “being affected by” a protected area. The 10 km buffer is used because it is a common reference perimeter in the impact evaluation literature, not because it represents a known threshold for social or ecological effects. The actual spatial extent of PA-related effects on livelihoods is an empirical question that varies by context and cannot be resolved by our data.

A related limitation concerns population differentiation. Our estimates treat all inhabitants within a given perimeter as demographically equivalent. In practice, populations near protected areas are heterogeneous in their dependence on natural resources, their legal status, and their vulnerability to land-use restrictions. Indigenous peoples and traditional communities - estimated at 476 million people worldwide across more than 90 countries (United Nations Department of Economic and Social Affairs 2021) - are disproportionately represented among populations living within or adjacent to protected areas, and often bear the most direct consequences of conservation management (Schleicher et al. 2019). Our gridded population data cannot identify these groups. The aggregate population magnitudes we report therefore encompass communities with very different relationships to protected areas, from urban residents who may experience diffuse ecosystem service benefits to indigenous groups whose livelihoods and territorial rights may be directly affected by PA governance. Disaggregating these populations remains an important challenge for the field.

5 Conclusion

Across 75 low- and lower-middle-income countries, protected areas are adjacent to populations on the order of several hundred million people. Restricting to PA interiors captures a relatively small population; extending to 10 km increases the encompassed population roughly twelvefold. Among PAs with recorded designation years, these figures are substantially larger in 2020 than in 2000, reflecting both PA expansion and population increase – though incomplete temporal metadata in the WDPA means that the precise magnitude of this change cannot be established with certainty. Populations near non-strict PAs (IUCN IV-VI) exceed those near strict protection regimes (IUCN I-III). India, the largest LMIC by population, had to be excluded because its national PA data are absent from the WDPA; if included, global counts of people living near protected areas would likely increase in absolute terms, though the corresponding population share might decrease.

These facts are relevant for how conservation impact evaluation results are interpreted, compared, and generalized. They do not tell us what those impacts are. But they establish the demographic scale that spatial exposure definitions imply – a scale that is almost never reported in the studies that use them.

Bibliography

- Adams, William. M., Ros Aveling, Dan Brockington, et al. 2004. "Biodiversity Conservation and the Eradication of Poverty." *Science* 306 (5699): 1146–49. <https://doi.org/10.1126/science.1097920>.
- Avouac, Romain, Thomas Faria, and Frédéric Comte. 2025. "A Cloud-Native Data Science Platform for Official Statistics." In *Foundations and Advances of Machine Learning in Official Statistics*, edited by Florian Dumpert. Springer Nature Switzerland. https://doi.org/10.1007/978-3-032-10004-7_8.
- Baston, Daniel. 2023. *Exactextractr: Fast Extraction from Raster Datasets Using Polygons*. <https://isciences.gitlab.io/exactextractr/>.
- Bingham, Heather C., Diego Juffe Bignoli, Edward Lewis, et al. 2019. "Sixty Years of Tracking Conservation Progress Using the World Database on Protected Areas." *Nature Ecology & Evolution* 3 (5): 737743. https://idp.nature.com/authorize/casa?redirect_uri=https://www.nature.com/articles/s41559-019-0869-3&casa_token=Zmluq7JmhmUAAAAA:Id6cAJrCDpmtypiga8LmkB5wb9a1O_bwCGwCeA1gz314-F6VT5ekCOeVMJo_Zrpbh_uuoXf0fwD3Txjg.
- Byerlee, Derek, and Alain de Janvry, eds. 2008. *World Development Report 2008: Agriculture for Development*. World Bank.
- Castañeda, Andrés, Dung Doan, David Newhouse, Minh Cong Nguyen, Hiroki Uematsu, and João Pedro Azevedo. 2018. "A New Profile of the Global Poor." *World Development* 101 (January): 250–67. <https://doi.org/10.1016/j.worlddev.2017.08.002>.
- Chen, Ruxia, Huimin Yan, Fang Liu, Wenpeng Du, and Yanzhao Yang. 2020. "Multiple Global Population Datasets: Differences and Spatial Distribution Characteristics." *ISPRS International Journal of Geo-Information* 9 (11): 637. <https://doi.org/10.3390/ijgi9110637>.
- Coetzer, Kaera L., Edward T. F. Witkowski, and Barend F. N. Erasmus. 2014. "Reviewing Biosphere Reserves Globally: Effective Conservation Action or Bureaucratic Label?" *Biological Reviews* 89 (1): 82–104. <https://doi.org/10.1111/brv.12044>.
- Convention on Biological Diversity. 2022. "Kunming-Montreal Global Biodiversity Framework." *Convention Biol. Divers. Kunming-Montreal Glob. Biodivers. Framew.(cbd. Int)* CBD/COP/15/L.25. https://seea.un.org/sites/seea.un.org/files/unsc_presentation_jillian_campbell_0.pdf.
- Fisher, Joshua, Summer Allen, Greg Yetman, and Linda Pistoletti. 2024. "Assessing the Influence of Landscape Conservation and Protected Areas on Social Wellbeing Using Random Forest Machine Learning." *Scientific Reports* 14 (1): 11357. <https://doi.org/10.1038/s41598-024-61924-4>.
- Freire, Sergio, Kytt MacManus, Martino Pesaresi, Erin Doxsey-Whitfield, and Jane Mills. 2016. "Development of New Open and Free Multi-Temporal Global Population Grids at 250 m Resolution." *Population* 250: 33.
- Gorelick, Noel, Matt Hancher, Mike Dixon, Simon Ilyushchenko, David Thau, and Rebecca Moore. 2017. "Google Earth Engine: Planetary-Scale Geospatial Analysis for Everyone." *Remote Sensing of Environment* 202: 1827. <https://www.sciencedirect.com/science/article/pii/S0034425717302900>.

- Hanson, Jeffrey O. 2022. "Wdpar: Interface to the World Database on Protected Areas." *Journal of Open Source Software* 7 (78): 4594. <https://doi.org/10.21105/joss.04594>.
- Hijmans, Robert J. 2025. *Terra: Spatial Data Analysis*. <https://rspatial.org/>.
- Kandel, Pratikshya, Ram Pandit, Benedict White, and Maksym Polyakov. 2022. "Do Protected Areas Increase Household Income? Evidence from a Meta-Analysis." *World Development* 159 (November): 106024. <https://doi.org/10.1016/j.worlddev.2022.106024>.
- Kittle, Andrew M., Anjali C. Watson, Samuel A. Cushman, and David W. Macdonald. 2018. "Forest Cover and Level of Protection Influence the Island-Wide Distribution of an Apex Carnivore and Umbrella Species, the Sri Lankan Leopard (*Panthera pardus kotiya*)." *Biodiversity and Conservation* 27 (1): 235–63. <https://doi.org/10.1007/s10531-017-1431-8>.
- Leberger, Roxanne, Isabel M. D. Rosa, Carlos A. Guerra, Florian Wolf, and Henrique M. Pereira. 2020. "Global Patterns of Forest Loss Across IUCN Categories of Protected Areas." *Biological Conservation* 241 (January): 108299. <https://doi.org/10.1016/j.biocon.2019.108299>.
- Leyk, Stefan, Andrea E. Gaughan, Susana B. Adamo, et al. 2019. "The Spatial Allocation of Population: A Review of Large-Scale Gridded Population Data Products and Their Fitness for Use." *Earth System Science Data* 11 (3): 1385–409. <https://essd.copernicus.org/articles/11/1385/2019/essd-11-1385-2019.html>.
- Maxwell, Sean L., Victor Cazalis, Nigel Dudley, et al. 2020. "Area-Based Conservation in the Twenty-First Century." *Nature* 586 (7828): 217227.
- Naidoo, R., D. Gerkey, D. Hole, et al. 2019. "Evaluating the Impacts of Protected Areas on Human Well-Being Across the Developing World." *Science Advances* 5 (4): eaav3006. <https://doi.org/10.1126/sciadv.aav3006>.
- Oldekop, Johan A., George Holmes, W. Edwin Harris, and Karl L. Evans. 2016. "A Global Assessment of the Social and Conservation Outcomes of Protected Areas." *Conservation Biology* 30 (1): 133141.
- Pebesma, Edzer. 2018. "Simple Features for R: Standardized Support for Spatial Vector Data." *The R Journal* 10 (1): 439–46. <https://doi.org/10.32614/RJ-2018-009>.
- R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Runfola, Daniel, Austin Anderson, Heather Baier, et al. 2020. "geoBoundaries: A Global Database of Political Administrative Boundaries." *PLOS ONE* 15 (4): e0231866. <https://doi.org/10.1371/journal.pone.0231866>.
- Schleicher, Judith, Julie G. Zaehring, Carine Fastré, Bhaskar Vira, and Terry C. H. Sunderland. 2019. "Protecting Half of the Planet Could Directly Affect over One Billion People." *Nature Sustainability* 2: 1094–96. <https://doi.org/10.1038/s41893-019-0423-y>.
- Stevens, Forrest R., Andrea E. Gaughan, Catherine Linard, and Andrew J. Tatem. 2015. "Disaggregating Census Data for Population Mapping Using Random Forests with Remotely-Sensed and

Ancillary Data.” *PloS One* 10 (2): e0107042. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0107042>.

Thomson, Dana R, Douglas R Leasure, Tomas Bird, Nikos Tzavidis, and Andrew J Tatem. 2022. “How Accurate Are WorldPop-Global-Unconstrained Gridded Population Data at the Cell-Level?: A Simulation Analysis in Urban Namibia.” *Plos One* 17 (7): e0271504.

UNEP-WCMC. 2026. *Protected Area Profile for India*. <https://www.protectedplanet.net/country/IND>.

UNEP-WCMC and IUCN. 2021. *Protected Planet Report 2021*. UNEP-WCMC; IUCN.

UNEP-WCMC and IUCN. 2024. *Protected Planet Report 2024*. UNEP-WCMC; IUCN.

UNEP-WCMC, and IUCN. 2023. *Protected Planet: The World Database on Protected Areas (WDPA)*. UNEP-WCMC and IUCN. Cambridge, UK. www.protectedplanet.net.

United Nations Department of Economic and Social Affairs. 2021. *State of the World’s Indigenous Peoples*. 5th ed. United Nations. <https://www.un.org/development/desa/indigenouspeoples/publications/state-of-the-worlds-indigenous-peoples.html>.

Vaggi, Gianni. 2017. “The Rich and the Poor: A Note on Countries’ Classification.” *PSL Quarterly Review* 70 (279). https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3100742.

Wani, Zubair A., Lal Singh, Sanjay Uniyal, Sunil Kumar Rana, Indra D. Bhatt, and Sunil Nautiyal. 2025. “Improving Ecosystem Vitality in India: Overcoming Barriers to Meet National and International Targets.” *Environmental Sustainability* 8 (1): 17–29. <https://doi.org/10.1007/s42398-025-00339-x>.

West, Paige, James Igoe, and Dan Brockington. 2006. “Parks and Peoples: The Social Impact of Protected Areas.” *Annual Review of Anthropology* 35 (1): 251–77. <https://doi.org/10.1146/annurev.anthro.35.081705.123308>.

Zhang, Tong, Luke Gibson, Jianzhang Ma, Rachakonda Sreekar, and David Lindenmayer. 2025. “Global Patterns of Border Protected Areas Reveal Gaps in Transboundary Conservation Efforts.” *One Earth*, ahead of print. <https://doi.org/10.1016/j.oneear.2025.101206>.