

Name:
J#:
Date:

Dr. Clontz

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

Standard V1.	Mark:
---------------------	-------

Let V be the set of all pairs of real numbers with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \odot (x_1, y_1) = (0, cy_1)$$

(a) Show that scalar multiplication **distributes vectors** over scalar addition:

$$(c + d) \odot (x, y) = c \odot (x, y) \oplus d \odot (x, y).$$

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$, and let $c, d \in \mathbb{R}$. Then

$$(c + d) \odot (x_1, y_1) = (0, (c + d)y_1) = (0, cy_1) \oplus (0, dy_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

However, V is not a vector space, as $1 \odot (x_1, y_1) = (0, y_1) \neq (x_1, y_1)$.

□

Standard V3.	Mark:
---------------------	-------

Determine if the vectors $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$, and $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$ span \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span \mathbb{R}^4 .

□

Standard V4.	Mark:
---------------------	-------

Let W be the set of all polynomials of the form $ax^3 + bx$. Determine if W is a subspace of \mathcal{P}^3 .

Solution: Yes because $s(a_1x^3 + b_1x) + t(a_2x^3 + b_2x) = (sa_1 + ta_2)x^3 + (sb_1 + tb_2)x$ also belongs to W . Alternately, yes because W is isomorphic to \mathbb{R}^2 .

□

Standard S2.	Mark:
---------------------	-------

Determine if the set $\{x^2 + x - 1, 3x^2 - x + 1, 2x - 2\}$ is a basis of \mathcal{P}_2

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

Additional Notes/Marks	
-------------------------------	--