

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Module V: Vector Spaces

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

What is a vector space?

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

At the end of this module, students will be able to...

- V1. Vector spaces.** ... explain why a given set with defined addition and scalar multiplication does satisfy a given vector space property, but nonetheless isn't a vector space.
- V2. Linear combinations.** ... determine if a Euclidean vector can be written as a linear combination of a given set of Euclidean vectors by solving an appropriate vector equation.
- V3. Spanning sets.** ... determine if a set of Euclidean vectors spans \mathbb{R}^n by solving appropriate vector equations.
- V4. Subspaces.** ... determine if a subset of \mathbb{R}^n is a subspace or not.
- V5. Linear independence.** ... determine if a set of Euclidean vectors is linearly dependent or independent by solving an appropriate vector equation.
- V6. Basis verification.** ... explain why a set of Euclidean vectors is or is not a basis of \mathbb{R}^n .
- V7. Basis computation.** ... compute a basis for the subspace spanned by a given set of Euclidean vectors, and determine the dimension of the subspace.
- V8. Polynomial and Matrix computation.** ... answer questions about vector spaces of polynomials or matrices.
- V9. Basis of solution space.** ... find a basis for the solution set of a homogeneous system of equations.

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Readiness Assurance Outcomes

Before beginning this module, each student should be able to...

- Use set builder notation to describe sets of vectors.
 - Add Euclidean vectors and multiply Euclidean vectors by scalars.
 - Add polynomials and multiply polynomials by scalars.
 - Perform basic manipulations of augmented matrices and linear systems
- E1,E2,E3.**

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

The following resources will help you prepare for this module.

- Set Builder Notation: <https://youtu.be/xnfUZ-NTsCE>
- Adding and subtracting Euclidean vectors (Khan Academy):
<http://bit.ly/2y8A0wa>
- Linear combinations of Euclidean vectors (Khan Academy):
<http://bit.ly/2nK3wne>
- Adding and subtracting polynomials (Khan Academy):
<http://bit.ly/2d5SLGZ>

Module V Section 1

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Today's goals: Today we will explore what properties are shared by \mathbb{R}^1 and \mathbb{R}^2 with an eye towards generalizing to higher dimensions.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Observation V.1

Several properties of the real numbers, such as commutativity:

$$x + y = y + x$$

also hold for Euclidean vectors with multiple components:

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} + \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.2 (~ 20 min) Consider each of the following properties of the real numbers \mathbb{R}^1 . Label each property as **valid** if the property also holds for two-dimensional Euclidean vectors $\vec{u}, \vec{v}, \vec{w} \in \mathbb{R}^2$ and scalars $a, b \in \mathbb{R}$, and **invalid** if it does not.

1. $\vec{u} + (\vec{v} + \vec{w}) = (\vec{u} + \vec{v}) + \vec{w}$.

2. $\vec{u} + \vec{v} = \vec{v} + \vec{u}$.

3. There exists some \vec{z} where $\vec{v} + \vec{z} = \vec{v}$.

4. There exists some $-\vec{v}$ where
 $\vec{v} + (-\vec{v}) = \vec{z}$.

5. If $\vec{u} \neq \vec{v}$, then $\frac{1}{2}(\vec{u} + \vec{v})$ is the only
 vector equally distant from both \vec{u} and
 \vec{v}

6. $a(b\vec{v}) = (ab)\vec{v}$.

7. $1\vec{v} = \vec{v}$.

8. If $\vec{u} \neq \vec{0}$, then there exists some scalar
 c such that $c\vec{u} = \vec{v}$.

9. $a(\vec{u} + \vec{v}) = a\vec{u} + a\vec{v}$.

10. $(a + b)\vec{v} = a\vec{v} + b\vec{v}$.

Definition V.3

A **vector space** V is any collection of mathematical objects with associated addition \oplus and scalar multiplication \odot operations that satisfy the following properties. Let $\vec{u}, \vec{v}, \vec{w}$ belong to V , and let a, b be scalar numbers.

- **Addition is associative:**

$$\vec{u} \oplus (\vec{v} \oplus \vec{w}) = (\vec{u} \oplus \vec{v}) \oplus \vec{w}.$$
- **Addition is commutative:**

$$\vec{u} \oplus \vec{v} = \vec{v} \oplus \vec{u}.$$
- **Additive identity exists:** There exists some \vec{z} where $\vec{v} \oplus \vec{z} = \vec{v}$.
- **Additive inverses exist:** There exists some $-\vec{v}$ where $\vec{v} \oplus (-\vec{v}) = \vec{z}$.
- **Scalar multiplication is associative:**

$$a \odot (b \odot \vec{v}) = (ab) \odot \vec{v}.$$
- **Scalar multiplication identity exists:** $1 \odot \vec{v} = \vec{v}$.
- **Scalar mult. distributes over vector addition:**

$$a \odot (\vec{u} \oplus \vec{v}) = a \odot \vec{u} \oplus a \odot \vec{v}.$$
- **Scalar mult. distributes over scalar addition:**

$$(a + b) \odot \vec{v} = a \odot \vec{v} \oplus b \odot \vec{v}.$$

Observation V.4

Every **Euclidean vector space**

$$\mathbb{R}^n = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \mid x_1, x_2, \dots, x_n \in \mathbb{R} \right\}$$

satisfies all eight requirements for the usual definitions of addition and scalar multiplication, but we will also study other types of vector spaces.

Observation V.5

The space of $m \times n$ matrices

$$M_{m,n} = \left\{ \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \mid a_{11}, \dots, a_{mn} \in \mathbb{R} \right\}$$

satisfies all eight requirements for component-wise addition and scalar multiplication.

Remark V.6

Every Euclidean space \mathbb{R}^n is a vector space, but there are other examples of vector spaces as well.

For example, consider the set \mathbb{C} of complex numbers with the usual definitions of addition and scalar multiplication, and let $\vec{u} = a + bi$, $\vec{v} = c + di$, and $\vec{w} = e + fi$. Then

$$\begin{aligned}\vec{u} + (\vec{v} + \vec{w}) &= (a + bi) + ((c + di) + (e + fi)) \\ &= (a + bi) + ((c + e) + (d + f)i) \\ &= (a + c + e) + (b + d + f)i \\ &= ((a + c) + (b + d)i) + (e + fi) \\ &= (\vec{u} + \vec{v}) + \vec{w}\end{aligned}$$

All eight properties can be verified in this way.

Remark V.7

The following sets are just a few examples of vector spaces, with the usual/natural operations for addition and scalar multiplication.

- \mathbb{R}^n : Euclidean vectors with n components.
- \mathbb{C} : Complex numbers.
- $M_{m,n}$: Matrices of real numbers with m rows and n columns.
- \mathcal{P}^n : Polynomials of degree n or less.
- \mathcal{P} : Polynomials of any degree.
- $C(\mathbb{R})$: Real-valued continuous functions.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Summary: Today we came up with a set of properties shared by all \mathbb{R}^n and used these to define **vector spaces** (Standard V1). Next class, we will practice determining when other sets are or are not vector spaces.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 2

Module V

- Section 1
- Section 2**
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Today's goals: Today we will practice showing when a set with given operations is or is not a vector space (Standard V1).

Remark V.8

Previously, we defined a **vector space** V to be any collection of mathematical objects with associated addition and scalar multiplication operations that satisfy the following eight properties for all $\vec{u}, \vec{v}, \vec{w}$ in V , and all scalars (i.e. real numbers) a, b .

- **Addition is associative:**

$$\vec{u} \oplus (\vec{v} \oplus \vec{w}) = (\vec{u} \oplus \vec{v}) \oplus \vec{w}.$$
- **Addition is commutative:**

$$\vec{u} \oplus \vec{v} = \vec{v} \oplus \vec{u}.$$
- **Additive identity exists:** There exists some \vec{z} where $\vec{v} \oplus \vec{z} = \vec{v}$.
- **Additive inverses exist:** There exists some $-\vec{v}$ where $\vec{v} \oplus (-\vec{v}) = \vec{z}$.
- **Scalar multiplication is associative:**

$$a \odot (b \odot \vec{v}) = (ab) \odot \vec{v}.$$
- **Scalar multiplication identity exists:** $1 \odot \vec{v} = \vec{v}$.
- **Scalar mult. distributes over vector addition:**

$$a \odot (\vec{u} \oplus \vec{v}) = a \odot \vec{u} \oplus a \odot \vec{v}.$$
- **Scalar mult. distributes over scalar addition:**

$$(a + b) \odot \vec{v} = a \odot \vec{v} \oplus b \odot \vec{v}.$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.9 (*~20 min*) Consider the set $V = \{(x, y) \mid y = e^x\}$ with operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 y_2) \qquad c \odot (x_1, y_1) = (cx_1, y_1^c)$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.9 (~ 20 min) Consider the set $V = \{(x, y) \mid y = e^x\}$ with operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 y_2) \qquad c \odot (x_1, y_1) = (cx_1, y_1^c)$$

Part 1: Show that V satisfies the distributive property

$$(a + b) \odot (x_1, y_1) = (a \odot (x_1, y_1)) \oplus (b \odot (x_1, y_1))$$

by simplifying both sides and verifying they are the same expression.

Activity V.9 (~ 20 min) Consider the set $V = \{(x, y) \mid y = e^x\}$ with operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 y_2) \qquad c \odot (x_1, y_1) = (cx_1, y_1^c)$$

Part 1: Show that V satisfies the distributive property

$$(a + b) \odot (x_1, y_1) = (a \odot (x_1, y_1)) \oplus (b \odot (x_1, y_1))$$

by simplifying both sides and verifying they are the same expression.

Part 2: Show that V contains an additive identity element satisfying

$$(x_1, y_1) \oplus \vec{z} = (x_1, y_1)$$

for all $(x_1, y_1) \in V$ by choosing appropriate values for $\vec{z} = (?, ?)$.

Remark V.10

It turns out $V = \{(x, y) \mid y = e^x\}$ with operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 y_2) \qquad c \odot (x_1, y_1) = (cx_1, y_1^c)$$

satisfies all eight properties.

- **Addition is associative:**

$$\vec{u} \oplus (\vec{v} \oplus \vec{w}) = (\vec{u} \oplus \vec{v}) \oplus \vec{w}.$$

- **Addition is commutative:**

$$\vec{u} \oplus \vec{v} = \vec{v} \oplus \vec{u}.$$

- **Additive identity exists:** There exists some \vec{z} where $\vec{v} \oplus \vec{z} = \vec{v}$.

- **Additive inverses exist:** There exists some $-\vec{v}$ where $\vec{v} \oplus (-\vec{v}) = \vec{z}$.

- **Scalar multiplication is associative:**

$$a \odot (b \odot \vec{v}) = (ab) \odot \vec{v}.$$

- **Scalar multiplication identity exists:** $1 \odot \vec{v} = \vec{v}$.

- **Scalar mult. distributes over vector addition:**

$$a \odot (\vec{u} \oplus \vec{v}) = a \odot \vec{u} \oplus a \odot \vec{v}.$$

- **Scalar mult. distributes over scalar addition:**

$$(a + b) \odot \vec{v} = a \odot \vec{v} \oplus b \odot \vec{v}.$$

Thus, V is a vector space.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.11 (*~15 min*) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + y_1 + x_2 + y_2, x_1^2 + x_2^2) \qquad c \odot (x_1, y_1) = (x_1^c, y_1 + c - 1).$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.11 (*~15 min*) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + y_1 + x_2 + y_2, x_1^2 + x_2^2) \quad c \odot (x_1, y_1) = (x_1^c, y_1 + c - 1).$$

Part 1: Show that 1 is the scalar multiplication identity element by simplifying $1 \odot (x, y)$ to (x, y) .

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.11 (*~15 min*) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + y_1 + x_2 + y_2, x_1^2 + x_2^2) \quad c \odot (x_1, y_1) = (x_1^c, y_1 + c - 1).$$

Part 1: Show that 1 is the scalar multiplication identity element by simplifying $1 \odot (x, y)$ to (x, y) .

Part 2: Show that V does not have an additive identity element by showing that $(0, -1) \oplus \vec{z} \neq (0, -1)$ no matter how $\vec{z} = (z, w)$ is chosen.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.11 (*~15 min*) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + y_1 + x_2 + y_2, x_1^2 + x_2^2) \quad c \odot (x_1, y_1) = (x_1^c, y_1 + c - 1).$$

Part 1: Show that 1 is the scalar multiplication identity element by simplifying $1 \odot (x, y)$ to (x, y) .

Part 2: Show that V does not have an additive identity element by showing that $(0, -1) \oplus \vec{z} \neq (0, -1)$ no matter how $\vec{z} = (z, w)$ is chosen.

Part 3: Is V a vector space?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.12 (*~15 min*) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + 3y_2) \qquad c \odot (x_1, y_1) = (cx_1, cy_1).$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.12 (~ 15 min) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + 3y_2) \quad c \odot (x_1, y_1) = (cx_1, cy_1).$$

Part 1: Show that scalar multiplication distributes over vector addition, i.e.

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)$$

for **all** $c \in \mathbb{R}$, $(x_1, y_1), (x_2, y_2) \in V$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.12 (~ 15 min) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + 3y_2) \quad c \odot (x_1, y_1) = (cx_1, cy_1).$$

Part 1: Show that scalar multiplication distributes over vector addition, i.e.

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)$$

for **all** $c \in \mathbb{R}$, $(x_1, y_1), (x_2, y_2) \in V$.

Part 2: Show that vector addition is not associative, i.e.

$$(x_1, y_1) \oplus ((x_2, y_2) \oplus (x_3, y_3)) \neq ((x_1, y_1) \oplus (x_2, y_2)) \oplus (x_3, y_3)$$

for **some** vectors $(x_1, y_1), (x_2, y_2), (x_3, y_3) \in V$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.12 (~ 15 min) Let $V = \{(x, y) \mid x, y \in \mathbb{R}\}$ have operations defined by

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + 3y_2) \quad c \odot (x_1, y_1) = (cx_1, cy_1).$$

Part 1: Show that scalar multiplication distributes over vector addition, i.e.

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)$$

for **all** $c \in \mathbb{R}$, $(x_1, y_1), (x_2, y_2) \in V$.

Part 2: Show that vector addition is not associative, i.e.

$$(x_1, y_1) \oplus ((x_2, y_2) \oplus (x_3, y_3)) \neq ((x_1, y_1) \oplus (x_2, y_2)) \oplus (x_3, y_3)$$

for **some** vectors $(x_1, y_1), (x_2, y_2), (x_3, y_3) \in V$.

Part 3: Is V a vector space?

Module V

- Section 1
- Section 2**
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Summary: Today we practiced showing when a set with given operations is or is not a vector space (Standard V1).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 3

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Today's goals: Today we will begin exploring the notions of **linear combinations** and **span**.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Definition V.13

A **linear combination** of a set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m\}$ is given by $c_1\vec{v}_1 + c_2\vec{v}_2 + \dots + c_m\vec{v}_m$ for any choice of scalar multiples c_1, c_2, \dots, c_m .

For example, we can say $\begin{bmatrix} 3 \\ 0 \\ 5 \end{bmatrix}$ is a linear combination of the vectors $\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$

since

$$\begin{bmatrix} 3 \\ 0 \\ 5 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

Definition V.14

The **span** of a set of vectors is the collection of all linear combinations of that set:

$$\text{span}\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m\} = \{c_1\vec{v}_1 + c_2\vec{v}_2 + \cdots + c_m\vec{v}_m \mid c_i \in \mathbb{R}\}.$$

For example:

$$\text{span}\left\{\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}\right\} = \left\{a \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} + b \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \mid a, b \in \mathbb{R}\right\}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.15 (*~10 min*) Consider $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.15 (*~10 min*) Consider $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$.

Part 1: Sketch

$$1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix},$$

$$3 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 6 \end{bmatrix},$$

$$0 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

$$\text{and } -2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -2 \\ -4 \end{bmatrix}$$

in the xy plane.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.15 (*~10 min*) Consider $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$.

Part 1: Sketch

$$1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad 3 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}, \quad 0 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \text{and } -2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -2 \\ -4 \end{bmatrix}$$

in the xy plane.

Part 2: Sketch a representation of all the vectors belonging to

$\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\} = \left\{ a \begin{bmatrix} 1 \\ 2 \end{bmatrix} \mid a \in \mathbb{R} \right\}$ in the xy plane.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.16 (*~10 min*) Consider $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.16 (*~10 min*) Consider $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$.

Part 1: Sketch the following linear combinations in the xy plane.

$$1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \qquad 0 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \qquad 1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$-2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \qquad -1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + -2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.16 (*~10 min*) Consider $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$.

Part 1: Sketch the following linear combinations in the xy plane.

$$1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \qquad 0 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \qquad 1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$-2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \qquad -1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + -2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Part 2: Sketch a representation of all the vectors belonging to $\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ in the xy plane.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.17 (~ 5 min) Sketch a representation of all the vectors belonging to $\text{span} \left\{ \begin{bmatrix} 6 \\ -4 \end{bmatrix}, \begin{bmatrix} -3 \\ 2 \end{bmatrix} \right\}$ in the xy plane.

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Summary: Today we begin exploring **linear combinations** and **span**.

Next class, we will learn how to check if a vector is a linear combination of a set of vectors or not (Standard V2).

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Module V Section 4

Module V

- Section 1
- Section 2
- Section 3
- Section 4**
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Today's goals: Today we will learn how to check if a vector is a linear combination of a set of vectors or not (Standard V2).

Remark V.18

Recall these definitions from last class:

- A **linear combination** of vectors is given by adding scalar multiples of those vectors, such as:

$$\begin{bmatrix} 3 \\ 0 \\ 5 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

- The **span** of a set of vectors is the collection of all linear combinations of that set, such as:

$$\text{span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \right\} = \left\{ a \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} + b \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.19 (*~15 min*) The vector $\begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}$ belongs to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} \right\}$ exactly when there exists a solution to the vector equation

$$x_1 \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}.$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.19 (*~15 min*) The vector $\begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}$ belongs to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} \right\}$ exactly when there exists a solution to the vector equation

$$x_1 \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}.$$

Part 1: Reinterpret this vector equation as a system of linear equations.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.19 (*~15 min*) The vector $\begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}$ belongs to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} \right\}$ exactly when there exists a solution to the vector equation

$$x_1 \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}.$$

Part 1: Reinterpret this vector equation as a system of linear equations.

Part 2: Find its solution set, using technology to find RREF of its corresponding augmented matrix.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.19 (*~15 min*) The vector $\begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}$ belongs to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} \right\}$

exactly when there exists a solution to the vector equation

$$x_1 \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}.$$

Part 1: Reinterpret this vector equation as a system of linear equations.

Part 2: Find its solution set, using technology to find RREF of its corresponding augmented matrix.

Part 3: Given this solution set, does $\begin{bmatrix} -1 \\ -6 \\ 1 \end{bmatrix}$ belong to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} \right\}$?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Fact V.20

A vector \vec{b} belongs to $\text{span}\{\vec{v}_1, \dots, \vec{v}_n\}$ if and only if the vector equation $x_1\vec{v}_1 + \dots + x_n\vec{v}_n = \vec{b}$ is consistent.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Quick Check V.21

The following are all equivalent statements:

- The vector \vec{b} belongs to $\text{span}\{\vec{v}_1, \dots, \vec{v}_n\}$.
- The vector equation $x_1\vec{v}_1 + \dots + x_n\vec{v}_n = \vec{b}$ is consistent.
- The linear system corresponding to $[\vec{v}_1 \dots \vec{v}_n \mid \vec{b}]$ is consistent.
- $\text{RREF}[\vec{v}_1 \dots \vec{v}_n \mid \vec{b}]$ doesn't have a row $[0 \dots 0 \mid 1]$ representing the contradiction $0 = 1$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.22 (*~10 min*) Determine if $\begin{bmatrix} 3 \\ -2 \\ 1 \\ 5 \end{bmatrix}$ belongs to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \\ 2 \end{bmatrix} \right\}$
by solving an appropriate vector equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.23 (*~5 min*) Determine if $\begin{bmatrix} -1 \\ -9 \\ 0 \end{bmatrix}$ belongs to $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix} \right\}$ by solving an appropriate vector equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.24 (*~10 min*) Does the third-degree polynomial $3y^3 - 2y^2 + y + 5$ in \mathcal{P}^3 belong to $\text{span}\{y^3 - 3y + 2, -y^3 - 3y^2 + 2y + 2\}$?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.24 (~ 10 min) Does the third-degree polynomial $3y^3 - 2y^2 + y + 5$ in \mathcal{P}^3 belong to $\text{span}\{y^3 - 3y + 2, -y^3 - 3y^2 + 2y + 2\}$?

Part 1: Reinterpret this question as a question about the solution(s) of a polynomial equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.24 (~ 10 min) Does the third-degree polynomial $3y^3 - 2y^2 + y + 5$ in \mathcal{P}^3 belong to $\text{span}\{y^3 - 3y + 2, -y^3 - 3y^2 + 2y + 2\}$?

Part 1: Reinterpret this question as a question about the solution(s) of a polynomial equation.

Part 2: Answer this equivalent question, and use its solution to answer the original question.

Module V

- Section 1
- Section 2
- Section 3
- Section 4**
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Activity V.25 (*~5 min*) Does the polynomial $x^2 + x + 1$ belong to $\text{span}\{x^2 - x, x + 1, x^2 - 1\}$?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.26 (*~5 min*) Does the matrix $\begin{bmatrix} 3 & -2 \\ 1 & 5 \end{bmatrix}$ belong to $\text{span} \left\{ \begin{bmatrix} 1 & 0 \\ -3 & 2 \end{bmatrix}, \begin{bmatrix} -1 & -3 \\ 2 & 2 \end{bmatrix} \right\}$?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.26 (~ 5 min) Does the matrix $\begin{bmatrix} 3 & -2 \\ 1 & 5 \end{bmatrix}$ belong to

$\text{span} \left\{ \begin{bmatrix} 1 & 0 \\ -3 & 2 \end{bmatrix}, \begin{bmatrix} -1 & -3 \\ 2 & 2 \end{bmatrix} \right\}?$

Part 1: Reinterpret this question as a question about the solution(s) of a matrix equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.26 (~ 5 min) Does the matrix $\begin{bmatrix} 3 & -2 \\ 1 & 5 \end{bmatrix}$ belong to

$$\text{span} \left\{ \begin{bmatrix} 1 & 0 \\ -3 & 2 \end{bmatrix}, \begin{bmatrix} -1 & -3 \\ 2 & 2 \end{bmatrix} \right\}?$$

Part 1: Reinterpret this question as a question about the solution(s) of a matrix equation.

Part 2: Answer this equivalent question, and use its solution to answer the original question.

Module V

- Section 1
- Section 2
- Section 3
- Section 4**
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Summary: Today we learned how to check if a vector is a linear combination of a set of vectors or not (Standard V2).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 5

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5**
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Today's goals: Today we will learn how to determine when a set of vectors spans the entire vector space (Standard V3).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8


Section 9

Section 10

Section 11

Observation V.27

Any single non-zero vector/number x in \mathbb{R}^1 spans \mathbb{R}^1 , since $\mathbb{R}^1 = \{cx \mid c \in \mathbb{R}\}$.



Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7


Section 8

Section 9

Section 10

Section 11

Activity V.28 (~ 5 min) How many vectors are required to span \mathbb{R}^2 ? Sketch a drawing in the xy plane to support your answer.



- (a) 1
- (b) 2
- (c) 3
- (d) 4
- (e) Infinitely Many

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7


Section 8

Section 9

Section 10

Section 11

Activity V.29 (*~5 min*) How many vectors are required to span \mathbb{R}^3 ?



- (a) 1
- (b) 2
- (c) 3
- (d) 4
- (e) Infinitely Many

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8


Section 9

Section 10

Section 11

Fact V.30

At least n vectors are required to span \mathbb{R}^n .



Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.31 (*~15 min*) Choose any vector $\begin{bmatrix} ? \\ ? \\ ? \end{bmatrix}$ in \mathbb{R}^3 that is not in

$\text{span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix} \right\}$ by using technology to verify that

$$\text{RREF} \left[\begin{array}{cc|c} 1 & -2 & ? \\ -1 & 0 & ? \\ 0 & 1 & ? \end{array} \right] = \left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right]. \text{ (Why does this work?)}$$

Fact V.32

The set $\{\vec{v}_1, \dots, \vec{v}_m\}$ fails to span all of \mathbb{R}^n exactly when the vector equation

$$x_1 \vec{v}_1 + \cdots + x_m \vec{v}_m = \vec{w}$$

is inconsistent for **some** vector \vec{w} .

Note that this happens exactly when $\text{RREF}[\vec{v}_1 \ \dots \ \vec{v}_m]$ has a non-pivot row of zeros.

$$\begin{bmatrix} 1 & -2 \\ -1 & 0 \\ 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\Rightarrow \left[\begin{array}{cc|c} 1 & -2 & a \\ -1 & 0 & b \\ 0 & 1 & c \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] \text{ for some choice of vector } \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.33 (*~5 min*) Consider the set of vectors

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -4 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 7 \\ -3 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \\ 5 \\ 7 \end{bmatrix}, \begin{bmatrix} 3 \\ 13 \\ 7 \\ 16 \end{bmatrix} \right\}. \text{ Does } \mathbb{R}^4 = \text{span } S?$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.33 (~ 5 min) Consider the set of vectors

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -4 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 7 \\ -3 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \\ 5 \\ 7 \end{bmatrix}, \begin{bmatrix} 3 \\ 13 \\ 7 \\ 16 \end{bmatrix} \right\}. \text{ Does } \mathbb{R}^4 = \text{span } S?$$

Part 1: Rewrite this as a question about the solutions to a vector equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.33 (~ 5 min) Consider the set of vectors

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -4 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 7 \\ -3 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \\ 5 \\ 7 \end{bmatrix}, \begin{bmatrix} 3 \\ 13 \\ 7 \\ 16 \end{bmatrix} \right\}. \text{ Does } \mathbb{R}^4 = \text{span } S?$$

Part 1: Rewrite this as a question about the solutions to a vector equation.

Part 2: Answer your new question, and use this to answer the original question.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.34 (*~10 min*) Consider the set of third-degree polynomials

$$S = \{2x^3 + 3x^2 - 1, 2x^3 + 3, 3x^3 + 13x^2 + 7x + 16, \\ -x^3 + 10x^2 + 7x + 14, 4x^3 + 3x^2 + 2\}.$$

Does $\mathcal{P}^3 = \text{span } S$?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.34 (*~10 min*) Consider the set of third-degree polynomials

$$S = \{2x^3 + 3x^2 - 1, 2x^3 + 3, 3x^3 + 13x^2 + 7x + 16, \\ -x^3 + 10x^2 + 7x + 14, 4x^3 + 3x^2 + 2\}.$$

Does $\mathcal{P}^3 = \text{span } S$?

Part 1: Rewrite this as a question about the solutions to a polynomial equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.34 (*~10 min*) Consider the set of third-degree polynomials

$$S = \{2x^3 + 3x^2 - 1, 2x^3 + 3, 3x^3 + 13x^2 + 7x + 16, \\ -x^3 + 10x^2 + 7x + 14, 4x^3 + 3x^2 + 2\}.$$

Does $\mathcal{P}^3 = \text{span } S$?

Part 1: Rewrite this as a question about the solutions to a polynomial equation.

Part 2: Answer your new question, and use this to answer the original question.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.35 (*~5 min*) Consider the set of matrices

$$S = \left\{ \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \right\}$$

Does $M_{2,2} = \text{span } S$?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.35 (*~5 min*) Consider the set of matrices

$$S = \left\{ \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \right\}$$

Does $M_{2,2} = \text{span } S$?

Part 1: Rewrite this as a question about the solutions to a matrix equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.35 (~ 5 min) Consider the set of matrices

$$S = \left\{ \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \right\}$$

Does $M_{2,2} = \text{span } S$?

Part 1: Rewrite this as a question about the solutions to a matrix equation.

Part 2: Answer your new question, and use this to answer the original question.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.36 (~ 5 min) Let $\vec{v}_1, \vec{v}_2, \vec{v}_3 \in \mathbb{R}^7$ be three vectors, and suppose \vec{w} is another vector with $\vec{w} \in \text{span}\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$. What can you conclude about $\text{span}\{\vec{w}, \vec{v}_1, \vec{v}_2, \vec{v}_3\}$?

- (a) $\text{span}\{\vec{w}, \vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is larger than $\text{span}\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$.
- (b) $\text{span}\{\vec{w}, \vec{v}_1, \vec{v}_2, \vec{v}_3\} = \text{span}\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$.
- (c) $\text{span}\{\vec{w}, \vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is smaller than $\text{span}\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Summary: Today we learned how to determine when a set of vectors spans the entire vector space (Standard V3).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 6

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10


Section 11

Today's goals: Today we will learn about **subspaces** (Standard V4)

Definition V.37

A subset of a vector space is called a **subspace** if it is a vector space on its own.

For example, the span of these two vectors forms a planar subspace inside of the larger vector space \mathbb{R}^3 .



Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Fact V.38

Any **subset** S of a vector space V that contains the additive identity $\vec{0}$ satisfies the eight vector space properties automatically, since it is a collection of known vectors.

However, to verify that it's a **subspace**, we need to check that addition and multiplication still make sense using only vectors from S . So we need to check two things:

- The set is **closed under addition**: for any $\vec{x}, \vec{y} \in S$, the sum $\vec{x} + \vec{y}$ is also in S .
- The set is **closed under scalar multiplication**: for any $\vec{x} \in S$ and scalar $c \in \mathbb{R}$, the product $c\vec{x}$ is also in S .

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.39 (*~15 min*) Let $S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + 2y + z = 0 \right\}.$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.39 (~ 15 min) Let $S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + 2y + z = 0 \right\}$.

Part 1: Let $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $\vec{w} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$ be vectors in S , so $x + 2y + z = 0$ and

$a + 2b + c = 0$. Show that $\vec{v} + \vec{w} = \begin{bmatrix} x + a \\ y + b \\ z + c \end{bmatrix}$ also belongs to S by verifying that $(x + a) + 2(y + b) + (z + c) = 0$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.39 (~ 15 min) Let $S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + 2y + z = 0 \right\}$.

Part 1: Let $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $\vec{w} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$ be vectors in S , so $x + 2y + z = 0$ and

$a + 2b + c = 0$. Show that $\vec{v} + \vec{w} = \begin{bmatrix} x + a \\ y + b \\ z + c \end{bmatrix}$ also belongs to S by verifying that $(x + a) + 2(y + b) + (z + c) = 0$.

Part 2: Let $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \in S$, so $x + 2y + z = 0$. Show that $c\vec{v} = \begin{bmatrix} cx \\ cy \\ cz \end{bmatrix}$ also belongs to S for any $c \in \mathbb{R}$ by verifying an appropriate equation.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.39 (~ 15 min) Let $S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + 2y + z = 0 \right\}$.

Part 1: Let $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $\vec{w} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$ be vectors in S , so $x + 2y + z = 0$ and

$a + 2b + c = 0$. Show that $\vec{v} + \vec{w} = \begin{bmatrix} x + a \\ y + b \\ z + c \end{bmatrix}$ also belongs to S by verifying that $(x + a) + 2(y + b) + (z + c) = 0$.

Part 2: Let $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \in S$, so $x + 2y + z = 0$. Show that $c\vec{v} = \begin{bmatrix} cx \\ cy \\ cz \end{bmatrix}$ also belongs to S for any $c \in \mathbb{R}$ by verifying an appropriate equation.

Part 3: Is S a subspace of \mathbb{R}^3 ?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.40 (*~10 min*) Let $S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + 2y + z = 4 \right\}$. Choose a vector

$\vec{v} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix}$ in S and a real number $c = ?$, and show that $c\vec{v}$ isn't in S . Is S a subspace of \mathbb{R}^3 ?

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Remark V.41

Since 0 is a scalar and $0\vec{v} = \vec{z}$ for any vector \vec{v} , a nonempty set that is closed under scalar multiplication must contain the zero vector \vec{z} for that vector space.

Put another way, you can check any of the following to show that a nonempty subset W isn't a subspace:

- Show that $\vec{0} \notin W$.
- Find $\vec{u}, \vec{v} \in W$ such that $\vec{u} + \vec{v} \notin W$.
- Find $c \in \mathbb{R}, \vec{v} \in W$ such that $c\vec{v} \notin W$.

If you cannot do any of these, then W can be proven to be a subspace by doing the following:

- Prove that $\vec{u} + \vec{v} \in W$ whenever $\vec{u}, \vec{v} \in W$.
- Prove that $c\vec{v} \in W$ whenever $c \in \mathbb{R}, \vec{v} \in W$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.42 (*~20 min*) Consider these subsets of \mathbb{R}^3 :

$$R = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = z + 1 \right\} \quad S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = |z| \right\} \quad T = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid z = xy \right\}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.42 (*~20 min*) Consider these subsets of \mathbb{R}^3 :

$$R = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = z + 1 \right\} \quad S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = |z| \right\} \quad T = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid z = xy \right\}$$

Part 1: Show R isn't a subspace by showing that $\vec{0} \notin R$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.42 (*~20 min*) Consider these subsets of \mathbb{R}^3 :

$$R = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = z + 1 \right\} \quad S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = |z| \right\} \quad T = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid z = xy \right\}$$

Part 1: Show R isn't a subspace by showing that $\vec{0} \notin R$.

Part 2: Show S isn't a subspace by finding two vectors $\vec{u}, \vec{v} \in S$ such that $\vec{u} + \vec{v} \notin S$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.42 (~ 20 min) Consider these subsets of \mathbb{R}^3 :

$$R = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = z + 1 \right\} \quad S = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid y = |z| \right\} \quad T = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid z = xy \right\}$$

Part 1: Show R isn't a subspace by showing that $\vec{0} \notin R$.

Part 2: Show S isn't a subspace by finding two vectors $\vec{u}, \vec{v} \in S$ such that $\vec{u} + \vec{v} \notin S$.

Part 3: Show T isn't a subspace by finding a vector $\vec{v} \in T$ such that $2\vec{v} \notin T$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.43 (*~5 min*) Let W be a subspace of a vector space V . How are $\text{span } W$ and W related?

- (a) $\text{span } W$ is bigger than W
- (b) $\text{span } W$ is the same as W
- (c) $\text{span } W$ is smaller than W

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6**
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Fact V.44

If S is any subset of a vector space V , then since $\text{span } S$ collects all possible linear combinations, $\text{span } S$ is automatically a subspace of V .

In fact, $\text{span } S$ is always the smallest subspace of V that contains all the vectors in S .

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6**
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Summary: Today we learned how to determine if a set of vectors forms a subspace of \mathbb{R}^n (Standard V4).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 7

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7**
- Section 8
- Section 9
- Section 10
- Section 11

Today's goals: Today we will learn about an important concept called **linear independence** (Standard V5).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.45 (~ 10 min) Consider the two sets

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix} \right\}$$


$$T = \left\{ \begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ -11 \end{bmatrix} \right\}$$

Which of the following is true?

- (A) $\text{span } S$ is bigger than $\text{span } T$.
- (B) $\text{span } S$ and $\text{span } T$ are the same size.
- (C) $\text{span } S$ is smaller than $\text{span } T$.

Definition V.46

We say that a set of vectors is **linearly dependent** if one vector in the set belongs to the span of the others. Otherwise, we say the set is **linearly independent**.



You can think of linearly dependent sets as containing a redundant vector, in the sense that you can drop a vector out without reducing the span of the set. In the above image, all three vectors lay on the same planar subspace, but only two vectors are needed to span the plane, so the set is linearly dependent.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.47 (~ 10 min) Let $\vec{v}_1, \vec{v}_2, \vec{v}_3$ be vectors in \mathbb{R}^n . Suppose $3\vec{v}_1 - 5\vec{v}_2 = \vec{v}_3$, so the set $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is linearly dependent. Which of the following is true of the vector equation $x_1\vec{v}_1 + x_2\vec{v}_2 + x_3\vec{v}_3 = \vec{0}$?

- (A) It is consistent with one solution
- (B) It is consistent with infinitely many solutions
- (C) It is inconsistent.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7**
- Section 8
- Section 9
- Section 10
- Section 11

Fact V.48

For any vector space, the set $\{\vec{v}_1, \dots, \vec{v}_n\}$ is linearly dependent if and only if the vector equation $x_1\vec{v}_1 + \dots + x_n\vec{v}_n = \vec{0}$ is consistent with infinitely many solutions.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.49 (*~10 min*) Find

$$\text{RREF} \left[\begin{array}{ccccc|c} 2 & 2 & 3 & -1 & 4 & 0 \\ 3 & 0 & 13 & 10 & 3 & 0 \\ 0 & 0 & 7 & 7 & 0 & 0 \\ -1 & 3 & 16 & 14 & 1 & 0 \end{array} \right]$$

and mark the part of the matrix that demonstrates that

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 13 \\ 7 \\ 16 \end{bmatrix}, \begin{bmatrix} -1 \\ 10 \\ 7 \\ 14 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}$$

is linearly dependent (the part that shows its linear system has infinitely many solutions).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Quick Check V.50

A set of Euclidean vectors $\{\vec{v}_1, \dots, \vec{v}_n\}$ is linearly dependent if and only if RREF $\begin{bmatrix} \vec{v}_1 & \dots & \vec{v}_n \end{bmatrix}$ has a column without a pivot position.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Observation V.51

Compare the following results:

- A set of \mathbb{R}^m vectors $\{\vec{v}_1, \dots, \vec{v}_n\}$ is linearly independent if and only if RREF $[\vec{v}_1 \ \dots \ \vec{v}_n]$ has all pivot columns.
- A set of \mathbb{R}^m vectors $\{\vec{v}_1, \dots, \vec{v}_n\}$ spans \mathbb{R}^m if and only if RREF $[\vec{v}_1 \ \dots \ \vec{v}_n]$ has all pivot rows.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.52 (~ 5 min) Is the set of Euclidean vectors

$$\left\{ \begin{bmatrix} -4 \\ 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 10 \\ 10 \\ 2 \\ 6 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 7 \\ 2 \\ 1 \end{bmatrix} \right\} \text{ linearly dependent or linearly independent?}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.52 (~ 5 min) Is the set of Euclidean vectors

$$\left\{ \begin{bmatrix} -4 \\ 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 10 \\ 10 \\ 2 \\ 6 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 7 \\ 2 \\ 1 \end{bmatrix} \right\} \text{ linearly dependent or linearly independent?}$$

Part 1: Reinterpret this question as an appropriate question about solutions to a vector equation.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7**
- Section 8
- Section 9
- Section 10
- Section 11

Activity V.52 (~ 5 min) Is the set of Euclidean vectors

$$\left\{ \begin{bmatrix} -4 \\ 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 10 \\ 10 \\ 2 \\ 6 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 7 \\ 2 \\ 1 \end{bmatrix} \right\} \text{ linearly dependent or linearly independent?}$$

Part 1: Reinterpret this question as an appropriate question about solutions to a vector equation.

Part 2: Use the solution to this question to answer the original question.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.53 (*~10 min*) Is the set of polynomials $\{x^3 + 1, x^2 + 2x, x^2 + 7x + 4\}$ linearly dependent or linearly independent?

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7**
- Section 8
- Section 9
- Section 10
- Section 11

Activity V.53 (~ 10 min) Is the set of polynomials $\{x^3 + 1, x^2 + 2x, x^2 + 7x + 4\}$ linearly dependent or linearly independent?

Part 1: Reinterpret this question as an appropriate question about solutions to a polynomial equation.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7**
- Section 8
- Section 9
- Section 10
- Section 11

Activity V.53 (~ 10 min) Is the set of polynomials $\{x^3 + 1, x^2 + 2x, x^2 + 7x + 4\}$ linearly dependent or linearly independent?

Part 1: Reinterpret this question as an appropriate question about solutions to a polynomial equation.

Part 2: Use the solution to this question to answer the original question.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7**
- Section 8
- Section 9
- Section 10
- Section 11

Summary: Today we learned how to determine if a set of vectors is **linearly independent** or **linearly dependent** (Standard V5).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 8

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Today's goals: Today we will learn about a **basis** of a vector space (Standard V6).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.54 (*~5 min*) What is the largest number of \mathbb{R}^4 vectors that can form a linearly independent set?

- (a) 3
- (b) 4
- (c) 5
- (d) You can have infinitely many vectors and still be linearly independent.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.55 (*~5 min*) What is the largest number of

$$\mathcal{P}^4 = \{ax^4 + bx^3 + cx^2 + dx + e \mid a, b, c, d, e \in \mathbb{R}\}$$

vectors that can form a linearly independent set?

- (a) 3
- (b) 4
- (c) 5
- (d) You can have infinitely many vectors and still be linearly independent.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.56 (~ 5 min) What is the largest number of

$$\mathcal{P} = \{f(x) \mid f(x) \text{ is any polynomial}\}$$

vectors that can form a linearly independent set?

- (a) 3
- (b) 4
- (c) 5
- (d) You can have infinitely many vectors and still be linearly independent.

Definition V.57

A **basis** is a linearly independent set that spans a vector space.

The **standard basis** of \mathbb{R}^n is the set $\{\vec{e}_1, \dots, \vec{e}_n\}$ where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix} \quad \dots \quad \vec{e}_n = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix}$$

For \mathbb{R}^3 , these are the vectors $\vec{e}_1 = \hat{i} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\vec{e}_2 = \hat{j} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, and $\vec{e}_3 = \hat{k} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

Observation V.58

A basis may be thought of as a collection of building blocks for a vector space, since every vector in the space can be expressed as a unique linear combination of basis vectors.

For example, in many calculus courses, vectors in \mathbb{R}^3 are often expressed in their component form

$$(3, -2, 4) = \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix}$$

or in their standard basic vector form

$$3\vec{e}_1 - 2\vec{e}_2 + 4\vec{e}_3 = 3\hat{i} - 2\hat{j} + 4\hat{k}.$$

Since every vector in \mathbb{R}^3 can be uniquely described as a linear combination of the vectors in $\{\vec{e}_1, \vec{e}_2, \vec{e}_3\}$, this set is indeed a basis.

Activity V.59 (~ 15 min) Label each of the sets A, B, C, D, E as

- SPANS \mathbb{R}^4 or DOES NOT SPAN \mathbb{R}^4
- LINEARLY INDEPENDENT or LINEARLY DEPENDENT
- BASIS FOR \mathbb{R}^4 or NOT A BASIS FOR \mathbb{R}^4

by finding RREF for their corresponding matrices.

$$A = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

$$B = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \\ 3 \end{bmatrix} \right\}$$

$$C = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 13 \\ 7 \\ 16 \end{bmatrix}, \begin{bmatrix} -1 \\ 10 \\ 7 \\ 14 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 2 \end{bmatrix} \right\}$$

$$D = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 6 \\ 1 \\ 5 \end{bmatrix} \right\}$$

$$E = \left\{ \begin{bmatrix} 5 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 1 \\ 3 \end{bmatrix} \right\}$$

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8**
- Section 9
- Section 10
- Section 11

Summary: Today we learned how to determine if a set of vectors form a **basis** of a vector space (Standard V6).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 9

Module V


Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Today's goals: Today we will learn how to find a basis of a subspace of \mathbb{R}^n .

Observation V.62

Recall that a **subspace** of a vector space is a subset that is itself a vector space.

One easy way to construct a subspace is to take the span of set, but a linearly dependent set contains “redundant” vectors. For example, only two of the three vectors in the following image are needed to span the planar subspace.



Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.63 (~ 10 min) Consider the subspace of \mathbb{R}^4 given by

$$W = \text{span} \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\}.$$

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Activity V.63 (~ 10 min) Consider the subspace of \mathbb{R}^4 given by

$$W = \text{span} \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\}.$$

Part 1: Mark the part of RREF $\begin{bmatrix} 2 & 2 & 2 & 1 \\ 3 & 0 & -3 & 5 \\ 0 & 1 & 2 & -1 \\ 1 & -1 & -3 & 0 \end{bmatrix}$ that shows that W 's spanning set is linearly dependent.

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Activity V.63 (~ 10 min) Consider the subspace of \mathbb{R}^4 given by

$$W = \text{span} \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\}.$$

Part 1: Mark the part of RREF $\begin{bmatrix} 2 & 2 & 2 & 1 \\ 3 & 0 & -3 & 5 \\ 0 & 1 & 2 & -1 \\ 1 & -1 & -3 & 0 \end{bmatrix}$ that shows that W 's spanning

set is linearly dependent.

Part 2: Find a basis for W by removing a vector from its spanning set to make it linearly independent.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9**
- Section 10
- Section 11

Fact V.64

Let $S = \{\vec{v}_1, \dots, \vec{v}_m\}$. The easiest basis describing $\text{span } S$ is the set of vectors in S given by the pivot columns of $\text{RREF}[\vec{v}_1 \ \dots \ \vec{v}_m]$.

Put another way, to compute a basis for the subspace $\text{span } S$, simply remove the vectors corresponding to the non-pivot columns of $\text{RREF}[\vec{v}_1 \ \dots \ \vec{v}_m]$. For example, since

$$\text{RREF} \begin{bmatrix} 1 & 2 & 3 \\ 0 & -2 & -2 \\ -3 & 1 & -2 \end{bmatrix} = \begin{bmatrix} \textcircled{1} & 0 & 1 \\ 0 & \textcircled{1} & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

the subspace $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \\ -2 \end{bmatrix} \right\}$ has $\left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \right\}$ as a basis.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9**
- Section 10
- Section 11

Activity V.65 (~ 10 min) Let W be the subspace of \mathbb{R}^4 given by

$$W = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ 2 \\ 1 \end{bmatrix} \right\}.$$

Find a basis for W .

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9**
- Section 10
- Section 11

Activity V.66 (*~10 min*) Let W be the subspace of \mathcal{P}^3 given by

$$W = \text{span} \{x^3 + 3x^2 + x - 1, 2x^3 - x^2 + x + 2, 4x^3 + 5x^2 + 3x, 3x^3 + 2x^2 + 2x + 1\}$$

Find a basis for W .

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9**
- Section 10
- Section 11

Activity V.67 (~ 10 min) Let W be the subspace of $M_{2,2}$ given by

$$W = \text{span} \left\{ \begin{bmatrix} 1 & 3 \\ 1 & -1 \end{bmatrix}, \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 5 \\ 3 & 0 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} \right\}.$$

Find a basis for W .

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9**
- Section 10
- Section 11

Summary: Today we learned how to find a basis of a subspace of \mathbb{R}^n .

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10**
- Section 11

Module V Section 10

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Today's goals: Today we will learn how to find the dimension of a subspace of \mathbb{R}^n .

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Observation V.68

In the previous section, we learned that computing a basis for the subspace $\text{span}\{\vec{v}_1, \dots, \vec{v}_m\}$, is as simple as removing the vectors corresponding to the non-pivot columns of $\text{RREF}[\vec{v}_1 \ \dots \ \vec{v}_m]$.

For example, since

$$\text{RREF} \begin{bmatrix} 1 & 2 & 3 \\ 0 & -2 & -2 \\ -3 & 1 & -2 \end{bmatrix} = \begin{bmatrix} \textcircled{1} & 0 & 1 \\ 0 & \textcircled{1} & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

the subspace $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \\ -2 \end{bmatrix} \right\}$ has $\left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \right\}$ as a basis.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.69 (~ 10 min) Let

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad T = \left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.69 (~ 10 min) Let

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad T = \left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Part 1: Find a basis for $\text{span } S$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.69 (~ 10 min) Let

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad T = \left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Part 1: Find a basis for $\text{span } S$.

Part 2: Find a basis for $\text{span } T$.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Observation V.70

Even though we found different bases for them, $\text{span } S$ and $\text{span } T$ are exactly the same subspace of \mathbb{R}^4 , since

$$S = \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix} \right\} = \left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\} = T$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Fact V.71

Any non-trivial vector space has infinitely-many different bases, but all the bases for a given vector space are exactly the same size.

For example,

$$\{\vec{e}_1, \vec{e}_2, \vec{e}_3\} \text{ and } \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\} \text{ and } \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \\ 5 \end{bmatrix} \right\}$$

are all valid bases for \mathbb{R}^3 , and they all contain three vectors.

Definition V.72

The **dimension** of a vector space is equal to the size of any basis for the vector space.

As you'd expect, \mathbb{R}^n has dimension n . For example, \mathbb{R}^3 has dimension 3 because any basis for \mathbb{R}^3 such as

$$\{\vec{e}_1, \vec{e}_2, \vec{e}_3\} \text{ and } \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\} \text{ and } \left\{ \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \\ 5 \end{bmatrix} \right\}$$

contains exactly three vectors.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.73 (*~10 min*) Find the dimension of each subspace of \mathbb{R}^4 by finding RREF for each corresponding matrix.

$$\begin{aligned} \text{span} \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \\ 3 \end{bmatrix} \right\} & \quad \text{span} \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 13 \\ 7 \\ 16 \end{bmatrix}, \begin{bmatrix} -1 \\ 10 \\ 7 \\ 14 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 2 \end{bmatrix} \right\} \\ \text{span} \left\{ \begin{bmatrix} 2 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 6 \\ 1 \\ 5 \end{bmatrix} \right\} & \quad \text{span} \left\{ \begin{bmatrix} 5 \\ 3 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 1 \\ 3 \end{bmatrix} \right\} \end{aligned}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Fact V.74

Every vector space with finite dimension, that is, every vector space V with a basis of the form $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is said to be **isomorphic** to a Euclidean space \mathbb{R}^n , since there exists a natural correspondance between vectors in V and vectors in \mathbb{R}^n :

$$c_1\vec{v}_1 + c_2\vec{v}_2 + \cdots + c_n\vec{v}_n \leftrightarrow \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10**
- Section 11

Observation V.75

We've already been taking advantage of the previous fact by converting polynomials and matrices into Euclidean vectors. Since \mathcal{P}^3 and $M_{2,2}$ are both four-dimensional:

$$4x^3 + 0x^2 - 1x + 5 \leftrightarrow \begin{bmatrix} 4 \\ 0 \\ -1 \\ 5 \end{bmatrix} \leftrightarrow \begin{bmatrix} 4 & 0 \\ -1 & 5 \end{bmatrix}$$

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.76 (~ 5 min) Suppose W is a subspace of \mathcal{P}^8 , and you know that the set $\{x^3 + x, x^2 + 1, x^4 - x\}$ is a linearly independent subset of W . What can you conclude about W ?

- (a) The dimension of W is at most 3.
- (b) The dimension of W is exactly 3.
- (c) The dimension of W is at least 3.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Activity V.77 (*~5 min*) Suppose W is a subspace of \mathcal{P}^8 , and you know that W is spanned by the six vectors

$$\{x^4 - x, x^3 + x, x^3 + x + 1, x^4 + 2x, x^3, 2x + 1\}.$$

What can you conclude about W ?

- (a) The dimension of W is at most 6.
- (b) The dimension of W is exactly 6.
- (c) The dimension of W is at least 6.

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Observation V.78

The space of polynomials \mathcal{P} (of *any* degree) has the basis $\{1, x, x^2, x^3, \dots\}$, so it is a natural example of an infinite-dimensional vector space.

Since \mathcal{P} and other infinite-dimensional spaces cannot be treated as an isomorphic finite-dimensional Euclidean space \mathbb{R}^n , vectors in such spaces cannot be studied by converting them into Euclidean vectors. Fortunately, most of the examples we will be interested in for this course will be finite-dimensional.

Definition V.79

A **homogeneous** system of linear equations is one of the form:

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0$$

This system is equivalent to the vector equation:

$$x_1\vec{v}_1 + \dots + x_n\vec{v}_n = \vec{0}$$

and the augmented matrix:

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & 0 \\ a_{21} & a_{22} & \cdots & a_{2n} & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & 0 \end{array} \right]$$

Activity V.80 (~ 5 min) Note that if $\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$ and $\begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$ are solutions to

$$x_1 \vec{v}_1 + \cdots + x_n \vec{v}_n = \vec{0} \text{ so is } \begin{bmatrix} a_1 + b_1 \\ \vdots \\ a_n + b_n \end{bmatrix}, \text{ since}$$

$$a_1 \vec{v}_1 + \cdots + a_n \vec{v}_n = \vec{0} \text{ and } b_1 \vec{v}_1 + \cdots + b_n \vec{v}_n = \vec{0}$$

implies

$$(a_1 + b_1) \vec{v}_1 + \cdots + (a_n + b_n) \vec{v}_n = \vec{0}.$$

Similarly, if $c \in \mathbb{R}$, $\begin{bmatrix} ca_1 \\ \vdots \\ ca_n \end{bmatrix}$ is a solution. Thus the solution set of a homogeneous system is...

a) A basis for \mathbb{R}^n .

b) A subspace of \mathbb{R}^n .

c) The empty set.

Activity V.81 (~ 10 min) Consider the homogeneous system of equations

$$x_1 + 2x_2 + x_4 = 0$$

$$2x_1 + 4x_2 - x_3 - 2x_4 = 0$$

$$3x_1 + 6x_2 - x_3 - x_4 = 0$$

Activity V.81 (~ 10 min) Consider the homogeneous system of equations

$$x_1 + 2x_2 + x_4 = 0$$

$$2x_1 + 4x_2 - x_3 - 2x_4 = 0$$

$$3x_1 + 6x_2 - x_3 - x_4 = 0$$

Part 1: Find its solution set (a subspace of \mathbb{R}^4).

Activity V.81 (~ 10 min) Consider the homogeneous system of equations

$$x_1 + 2x_2 + x_4 = 0$$

$$2x_1 + 4x_2 - x_3 - 2x_4 = 0$$

$$3x_1 + 6x_2 - x_3 - x_4 = 0$$

Part 1: Find its solution set (a subspace of \mathbb{R}^4).

Part 2: Rewrite this solution space in the form

$$\left\{ a \begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix} + b \begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix} \mid a, b \in \mathbb{R} \right\}.$$

Activity V.81 (~ 10 min) Consider the homogeneous system of equations

$$x_1 + 2x_2 + x_4 = 0$$

$$2x_1 + 4x_2 - x_3 - 2x_4 = 0$$

$$3x_1 + 6x_2 - x_3 - x_4 = 0$$

Part 1: Find its solution set (a subspace of \mathbb{R}^4).

Part 2: Rewrite this solution space in the form

$$\left\{ a \begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix} + b \begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix} \mid a, b \in \mathbb{R} \right\}.$$

Part 3: Rewrite this solution space in the form

$$\text{span} \left\{ \begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix}, \begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix} \right\}.$$

Fact V.82

The coefficients of the free variables in the solution set of a linear system always yield linearly independent vectors.

Thus if

$$\left\{ a \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} -1 \\ 0 \\ -4 \\ 1 \end{bmatrix} \mid a, b \in \mathbb{R} \right\} = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ -4 \\ 1 \end{bmatrix} \right\}$$

is the solution space for a homogeneous system, then

$$\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ -4 \\ 1 \end{bmatrix} \right\}$$

is a basis for the solution space.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10**
- Section 11

Summary: Today we learned how to find the dimension of a subspace of \mathbb{R}^n . We also started learning how to find a basis of the solution space of a **homogeneous** system of equations (Standard V9).

Module V

Section 1

Section 2

Section 3

Section 4

Section 5

Section 6

Section 7

Section 8

Section 9

Section 10

Section 11

Module V Section 11

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11**

Today's goals: Today we will practice finding a basis of the solution space of a **homogeneous** system of equations (Standard V9). This will come up again later when we find the **kernel** of a **linear transformation**.

Module V

Section 1
Section 2
Section 3
Section 4
Section 5
Section 6
Section 7
Section 8
Section 9
Section 10
Section 11

Activity V.83 (*~10 min*) Consider the homogeneous system of equations

$$2x_1 + 4x_2 + 2x_3 - 4x_4 = 0$$

$$-2x_1 - 4x_2 + x_3 + x_4 = 0$$

$$3x_1 + 6x_2 - x_3 - 4x_4 = 0$$

Find a basis for its solution space.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Activity V.84 (*~10 min*) Consider the homogeneous vector equation

$$x_1 \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ -4 \\ 6 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} + x_4 \begin{bmatrix} -4 \\ 1 \\ -4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Find a basis for its solution space.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11

Activity V.85 (*~5 min*) Consider the homogeneous system of equations

$$x_1 - 3x_2 + 2x_3 = 0$$

$$2x_1 + 6x_2 + 4x_3 = 0$$

$$x_1 + 6x_2 - 4x_3 = 0$$

Find a basis for its solution space.

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11**

Observation V.86

The basis of the trivial vector space is the empty set. You can denote this as either \emptyset or $\{\}$.

Thus, if $\vec{0}$ is the only solution of a homogeneous system, the basis of the solution space is \emptyset .

Module V

- Section 1
- Section 2
- Section 3
- Section 4
- Section 5
- Section 6
- Section 7
- Section 8
- Section 9
- Section 10
- Section 11**

Summary: Today we learned how to find a basis of the solution space of a **homogeneous** system of equations (Standard V9).