


Application Activities - Module G Part 2 - Class Day 26

Definition 26.1 The **transpose** of a matrix is given by rewriting its columns as rows and vice versa:

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Fact 26.2 It is possible to prove that the determinant of a matrix and its transpose are the same. For example, let $A = \begin{bmatrix} 3 & 3 \\ 0 & 2 \end{bmatrix}$, so $A^T = \begin{bmatrix} 3 & 0 \\ 3 & 2 \end{bmatrix}$; both matrices scale the unit square by 6, even though the parallelograms are not congruent.



Fact 26.3 We previously figured out that column operations can be used to simplify determinants; since $\det(A) = \det(A^T)$, we can also use row operations:

1. Multiplying rows by scalars: $\det \begin{bmatrix} \vdots \\ cR \\ \vdots \end{bmatrix} = c \det \begin{bmatrix} \vdots \\ R \\ \vdots \end{bmatrix}$

2. Swapping two rows: $\det \begin{bmatrix} \vdots \\ R \\ \vdots \\ S \\ \vdots \end{bmatrix} = -\det \begin{bmatrix} \vdots \\ S \\ \vdots \\ R \\ \vdots \end{bmatrix}$

3. Adding multiples of rows to other rows: $\det \begin{bmatrix} \vdots \\ R \\ \vdots \\ S \\ \vdots \end{bmatrix} = \det \begin{bmatrix} \vdots \\ R + cS \\ \vdots \\ S \\ \vdots \end{bmatrix}$

Activity 26.4 Compute the determinant of $\begin{bmatrix} 2 & 4 \\ 2 & 3 \end{bmatrix}$ by row reducing it to a nicer matrix.


For example, $\det \begin{bmatrix} 2 & 4 \\ 2 & 3 \end{bmatrix} = 2 \det \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$.

Fact 26.5 This same process allows us to prove a more convenient formula:

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

In higher dimensions, the formulas become unreasonable. For example, the formula for 4×4 matrices has 24 terms!

Activity 26.6 The following image illustrates the transformation of the unit cube by the matrix $\begin{bmatrix} 3 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.



This volume is equal to which of the following areas?

(a) $\det \begin{bmatrix} 3 & 1 \\ 1 & 1 \end{bmatrix}$

(b) $\det \begin{bmatrix} 3 & 1 \\ 1 & 0 \end{bmatrix}$

(c) $\det \begin{bmatrix} 3 & 1 \\ 0 & 1 \end{bmatrix}$

(d) $\det \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$

Fact 26.7 If column i of a matrix is \mathbf{e}_i , then both column and row i may be removed without changing the value of the determinant. For example, the second column of the following matrix is \mathbf{e}_2 , so:

$$\det \begin{bmatrix} 3 & 0 & -1 & 5 \\ 2 & 1 & 4 & 0 \\ -1 & 0 & 1 & 11 \\ 3 & 0 & 0 & 1 \end{bmatrix} = \det \begin{bmatrix} 3 & -1 & 5 \\ -1 & 1 & 11 \\ 3 & 0 & 1 \end{bmatrix}$$

Therefore the same holds for the transpose:

$$\det \begin{bmatrix} 3 & 2 & -1 & 3 \\ 0 & 1 & 0 & 0 \\ -1 & 4 & 1 & 0 \\ 5 & 0 & 11 & 1 \end{bmatrix} = \det \begin{bmatrix} 3 & -1 & 3 \\ -1 & 1 & 0 \\ 5 & 11 & 1 \end{bmatrix}$$

Geometrically, this is the fact that if the height is 1, the base \times height formula reduces to the area/volume/etc. of the $n - 1$ dimensional base.

Activity 26.8 Compute $\det \begin{bmatrix} 1 & 0 & 0 \\ 1 & 5 & 12 \\ 3 & 2 & -1 \end{bmatrix}$.

Activity 26.9 Compute $\det \begin{bmatrix} 0 & 3 & -2 \\ 1 & 5 & 12 \\ 0 & 2 & -1 \end{bmatrix}$.

(a) -1

(b) 0

(c) 1

Activity 26.10 Compute $\det \begin{bmatrix} 1 & 2 & 3 \\ 1 & -2 & -5 \\ 0 & 3 & 3 \end{bmatrix}$

Hint: $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$.

(a) 3

(b) 6

(c) 9

(d) 12

Activity 26.11 Compute $\det \begin{bmatrix} 2 & 3 & 5 & 0 \\ 0 & 3 & 2 & 0 \\ 1 & 2 & 0 & 3 \\ -1 & -1 & 2 & 2 \end{bmatrix}$.

Observation 26.12

$$\begin{aligned} \det \begin{bmatrix} 2 & 3 & 5 & 0 \\ 0 & 3 & 2 & 0 \\ 1 & 2 & 0 & 3 \\ -1 & -1 & 2 & 2 \end{bmatrix} &= (-1)(0) \det \begin{bmatrix} 2 & 3 & 5 & 0 \\ 0 & 3 & 2 & 0 \\ 1 & 2 & 0 & 3 \\ -1 & -1 & 2 & 2 \end{bmatrix} + (1)(3) \det \begin{bmatrix} 2 & 3 & 5 & 0 \\ 0 & 3 & 2 & 0 \\ 1 & 2 & 0 & 3 \\ -1 & -1 & 2 & 2 \end{bmatrix} + \\ &\quad (-1)(2) \det \begin{bmatrix} 2 & 3 & 5 & 0 \\ 0 & 3 & 2 & 0 \\ 1 & 2 & 0 & 3 \\ -1 & -1 & 2 & 2 \end{bmatrix} + (1)(0) \det \begin{bmatrix} 2 & 3 & 5 & 0 \\ 0 & 3 & 2 & 0 \\ 1 & 2 & 0 & 3 \\ -1 & -1 & 2 & 2 \end{bmatrix} \\ &= 3 \det \begin{bmatrix} 2 & 5 & 0 \\ 1 & 0 & 3 \\ -1 & 2 & 2 \end{bmatrix} + (-1)(2) \det \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 3 \\ -1 & -1 & 2 \end{bmatrix} \end{aligned}$$

This technique is called **Laplace expansion** or **cofactor expansion**.

Activity 26.13 Compute $\det \begin{bmatrix} 1 & 2 & 1 & 0 \\ 0 & 3 & 2 & -1 \\ 1 & 2 & 0 & 3 \\ -1 & -3 & 2 & -2 \end{bmatrix}$.