

Name:
J#:
Date:

Dr. Clontz

FINAL EXAM

Math 237 – Linear Algebra

Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

Standard E1.	Mark:
---------------------	-------

Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{cccc|c} 3 & -1 & 0 & 1 & 5 \\ -1 & 9 & 1 & -7 & 0 \\ 1 & 0 & -1 & 0 & -3 \end{array} \right]$$

Solution:

$$\begin{aligned} 3x_1 - x_2 + x_4 &= 5 \\ -x_1 + 9x_2 + x_3 - 7x_4 &= 0 \\ x_1 - x_3 &= -3 \end{aligned}$$

□

Standard E2.	Mark:
---------------------	-------

Find the reduced row echelon form of the matrix below.

$$\left[\begin{array}{cccc|c} 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \\ -1 & 0 & 5 & 0 & -1 \end{array} \right]$$

Solution:

$$\begin{aligned} &\left[\begin{array}{cccc|c} 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \\ -1 & 0 & 5 & 0 & -1 \end{array} \right] \sim \left[\begin{array}{cccc|c} -1 & 0 & 5 & 0 & -1 \\ 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \end{array} \right] \sim \left[\begin{array}{cccc|c} 1 & 0 & -5 & 0 & 1 \\ 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \end{array} \right] \\ &\sim \left[\begin{array}{cccc|c} 1 & 0 & -5 & 0 & 1 \\ 0 & 1 & 9 & 0 & 3 \\ 0 & -1 & 15 & -2 & -3 \end{array} \right] \sim \left[\begin{array}{cccc|c} 1 & 0 & -5 & 0 & 1 \\ 0 & 1 & 9 & 0 & 3 \\ 0 & 0 & 24 & -2 & 0 \end{array} \right] \sim \left[\begin{array}{cccc|c} 1 & 0 & -5 & 0 & 1 \\ 0 & 1 & 9 & 0 & 3 \\ 0 & 0 & 1 & -\frac{1}{12} & 0 \end{array} \right] \sim \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{5}{12} & 1 \\ 0 & 1 & 0 & \frac{3}{4} & 3 \\ 0 & 0 & 1 & -\frac{1}{12} & 0 \end{array} \right] \end{aligned}$$

□

Standard E3.	Mark:
---------------------	-------

Find the solution set for the following system of linear equations.

$$2x_1 - 2x_2 + 6x_3 - x_4 = -1$$

$$3x_1 + 6x_3 + x_4 = 5$$

$$-4x_1 + x_2 - 9x_3 + 2x_4 = -7$$

Solution: Let $A = \left[\begin{array}{cccc|c} 2 & -2 & 6 & -1 & -1 \\ 3 & 0 & 6 & 1 & 5 \\ -4 & 1 & -9 & 2 & -7 \end{array} \right]$, so $\text{RREF } A = \left[\begin{array}{cccc|c} 1 & 0 & 2 & 0 & 2 \\ 0 & 1 & -1 & 0 & 3 \\ 0 & 0 & 0 & 1 & -1 \end{array} \right]$. It follows that the solution set is given by $\begin{bmatrix} 2-2a \\ 3+a \\ a \\ -1 \end{bmatrix}$ for all real numbers a .

□

Standard E4.	Mark:
---------------------	-------

Find a basis for the solution set of the system of equations

$$x + 2y + 3z + w = 0$$

$$3x - y + z + w = 0$$

$$2x - 3y - 2z = 0$$

Solution:

$$\text{RREF} \left(\left[\begin{array}{cccc} 1 & -2 & 3 & 1 \\ 3 & -1 & 1 & 1 \\ 2 & -3 & -2 & 0 \end{array} \right] \right) = \left[\begin{array}{cccc} 1 & 0 & \frac{5}{7} & \frac{3}{7} \\ 0 & 1 & \frac{8}{7} & \frac{2}{7} \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Then the solution set is

$$\left\{ \left[\begin{array}{c} -\frac{5}{7}a - \frac{3}{7}b \\ -\frac{8}{7}a - \frac{2}{7}b \\ a \\ b \end{array} \right] \mid a, b \in \mathbb{R} \right\}$$

So a basis for the solution set is $\left\{ \left[\begin{array}{c} -\frac{5}{7} \\ -\frac{8}{7} \\ 1 \\ 0 \end{array} \right], \left[\begin{array}{c} -\frac{3}{7} \\ -\frac{2}{7} \\ 0 \\ 1 \end{array} \right] \right\}$, or $\left\{ \left[\begin{array}{c} 5 \\ 8 \\ -7 \\ 0 \end{array} \right], \left[\begin{array}{c} 3 \\ 2 \\ 0 \\ -7 \end{array} \right] \right\}$.

□

Standard V1.	Mark:
---------------------	-------

Let V be the set of all polynomials with the operations, for any $f, g \in V$, $c \in \mathbb{R}$,

$$f \oplus g = f' + g'$$

$$c \odot f = cf'$$

(here f' denotes the derivative of f).

(a) Show that scalar multiplication **distributes scalars** over vector addition: $c \odot (f \oplus g) = c \odot f \oplus c \odot g$.

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $f, g \in \mathcal{P}$, and let $c \in \mathbb{R}$.

$$c \odot (f \oplus g) = c \odot (f' + g') = c(f' + g')' = cf'' + cg'' = cf' \oplus cg' = c \odot f \oplus c \odot g.$$

However, this is not a vector space, as there is no zero vector. Additionally, $1 \odot f \neq f$ for any nonzero polynomial f .

□

Standard V2.	Mark:
---------------------	-------

Determine if $\begin{bmatrix} 4 \\ -1 \\ 6 \\ -7 \end{bmatrix}$ belongs to the span of the set $\left\{ \begin{bmatrix} 2 \\ 0 \\ -1 \\ 5 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 4 \\ 3 \end{bmatrix} \right\}$.

Solution: Since

$$\text{RREF} \left(\left[\begin{array}{cc|c} 2 & 4 & 4 \\ 0 & -1 & -1 \\ -1 & 4 & 6 \\ 5 & 3 & -7 \end{array} \right] \right) = \left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array} \right]$$

contains the contradiction $0 = 1$, $\begin{bmatrix} 4 \\ -1 \\ 6 \\ -7 \end{bmatrix}$ is not a linear combination of the three vectors.

□

Standard V3.	Mark:
---------------------	-------

Determine if the vectors $\begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$, and $\begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}$ span \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, the vectors do not span \mathbb{R}^4 .

□

Standard V4.	Mark:
---------------------	-------

Let W be the set of all complex numbers that are purely real (i.e of the form $a + 0i$) or purely imaginary (i.e. of the form $0 + bi$). Determine if W is a subspace of \mathbb{C} .

Solution: No, because 1 is purely real and i is purely imaginary, but the linear combination $1 + i$ is neither.

□

Standard S1.	Mark:
---------------------	-------

Determine if the set of matrices $\left\{ \begin{bmatrix} 3 & -1 \\ 0 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 3 & -8 \\ 6 & 5 \end{bmatrix} \right\}$ is linearly dependent or linearly independent.

Solution:

$$\text{RREF} \left(\begin{bmatrix} 3 & 1 & 3 \\ -1 & 2 & -8 \\ 0 & -2 & 6 \\ 4 & 1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Since the reduced row echelon form has a nonpivot column, the vectors are linearly dependent.

□

Standard S2.	Mark:
---------------------	-------

Determine if the set $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$ is a basis of $\mathbb{R}^{2 \times 2}$ or not.

Solution:

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

Standard S3.	Mark:
---------------------	-------

Let $W = \text{span} \left(\left(\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix} \right) \right)$. Find a basis for W .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then a basis is $\left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix} \right\}$.

□

Standard S4.	Mark:
---------------------	-------

Let $W = \text{span} \left(\left(\begin{bmatrix} 1 \\ -1 \\ 3 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ -7 \end{bmatrix} \right) \right)$. Compute the dimension of W .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 2 & 3 & 1 \\ -1 & 0 & -1 & 1 \\ 3 & 1 & 4 & 1 \\ -3 & 1 & -2 & -7 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This has 3 pivot columns so $\dim(W) = 3$.

□

Standard A1.	Mark:
---------------------	-------

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}$ be the linear transformation given by

$$T \left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \right) = [x_2 + 3x_3].$$

Write the matrix for T with respect to the standard bases of \mathbb{R}^3 and \mathbb{R} .

Solution:

$$\begin{bmatrix} 0 & 1 & 3 \end{bmatrix}$$

□

Standard A2.	Mark:
---------------------	-------

Determine if $D : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ given by $D \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = ad - bc$ is a linear transformation or not.

Solution: $D(I) = 1$ but $D(2I) = 4 \neq 2D(I)$, so D is not linear.

□

Standard A3.	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a) $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$ given by the standard matrix $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$.

(b) $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ given by the standard matrix $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix}$

Solution:

(a) $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$. Since each column is a pivot column, S is injective. Since there is no zero row, S is not surjective.

(b) Since $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$, T is not injective.

$$\text{RREF} \left(\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{4}{5} & -\frac{2}{5} \\ 0 & 1 & \frac{1}{5} & \frac{3}{5} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, T is not surjective.

□

Standard A4.	Mark:
---------------------	-------

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation given by

$$T \left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} -3x + y \\ -8x + 2y - z \\ 2y + 3z \end{bmatrix}$$

Compute a basis for the kernel and a basis for the image of T .

Solution: Let $A = \begin{bmatrix} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \end{bmatrix}$, and compute $\text{RREF}(A) = \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$. Then a basis for the image is

its columns,

$$\left\{ \begin{bmatrix} -3 \\ -8 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \right\}$$

And the kernel is the solution set of $AX = 0$, so a basis would be

$$\left\{ \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} \right\}$$

□

Standard M1.	Mark:
---------------------	-------

Let

$$A = \begin{bmatrix} 3 \\ 5 \\ -1 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 1 & -1 & 2 \\ 1 & -1 & 3 & -3 \end{bmatrix}$$

$$C = \begin{bmatrix} 2 & -1 \\ 0 & 4 \\ 3 & 1 \end{bmatrix}$$

Exactly one of the six products AB , AC , BA , BC , CA , CB can be computed. Determine which one, and compute it.

Solution: CB is the only one that can be computed, and

$$CB = \begin{bmatrix} 3 & 3 & -5 & 7 \\ 4 & -4 & 12 & -12 \\ 7 & 2 & 0 & 3 \end{bmatrix}$$

□

Standard M2.	Mark:
---------------------	-------

Determine if the matrix $\begin{bmatrix} 3 & -1 & 0 \\ 2 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$ is invertible.

Solution: It is row equivalent to the identity matrix, so it is invertible.

□

Standard M3.	Mark:
---------------------	-------

Find the inverse of the matrix $\begin{bmatrix} 8 & 5 & 3 & 0 \\ 3 & 2 & 1 & 1 \\ 5 & -3 & 1 & -2 \\ -1 & 2 & 0 & 1 \end{bmatrix}$.

Solution:

$$\text{RREF} \left(\left(\begin{array}{cccc|cccc} 8 & 5 & 3 & 0 & 1 & 0 & 0 & 0 \\ 3 & 2 & 1 & 1 & 0 & 1 & 0 & 0 \\ 5 & -3 & 1 & -2 & 0 & 0 & 1 & 0 \\ -1 & 2 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right) \right) = \left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 1 & 2 & -5 & 12 \\ 0 & 1 & 0 & 0 & 1 & 1 & -4 & -9 \\ 0 & 0 & 1 & 0 & -4 & -7 & 20 & 47 \\ 0 & 0 & 0 & 1 & -1 & 0 & 3 & 7 \end{array} \right]$$

So the inverse is $\begin{bmatrix} 1 & 2 & -5 & 12 \\ 1 & 1 & -4 & -9 \\ -4 & -7 & 20 & 47 \\ -1 & 0 & 3 & 7 \end{bmatrix}$.

□

Standard G1.	Mark:
---------------------	-------

Compute the determinant of the matrix $\begin{bmatrix} 2 & 3 & 0 & 1 \\ -1 & 3 & 1 & 4 \\ 0 & 2 & 0 & 3 \\ 1 & -1 & 3 & 5 \end{bmatrix}$.

Solution: -60 .

□

Standard G2.	Mark:
---------------------	-------

Compute the eigenvalues, along with their algebraic multiplicities, of the matrix $\begin{bmatrix} 8 & -3 & -1 \\ 21 & -8 & -3 \\ -7 & 3 & 2 \end{bmatrix}$.

Solution:

$$\begin{aligned} \det(A - \lambda I) &= (8 - \lambda) \det \begin{bmatrix} -8 - \lambda & -3 \\ 3 & 2 - \lambda \end{bmatrix} - (-3) \det \begin{bmatrix} 21 & -3 \\ -7 & 2 - \lambda \end{bmatrix} + (-1) \det \begin{bmatrix} 21 & -8 - \lambda \\ -7 & 3 \end{bmatrix} \\ &= (8 - \lambda) (\lambda^2 + 6\lambda - 7) + 3(-21\lambda + 21) - (-7\lambda + 7) \\ &= (\lambda - 1)((8 - \lambda)(\lambda + 7) - 63 + 7) \\ &= (\lambda - 1)(\lambda - \lambda^2) \\ &= -\lambda(\lambda - 1)^2 \end{aligned}$$

So the eigenvalues are 0 (with algebraic multiplicity 1) and 1 (with algebraic multiplicity 2).

□

Standard G3.	Mark:
---------------------	-------

Find the eigenspace associated to the eigenvalue 3 in the matrix $A = \begin{bmatrix} 1 & -2 & -1 & 0 \\ -4 & -1 & -2 & 0 \\ 14 & 12 & 11 & 2 \\ -14 & -10 & -9 & -1 \end{bmatrix}$.

Solution: The eigenspace is spanned by $\begin{bmatrix} -1 \\ 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix}$.

□

Standard G4.	Mark:
--------------	-------

Compute the geometric multiplicity of the eigenvalue 2 in the matrix $A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -4 & 4 & 0 & 0 \\ 11 & -6 & 1 & -1 \\ -9 & 5 & 1 & 3 \end{bmatrix}$.

Solution: The eigenspace is spanned by $\begin{bmatrix} -1 \\ -2 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ -2 \\ 0 \\ 1 \end{bmatrix}$, so the geometric multiplicity is 2.

□

Additional Notes/Marks	
------------------------	--