Application Activities - Module M Part 3 - Class Day 23

Definition 23.1 Let $T : \mathbb{R}^n \to \mathbb{R}^n$ be a linear map with standard matrix A.

- If T is a bijection and B is any $\mathbb{R}^n$ vector, then $T(X) = AX = B$ has a unique solution X.
- So we may define an **inverse map** $T^{-1} : \mathbb{R}^n \to \mathbb{R}^n$ by setting $T^{-1}(B) = X$ to be this unique solution.
- Let A^{-1} be the standard matrix for T^{-1}. We call A^{-1} the **inverse matrix** of A, so we also say that A is **invertible**.

Activity 23.2 Let $T : \mathbb{R}^n \to \mathbb{R}^n$ be a bijective linear map with standard matrix A, and let $X \in \mathbb{R}^n$. Compute $(T^{-1} \circ T)(X)$ and $A^{-1}A$.

Observation 23.3 By definition, a linear map T being bijective is equivalent to its standard matrix being invertible. Furthermore, $T^{-1} \circ T = T \circ T^{-1}$ is the identity map, and $A^{-1}A = AA^{-1} = I$ is the identity matrix.

Activity 23.4 Let $T : \mathbb{R}^3 \to \mathbb{R}^3$ be given by the matrix $A = \begin{bmatrix} 2 & -1 & -6 \\ 2 & 1 & 3 \\ 1 & 1 & 4 \end{bmatrix}$.

Part 1: Solve $T(X) = \mathbf{e}_1$ to find $T^{-1}(\mathbf{e}_1)$.
Part 2: Solve $T(X) = \mathbf{e}_2$ to find $T^{-1}(\mathbf{e}_2)$.
Part 3: Solve $T(X) = \mathbf{e}_3$ to find $T^{-1}(\mathbf{e}_3)$.
Part 4: Compute A^{-1}, the standard matrix for T^{-1}.

Observation 23.5 We could have solved these three systems simultaneously by row reducing the matrix $[A \,|\, I]$ at once.

$$A = \left[\begin{array}{ccc|ccc} 2 & -1 & -6 & 1 & 0 & 0 \\ 2 & 1 & 3 & 0 & 1 & 0 \\ 1 & 1 & 4 & 0 & 0 & 1 \end{array}\right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -2 & 3 \\ 0 & 1 & 0 & -5 & 14 & -18 \\ 0 & 0 & 1 & 1 & -3 & 4 \end{array}\right]$$

Activity 23.6 Find the inverse A^{-1} of the matrix $A = \begin{bmatrix} 1 & 3 \\ 0 & -2 \end{bmatrix}$ by row-reducing $[A \,|\, I]$.

Activity 23.7 Is the matrix $\begin{bmatrix} 2 & 3 & 1 \\ -1 & -4 & 2 \\ 0 & -5 & 5 \end{bmatrix}$ invertible? Give a reason for your answer.

Observation 23.8 A matrix $A \in \mathbb{R}^{n \times n}$ is invertible if and only if $\text{RREF}(A) = I_n$.