

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 15**

Math 237 – Linear Algebra

**Version 4**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V2.** Determine if  $\begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$  can be written as a linear combination of the vectors  $\begin{bmatrix} -1 \\ -9 \\ 15 \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ 5 \\ -5 \end{bmatrix}$ .

**Solution:**

$$\text{RREF} \left( \left[ \begin{array}{cc|c} -1 & 1 & 0 \\ -9 & 5 & 0 \\ 15 & -5 & 2 \end{array} \right] \right) = \left[ \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right]$$

Since this system has no solution,  $\begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$  cannot be written as a linear combination of the vectors  $\begin{bmatrix} -1 \\ -9 \\ 15 \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ 5 \\ -5 \end{bmatrix}$ .

□

**S1.** Determine if the set of polynomials  $\{-3x^3 - 8x^2, x^3 + 2x^2 + 2, -x^2 + 3\}$  is linearly dependent or linearly independent

**Solution:**

$$\text{RREF} \left( \left[ \begin{array}{ccc} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 0 & 0 \\ 0 & 2 & 3 \end{array} \right] \right) = \left[ \begin{array}{ccc} 1 & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

This has a non pivot column, therefore the set is linearly dependent.

□

**S3.** Let  $W = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ -1 \\ 3 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ -7 \end{bmatrix} \right\} \right)$ . Find a basis of  $W$ .

**Solution:**

$$\text{RREF} \left( \left[ \begin{array}{cccc} 1 & 2 & 3 & 1 \\ -1 & 0 & -1 & 1 \\ 3 & 1 & 4 & 1 \\ -3 & 1 & -2 & -7 \end{array} \right] \right) = \left[ \begin{array}{cccc} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Then  $\left\{ \begin{bmatrix} 1 \\ -1 \\ 3 \\ -3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ -7 \end{bmatrix} \right\}$  is a basis for  $W$ .

□

**S4.** Let  $W$  be the subspace of  $\mathcal{P}_3$  given by  
 $W = \text{span}(\{x^3 - x^2 + 3x - 3, 2x^3 + x + 1, 3x^3 - x^2 + 4x - 2, x^3 + x^2 + x - 7\})$ . Compute the dimension of  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 2 & 3 & 1 \\ -1 & 0 & -1 & 1 \\ 3 & 1 & 4 & 1 \\ -3 & 1 & -2 & -7 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This has 3 pivot columns so  $\dim(W) = 3$ .

□

**V2:**

**S1:**

**S3:**

**S4:**