

Name: \_\_\_\_\_

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all polynomials with the operations, for any  $f, g \in V$ ,  $c \in \mathbb{R}$ ,

$$f \oplus g = f' + g'$$

$$c \odot f = cf'$$

(here  $f'$  denotes the derivative of  $f$ ).

(a) Show that scalar multiplication **distributes scalars** over vector addition:  $c \odot (f \oplus g) = c \odot f \oplus c \odot g$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $f, g \in \mathcal{P}$ , and let  $c \in \mathbb{R}$ .

$$c \odot (f \oplus g) = c \odot (f' + g') = c(f' + g')' = cf'' + cg'' = cf' \oplus cg' = c \odot f \oplus c \odot g.$$

However, this is not a vector space, as there is no zero vector. Additionally,  $1 \odot f \neq f$  for any nonzero polynomial  $f$ .

□

**V3.** Determine if the vectors  $\begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, the vectors do not span  $\mathbb{R}^4$ .

□

**V4.** Let  $W$  be the set of all polynomials of even degree. Determine if  $W$  is a subspace of the vector space of all polynomials.

**Solution:**  $W$  is closed under scalar multiplication, but not under addition. For example,  $x - x^2$  and  $x^2$  are both in  $W$ , but  $(x - x^2) + (x^2) = x \notin W$ .

□

**S2.** Determine if the set  $\{x^3 - x, x^2 + x + 1, x^3 - x^2 + 2, 2x^2 - 1\}$  is a basis of  $\mathcal{P}_3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 \\ -1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix is not the identity matrix, it is not a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**