

Name: _____

MASTERY QUIZ DAY 10

Math 237 – Linear Algebra

Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{ccc|c} -4 & -1 & 3 & 2 \\ 1 & 2 & -1 & 0 \\ -1 & 4 & 1 & 4 \end{array} \right]$$

Solution:

$$-4x_1 - x_2 + 3x_3 = 2$$

$$x_1 + 2x_2 - x_3 = 0$$

$$-x_1 + 4x_2 + x_3 = 4$$

□

E3. Solve the following linear system.

$$4x_1 + 4x_2 + 3x_3 - 6x_4 = 5$$

$$-2x_3 - 4x_4 = 3$$

$$2x_1 + 2x_2 + x_3 - 4x_4 = -1$$

Solution: Let $A = \left[\begin{array}{cccc|c} 4 & 4 & 3 & -6 & 5 \\ 0 & 0 & -2 & -4 & 3 \\ 2 & 2 & 1 & -4 & -1 \end{array} \right]$, so $\text{RREF } A = \left[\begin{array}{cccc|c} 1 & 1 & 0 & -3 & 0 \\ 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$. It follows that the system is inconsistent with no solutions (since the bottom row implies the contradiction $0 = 1$).

□

E4. Find a basis for the solution set to the homogeneous system of equations

$$2x_1 + 3x_2 - 5x_3 + 14x_4 = 0$$

$$x_1 + x_2 - x_3 + 5x_4 = 0$$

Solution: Let $A = \left[\begin{array}{cccc|c} 2 & 3 & -5 & 14 & 0 \\ 1 & 1 & -1 & 5 & 0 \end{array} \right]$, so $\text{RREF } A = \left[\begin{array}{cccc|c} 1 & 0 & 2 & 1 & 1 \\ 0 & 1 & -3 & 4 & 2 \end{array} \right]$. It follows that the basis for the solution set is given by $\left\{ \begin{bmatrix} -2 \\ 3 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ -4 \\ 0 \\ 1 \end{bmatrix} \right\}$.

□

V1. Let V be the set of all real numbers together with the operations \oplus and \odot defined by, for any $x, y \in V$ and $c \in \mathbb{R}$,

$$\begin{aligned}x \oplus y &= x + y - 3 \\c \odot x &= cx - 3(c - 1)\end{aligned}$$

- (a) Show that **scalar multiplication** is **associative**: $a \odot (b \odot x) = (ab) \odot x$.
 (b) Determine if V is a vector space or not. Justify your answer

Solution: Let $x, y \in V$, $c, d \in \mathbb{R}$. To show associativity:

$$\begin{aligned}c \odot (d \odot x) &= c \odot (dx - 3(d - 1)) \\&= c(dx - 3(d - 1)) - 3(c - 1) \\&= cdx - 3(cd - 1) \\&= (cd) \odot x\end{aligned}$$

We verify the remaining 7 properties to see that V is a vector space.

- 1) Real addition is associative, so \oplus is associative.
- 2) $x \oplus 3 = x + 3 - 3 = x$, so 3 is the additive identity.
- 3) $x \oplus (6 - x) = x + (6 - x) - 3 = 3$, so $6 - x$ is the additive inverse of x .
- 4) Real addition is commutative, so \oplus is commutative.
- 5) Associativity shown above
- 6) $1 \odot x = x - 3(1 - 1) = x$
- 7)

$$\begin{aligned}c \odot (x \oplus y) &= c \odot (x + y - 3) \\&= c(x + y - 3) - 3(c - 1) \\&= cx - 3(c - 1) + cy - 3(c - 1) - 3 \\&= (c \odot x) \oplus (c \odot y)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot x &= (c + d)x - 3(c + d - 1) \\&= cx - 3(c - 1) + dx - 3(d - 1) - 3 \\&= (c \odot x) \oplus (d \odot x)\end{aligned}$$

Therefore V is a vector space.

□

E1:

E3:

E4:

V1:

E2:

Name: _____

MASTERY QUIZ DAY 10

Math 237 – Linear Algebra

Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{ccc|c} 2 & -1 & 0 & 1 \\ -1 & 4 & 1 & -7 \\ 1 & 2 & -1 & 0 \end{array} \right]$$

Solution:

$$\begin{aligned} 2x_1 - x_2 &= 1 \\ -x_1 + 4x_2 + x_3 &= -7 \\ x_1 + 2x_2 - x_3 &= 0 \end{aligned}$$

□

E3. Solve the following linear system.

$$\begin{aligned} 4x_1 + 4x_2 + 3x_3 - 6x_4 &= 5 \\ -2x_3 - 4x_4 &= 3 \\ 2x_1 + 2x_2 + x_3 - 4x_4 &= -1 \end{aligned}$$

Solution: Let $A = \left[\begin{array}{cccc|c} 4 & 4 & 3 & -6 & 5 \\ 0 & 0 & -2 & -4 & 3 \\ 2 & 2 & 1 & -4 & -1 \end{array} \right]$, so $\text{RREF } A = \left[\begin{array}{cccc|c} 1 & 1 & 0 & -3 & 0 \\ 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$. It follows that the system is inconsistent with no solutions (since the bottom row implies the contradiction $0 = 1$).

□

E4. Find a basis for the solution set of the system of equations

$$\begin{aligned} x + 3y + 3z + 7w &= 0 \\ x + 3y - z - w &= 0 \\ 2x + 6y + 3z + 8w &= 0 \\ x + 3y - 2z - 3w &= 0 \end{aligned}$$

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the solution set is

$$\left\{ \begin{bmatrix} -3a - b \\ a \\ -2b \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

So a basis for the solution set is

$$\left\{ \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \\ -1 \end{bmatrix} \right\}$$

□

V1. Let V be the set of all polynomials with the operations, for any $f, g \in V$, $c \in \mathbb{R}$,

$$\begin{aligned} f \oplus g &= f' + g' \\ c \odot f &= cf' \end{aligned}$$

(here f' denotes the derivative of f).

- (a) Show that scalar multiplication **distributes scalars** over vector addition: $c \odot (f \oplus g) = c \odot f \oplus c \odot g$.
- (b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $f, g \in \mathcal{P}$, and let $c \in \mathbb{R}$.

$$c \odot (f \oplus g) = c \odot (f' + g') = c(f' + g')' = cf'' + cg'' = cf' \oplus cg' = c \odot f \oplus c \odot g.$$

However, this is not a vector space, as there is no zero vector. Additionally, $1 \odot f \neq f$ for any nonzero polynomial f .

□

E1:

E3:

E4:

V1:

E2:

Name: _____

MASTERY QUIZ DAY 10

Math 237 – Linear Algebra

Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write an augmented matrix corresponding to the following system of linear equations.

$$\begin{aligned}x_1 + 4x_3 &= 1 \\x_2 - x_3 &= 7 \\x_1 - x_2 + 3x_4 &= -1\end{aligned}$$

Solution:

$$\left[\begin{array}{cccc|c} 1 & 0 & 4 & 0 & 1 \\ 0 & 1 & -1 & 0 & 7 \\ 1 & -1 & 0 & 3 & -1 \end{array} \right]$$

□

E3. Solve the system of linear equations.

$$\begin{aligned}2x + y - z + w &= 5 \\3x - y - 2w &= 0 \\-x + 5z + 3w &= -1\end{aligned}$$

Solution:

$$\text{RREF} \left(\left[\begin{array}{cccc|c} 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \\ -1 & 0 & 5 & 0 & -1 \end{array} \right] \right) = \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{12} & 1 \\ 0 & 1 & 0 & \frac{7}{12} & 3 \\ 0 & 0 & 1 & \frac{4}{12} & 0 \end{array} \right]$$

So the solutions are

$$\left\{ \left[\begin{array}{c} 1 + a \\ 3 - 21a \\ -7a \\ 12a \end{array} \right] \mid a \in \mathbb{R} \right\}$$

□

E4. Find a basis for the solution set of the system of equations

$$\begin{aligned}x + 2y + 3z + w &= 0 \\3x - y + z + w &= 0 \\2x - 3y - 2z &= 0 \\-x + 2z + 5w &= 0\end{aligned}$$

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -1 & 1 & 1 \\ 2 & -3 & -2 & 0 \\ -1 & 0 & 2 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the solution set is

$$\left\{ \begin{bmatrix} a \\ 2a \\ -2a \\ a \end{bmatrix} \mid a \in \mathbb{R} \right\}$$

So a basis for the solution set is $\left\{ \begin{bmatrix} 1 \\ 2 \\ -2 \\ 1 \end{bmatrix} \right\}$.

□

V1. Let V be the set of all real numbers together with the operations \oplus and \odot defined by, for any $x, y \in V$ and $c \in \mathbb{R}$,

$$\begin{aligned} x \oplus y &= x + y - 3 \\ c \odot x &= cx - 3(c - 1) \end{aligned}$$

- (a) Show that **scalar multiplication** is **associative**: $a \odot (b \odot x) = (ab) \odot x$.
- (b) Determine if V is a vector space or not. Justify your answer

Solution: Let $x, y \in V$, $c, d \in \mathbb{R}$. To show associativity:

$$\begin{aligned} c \odot (d \odot x) &= c \odot (dx - 3(d - 1)) \\ &= c(dx - 3(d - 1)) - 3(c - 1) \\ &= cdx - 3(cd - 1) \\ &= (cd) \odot x \end{aligned}$$

We verify the remaining 7 properties to see that V is a vector space.

- 1) Real addition is associative, so \oplus is associative.
- 2) $x \oplus 3 = x + 3 - 3 = x$, so 3 is the additive identity.
- 3) $x \oplus (6 - x) = x + (6 - x) - 3 = 3$, so $6 - x$ is the additive inverse of x .
- 4) Real addition is commutative, so \oplus is commutative.
- 5) Associativity shown above
- 6) $1 \odot x = x - 3(1 - 1) = x$
- 7)

$$\begin{aligned} c \odot (x \oplus y) &= c \odot (x + y - 3) \\ &= c(x + y - 3) - 3(c - 1) \\ &= cx - 3(c - 1) + cy - 3(c - 1) - 3 \\ &= (c \odot x) \oplus (c \odot y) \end{aligned}$$

8)

$$\begin{aligned}(c+d) \odot x &= (c+d)x - 3(c+d-1) \\ &= cx - 3(c-1) + dx - 3(d-1) - 3 \\ &= (c \odot x) \oplus (d \odot x)\end{aligned}$$

Therefore V is a vector space.

□

E1:

E3:

E4:

V1:

E2:

Name: _____

MASTERY QUIZ DAY 10

Math 237 – Linear Algebra

Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{cccc|c} 3 & -1 & 0 & 1 & 5 \\ -1 & 9 & 1 & -7 & 0 \\ 1 & 0 & -1 & 0 & -3 \end{array} \right]$$

Solution:

$$\begin{aligned} 3x_1 - x_2 + x_4 &= 5 \\ -x_1 + 9x_2 + x_3 - 7x_4 &= 0 \\ x_1 - x_3 &= -3 \end{aligned}$$

□

E3. Solve the system of equations

$$\begin{aligned} -3x + y &= 2 \\ -8x + 2y - z &= 6 \\ 2y + 3z &= -2 \end{aligned}$$

Solution:

$$\text{RREF} \left(\left[\begin{array}{ccc|c} -3 & 1 & 0 & 2 \\ -8 & 2 & -1 & 6 \\ 0 & 2 & 3 & -2 \end{array} \right] \right) = \left[\begin{array}{ccc|c} 1 & 0 & \frac{1}{2} & -1 \\ 0 & 1 & \frac{3}{2} & -1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

The solutions are

$$\left\{ \left[\begin{array}{c} -1 - \frac{c}{2} \\ -1 - \frac{3c}{2} \\ c \end{array} \right] \mid c \in \mathbb{R} \right\} = \left\{ \left[\begin{array}{c} c - 1 \\ 3c - 1 \\ -2c \end{array} \right] \mid c \in \mathbb{R} \right\}$$

□

E4. Find a basis for the solution set to the system of equations

$$\begin{aligned} x + 2y - 3z &= 0 \\ 2x + y - 4z &= 0 \\ 3y - 2z &= 0 \\ x - y - z &= 0 \end{aligned}$$

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -4 \\ 0 & 3 & -2 \\ 1 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{5}{3} \\ 0 & 1 & -\frac{2}{3} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Then the solution set is

$$\left\{ \begin{bmatrix} \frac{5}{3}a \\ \frac{2}{3}a \\ \frac{2}{3}a \\ a \end{bmatrix} \mid a \in \mathbb{R} \right\}$$

So a basis is $\left\{ \begin{bmatrix} \frac{5}{3} \\ \frac{2}{3} \\ \frac{2}{3} \\ 1 \end{bmatrix} \right\}$ or $\left\{ \begin{bmatrix} 5 \\ 2 \\ 2 \\ 3 \end{bmatrix} \right\}$.

□

V1. Let V be the set of all pairs of real numbers with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned} (x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2, y_1 + y_2) \\ c \odot (x_1, y_1) &= (0, cy_1) \end{aligned}$$

(a) Show that scalar multiplication **distributes vectors** over scalar addition:

$$(c + d) \odot (x, y) = c \odot (x, y) \oplus d \odot (x, y).$$

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$, and let $c, d \in \mathbb{R}$. Then

$$(c + d) \odot (x_1, y_1) = (0, (c + d)y_1) = (0, cy_1) \oplus (0, dy_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

However, V is not a vector space, as $1 \odot (x_1, y_1) = (0, y_1) \neq (x_1, y_1)$.

□

E1:

E3:

E4:

V1:

E2:

Name: _____

MASTERY QUIZ DAY 10

Math 237 – Linear Algebra

Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{ccc|c} -4 & -1 & 3 & 2 \\ 1 & 2 & -1 & 0 \\ -1 & 4 & 1 & 4 \end{array} \right]$$

Solution:

$$-4x_1 - x_2 + 3x_3 = 2$$

$$x_1 + 2x_2 - x_3 = 0$$

$$-x_1 + 4x_2 + x_3 = 4$$

□

E3. Solve the system of linear equations.

$$2x + y - z + w = 5$$

$$3x - y - 2w = 0$$

$$-x + 5z + 3w = -1$$

Solution:

$$\text{RREF} \left(\left[\begin{array}{cccc|c} 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \\ -1 & 0 & 5 & 0 & -1 \end{array} \right] \right) = \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{12} & 1 \\ 0 & 1 & 0 & \frac{4}{7} & 3 \\ 0 & 0 & 1 & \frac{4}{12} & 0 \end{array} \right]$$

So the solutions are

$$\left\{ \left[\begin{array}{c} 1+a \\ 3-21a \\ -7a \\ 12a \end{array} \right] \mid a \in \mathbb{R} \right\}$$

□

E4. Find a basis for the solution set of the system of equations

$$x + 2y + 3z + w = 0$$

$$3x - y + z + w = 0$$

$$2x - 3y - 2z = 0$$

$$-x + 2z + 5w = 0$$

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & -2 & 3 & 1 \\ 3 & -1 & 1 & 1 \\ 2 & -3 & -2 & 0 \\ -1 & 0 & 2 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the solution set is

$$\left\{ \begin{bmatrix} a \\ 2a \\ -2a \\ a \end{bmatrix} \mid a \in \mathbb{R} \right\}$$

So a basis for the solution set is $\left\{ \begin{bmatrix} 1 \\ 2 \\ -2 \\ 1 \end{bmatrix} \right\}$.

□

V1. Let V be the set of all real numbers with the operations, for any $x, y \in V$, $c \in \mathbb{R}$,

$$x \oplus y = \sqrt{x^2 + y^2}$$

$$c \odot x = cx$$

(a) Show that the vector **addition** \oplus is **associative**: $x \oplus (y \oplus z) = (x \oplus y) \oplus z$.

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $x, y, z \in \mathbb{R}$. Then

$$\begin{aligned} (x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\ &= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\ &= \sqrt{x^2 + y^2 + z^2} \\ &= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\ &= x \oplus \sqrt{y^2 + z^2} \\ &= x \oplus (y \oplus z) \end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

E1:

E3:

E4:

V1:

E2:

Name: _____

MASTERY QUIZ DAY 10

Math 237 – Linear Algebra

Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{ccc|c} 2 & -1 & 0 & 1 \\ -1 & 4 & 1 & -7 \\ 1 & 2 & -1 & 0 \end{array} \right]$$

Solution:

$$\begin{aligned} 2x_1 - x_2 &= 1 \\ -x_1 + 4x_2 + x_3 &= -7 \\ x_1 + 2x_2 - x_3 &= 0 \end{aligned}$$

□

E3. Solve the system of linear equations.

$$\begin{aligned} 2x + y - z + w &= 5 \\ 3x - y - 2w &= 0 \\ -x + 5z + 3w &= -1 \end{aligned}$$

Solution:

$$\text{RREF} \left(\left[\begin{array}{cccc|c} 2 & 1 & -1 & 0 & 5 \\ 3 & -1 & 0 & -2 & 0 \\ -1 & 0 & 5 & 0 & -1 \end{array} \right] \right) = \left[\begin{array}{cccc|c} 1 & 0 & 0 & -\frac{1}{12} & 1 \\ 0 & 1 & 0 & \frac{7}{12} & 3 \\ 0 & 0 & 1 & \frac{4}{12} & 0 \end{array} \right]$$

So the solutions are

$$\left\{ \left[\begin{array}{c} 1+a \\ 3-21a \\ -7a \\ 12a \end{array} \right] \mid a \in \mathbb{R} \right\}$$

□

E4. Find a basis for the solution set to the homogeneous system of equations

$$\begin{aligned} 4x_1 + 4x_2 + 3x_3 - 6x_4 &= 0 \\ -2x_3 - 4x_4 &= 0 \\ 2x_1 + 2x_2 + x_3 - 4x_4 &= 0 \end{aligned}$$

Solution: Let $A = \left[\begin{array}{cccc|c} 4 & 4 & 3 & -6 & 0 \\ 0 & 0 & -2 & -4 & 0 \\ 2 & 2 & 1 & -4 & 0 \end{array} \right]$, so RREF $A = \left[\begin{array}{cccc|c} 1 & 1 & 0 & -3 & 0 \\ 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$. It follows that the basis for the solution set is given by $\left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 0 \\ -2 \\ 1 \end{bmatrix} \right\}$.

□

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2 - 1, y_1 + y_2 - 1)$$

$$c \odot (x_1, y_1) = (cx_1 - (c - 1), cy_1 - (c - 2))$$

- (a) Show that this vector space has an **additive identity** element $\mathbf{0}$ satisfying $(x, y) \oplus \mathbf{0} = (x, y)$.
 (b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$; then $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

Now we will show the other seven properties. Let $(x_1, y_1), (x_2, y_2) \in V$, and let $c, d \in \mathbb{R}$.

- 1) Since real addition is associative, \oplus is associative.
- 2) Since real addition is commutative, \oplus is commutative.
- 3) The additive identity is $(1, 1)$.
- 4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .
- 5)

$$\begin{aligned} c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1) \end{aligned}$$

$$6) 1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$$

7)

$$\begin{aligned} c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2) \end{aligned}$$

8)

$$\begin{aligned} (c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1) \end{aligned}$$

Therefore V is a vector space.

□

E1: **E3:** **E4:** **V1:** **E2:**