

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 1**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 5 \\ -1 \\ -2 \end{bmatrix}$ ,  $\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$ , and  $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} -3 & 5 & 2 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & -2 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix has only two pivot columns, the vectors do not span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all polynomials of even degree. Determine if  $W$  is a subspace of the vector space of all polynomials.

**Solution:**  $W$  is closed under scalar multiplication, but not under addition. For example,  $x - x^2$  and  $x^2$  are both in  $W$ , but  $(x - x^2) + (x^2) = x \notin W$ .

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 2**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span  $\mathbb{R}^4$ .

□

**V4.** Let  $W$  be the set of all complex numbers that are purely real (i.e. of the form  $a + 0i$ ) or purely imaginary (i.e. of the form  $0 + bi$ ). Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** No, because 1 is purely real and  $i$  is purely imaginary, but the linear combination  $1 + i$  is neither.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 0 \\ -1 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & 0 & -1 & 0 \\ 1 & 2 & 0 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 3**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 5 \\ -1 \\ -2 \end{bmatrix}$ ,  $\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$ , and  $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} -3 & 5 & 2 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & -2 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix has only two pivot columns, the vectors do not span  $\mathbb{R}^3$ .

□

**V4.** Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

**S2.** Determine if the set  $\{x^2 + x - 1, 3x^2 - x + 1, 2x - 2\}$  is a basis of  $\mathcal{P}_2$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 4**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, the vectors do not span  $\mathbb{R}^4$ .

□

**V4.** Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

**S2.** Determine if the set  $\{x^2 + x - 1, 3x^2 - x + 1, 2x - 2\}$  is a basis of  $\mathcal{P}_2$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 5**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Does  $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$ ?

**Solution:** Since

$$\text{RREF} \begin{bmatrix} 2 & 3 & 1 & -4 \\ -1 & 12 & 2 & 2 \\ 4 & -9 & 3 & -8 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

lacks a zero row, the vectors span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all polynomials of even degree. Determine if  $W$  is a subspace of the vector space of all polynomials.

**Solution:**  $W$  is closed under scalar multiplication, but not under addition. For example,  $x - x^2$  and  $x^2$  are both in  $W$ , but  $(x - x^2) + (x^2) = x \notin W$ .

□

**S2.** Determine if the set  $\{x^2 + x - 1, 3x^2 - x + 1, 2x - 2\}$  is a basis of  $\mathcal{P}_2$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 6**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 5 \\ -1 \\ -2 \end{bmatrix}$ ,  $\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$ , and  $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} -3 & 5 & 2 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & -2 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix has only two pivot columns, the vectors do not span  $\mathbb{R}^3$ .

□

**V4.** Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**