

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 25

Math 237 – Linear Algebra

### Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the standard matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the standard matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 7 & -1 & 5 \end{bmatrix}$

### Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 7 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$

Since there is not a zero row,  $T$  is surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^{2 \times 3} \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \right) = \begin{bmatrix} a+x \\ b+y \\ c+z \end{bmatrix}$ . Compute a basis for the kernel and a basis for the image of  $T$ .

**Solution:** Rewrite as  $T' \begin{pmatrix} a \\ b \\ c \\ x \\ y \\ z \end{pmatrix} = \begin{bmatrix} a+x \\ b+y \\ c+z \end{bmatrix}$ .

$$\text{RREF} \left( \begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$

Thus  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$  is a basis for the image, and  $\left\{ \begin{bmatrix} -1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -1 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 1 \end{bmatrix} \right\}$  is a basis for the kernel.

□

Additional Notes/Marks	
------------------------	--

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 25

Math 237 – Linear Algebra

### Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  where  $S(\vec{e}_1) = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$ ,  $S(\vec{e}_2) = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ ,  $S(\vec{e}_3) = \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$ , and  $S(\vec{e}_4) = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ ,

(b)  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  where  $T(\vec{e}_1) = \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$ ,  $T(\vec{e}_2) = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}$ , and  $T(\vec{e}_3) = \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$ .

**Solution:**

(a)  $\text{RREF} \begin{bmatrix} 2 & 1 & 0 & 3 \\ 1 & 2 & -1 & 2 \\ 0 & 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$ . The map is not injective since it has a column without pivot, but it is surjective because every row has a pivot.

(b)  $\text{RREF} \begin{bmatrix} 2 & 1 & 1 \\ 2 & 0 & 2 \\ 1 & 4 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$ . The map is not injective since there is a column without a pivot, and it is not surjective because there is a row without a pivot.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} 8x - 3y - z \\ y + 3z \\ -7x + 3y + 2z \end{bmatrix}$ . Compute a basis for the kernel and a basis for the image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 \\ 0 & 1 & 3 \\ -7 & 3 & 2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

Thus  $\left\{ \begin{bmatrix} 8 \\ 0 \\ -7 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 3 \end{bmatrix} \right\}$  is a basis for the image, and  $\left\{ \begin{bmatrix} -1 \\ -3 \\ 1 \end{bmatrix} \right\}$  is a basis for the kernel.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 25

Math 237 – Linear Algebra

## Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the standard matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the standard matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix}$

## Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{4}{5} & -\frac{2}{5} \\ 0 & 1 & \frac{1}{5} & \frac{3}{5} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row,  $T$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} a & b \\ x & y \end{bmatrix} \right) = \begin{bmatrix} a+x \\ 0 \\ b+y \end{bmatrix}$ . Compute a basis for the kernel and a basis for the image of  $T$ .

**Solution:** Rewrite as  $T' \left( \begin{bmatrix} a \\ b \\ x \\ y \end{bmatrix} \right) = \begin{bmatrix} a+x \\ 0 \\ b+y \end{bmatrix}$ .

$$\text{RREF} \left( \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

Thus  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$  is a basis for the image, and  $\left\{ \begin{bmatrix} -1 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ 0 & 1 \end{bmatrix} \right\}$  is a basis for the kernel.

□

Additional Notes/Marks	
------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 25

Math 237 – Linear Algebra

## Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  where  $S(\vec{e}_1) = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$ ,  $S(\vec{e}_2) = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ ,  $S(\vec{e}_3) = \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$ , and  $S(\vec{e}_4) = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ ,

(b)  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  where  $T(\vec{e}_1) = \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$ ,  $T(\vec{e}_2) = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}$ , and  $T(\vec{e}_3) = \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$ .

**Solution:**

(a)  $\text{RREF} \begin{bmatrix} 2 & 1 & 0 & 3 \\ 1 & 2 & -1 & 2 \\ 0 & 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$ . The map is not injective since it has a column without pivot, but it is surjective because every row has a pivot.

(b)  $\text{RREF} \begin{bmatrix} 2 & 1 & 1 \\ 2 & 0 & 2 \\ 1 & 4 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$ . The map is not injective since there is a column without a pivot, and it is not surjective because there is a row without a pivot.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathcal{P}^3 \rightarrow \mathcal{P}^3$  be the linear transformation given by

$$T(ax^3 + bx^2 + cx + d) = (a + 3b + 3c + 7d)x^3 + (a + 3b - c - d)x^2 + (2a + 6b + 3c + 8d)x + (a + 3b - 2c - 3d)$$

Compute a basis for the kernel and a basis for the image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then a basis for the kernel is

$$\{-3x^3 + x^2, -x^3 - 2x + 1\}$$

and a basis for the image is

$$\{x^3 + x^2 + 2x + 1, 3x^3 - x^2 + 3x - 2\}$$

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 25

Math 237 – Linear Algebra

### Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the standard matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the standard matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix}$

### Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{4}{5} & -\frac{2}{5} \\ 0 & 1 & \frac{1}{5} & \frac{3}{5} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row,  $T$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} 8x - 3y - z + 4w \\ y + 3z - 4w \\ -7x + 3y + 2z - 5w \end{bmatrix}$ . Compute a basis for the kernel and a basis for the image of  $T$ .



**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 0 & 1 & 3 & -4 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Thus  $\left\{ \begin{bmatrix} 8 \\ 0 \\ -7 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 3 \end{bmatrix} \right\}$  is a basis for the image, and  $\left\{ \begin{bmatrix} 1 \\ 3 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ 0 \\ 1 \end{bmatrix} \right\}$  is a basis for the kernel.

□

Additional Notes/Marks	
------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 25

Math 237 – Linear Algebra

## Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by the standard matrix  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the standard matrix  $\begin{bmatrix} 2 & 3 & -1 & -2 \\ 0 & 1 & 3 & 1 \\ 2 & 1 & -7 & -4 \end{bmatrix}$

## Solution:

(a)  $\text{RREF} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is no zero row,  $S$  is surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & -2 \\ 0 & 1 & 3 & 1 \\ 2 & 1 & -7 & -4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -5 & -\frac{5}{2} \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there are only two pivot columns,  $T$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the linear transformation given by

$$T \left( \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} -3x + y \\ -8x + 2y - z \\ 2y + 3z \end{bmatrix}$$

Compute a basis for the kernel and a basis for the image of  $T$ .

**Solution:** Let  $A = \begin{bmatrix} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \end{bmatrix}$ , and compute  $\text{RREF}(A) = \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$ . Then a basis for the image is

its columns,

$$\left\{ \begin{bmatrix} -3 \\ -8 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \right\}$$

And the kernel is the solution set of  $AX = 0$ , so a basis would be

$$\left\{ \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} \right\}$$

□

Additional Notes/Marks	
------------------------	--