

Name:
J#:
Date:

Dr. Clontz

# MIDTERM EXAM

Math 237 – Linear Algebra

## Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard E1.</b>	Mark:
---------------------	-------

Write a system of linear equations corresponding to the following augmented matrix.

$$\left[ \begin{array}{cccc|c} 3 & -1 & 0 & 1 & 5 \\ -1 & 9 & 1 & -7 & 0 \\ 1 & 0 & -1 & 0 & -3 \end{array} \right]$$

**Solution:**

$$\begin{aligned} 3x_1 - x_2 + x_4 &= 5 \\ -x_1 + 9x_2 + x_3 - 7x_4 &= 0 \\ x_1 - x_3 &= -3 \end{aligned}$$

□

<b>Standard E2.</b>	Mark:
---------------------	-------

Put the following matrix in reduced row echelon form.

$$\left[ \begin{array}{ccc|c} -3 & 1 & 0 & 2 \\ -8 & 2 & -1 & 6 \\ 0 & 2 & 3 & -2 \end{array} \right]$$

**Solution:**

$$\begin{aligned} \left[ \begin{array}{ccc|c} -3 & 1 & 0 & 2 \\ -8 & 2 & -1 & 6 \\ 0 & 2 & 3 & -2 \end{array} \right] &\sim \left[ \begin{array}{ccc|c} 1 & -\frac{1}{3} & 0 & -\frac{2}{3} \\ -8 & 2 & -1 & 6 \\ 0 & 2 & 3 & -2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & -\frac{1}{3} & 0 & -\frac{2}{3} \\ 0 & -\frac{2}{3} & -1 & \frac{2}{3} \\ 0 & 2 & 3 & -2 \end{array} \right] \\ &\sim \left[ \begin{array}{ccc|c} 1 & -\frac{1}{3} & 0 & -\frac{2}{3} \\ 0 & 1 & \frac{3}{2} & -1 \\ 0 & 2 & 3 & -2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & \frac{1}{2} & -1 \\ 0 & 1 & \frac{3}{2} & -1 \\ 0 & 0 & 0 & 0 \end{array} \right] \end{aligned}$$

□

<b>Standard E3.</b>	Mark:
---------------------	-------

Solve the following linear system.

$$\begin{aligned} 3x + 2y + z &= 7 \\ x + y + z &= 1 \\ -2x + 3z &= -11 \end{aligned}$$

**Solution:** Let  $A = \left[ \begin{array}{ccc|c} 3 & 2 & 1 & 7 \\ 1 & 1 & 1 & 1 \\ -2 & 0 & 3 & 11 \end{array} \right]$ , so  $\text{RREF } A = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 4 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & -1 \end{array} \right]$ . It follows that the system has exactly one solution:  $\begin{bmatrix} 4 & -2 & -1 \end{bmatrix}$

□

<b>Standard E4.</b>	Mark:
---------------------	-------

Find a basis for the solution set to the homogeneous system of equations

$$\begin{aligned} 2x_1 + 3x_2 - 5x_3 + 14x_4 &= 0 \\ x_1 + x_2 - x_3 + 5x_4 &= 0 \end{aligned}$$

**Solution:** Let  $A = \left[ \begin{array}{cccc|c} 2 & 3 & -5 & 14 & 0 \\ 1 & 1 & -1 & 5 & 0 \end{array} \right]$ , so  $\text{RREF } A = \left[ \begin{array}{cccc|c} 1 & 0 & 2 & 1 & 1 \\ 0 & 1 & -3 & 4 & 2 \end{array} \right]$ . It follows that the basis for the solution set is given by  $\left\{ \begin{bmatrix} -2 \\ 3 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ -4 \\ 0 \\ 1 \end{bmatrix} \right\}$ .

□

<b>Standard V1.</b>	Mark:
---------------------	-------

Let  $V$  be the set of all polynomials with the operations, for any  $f, g \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned} f \oplus g &= f' + g' \\ c \odot f &= cf' \end{aligned}$$

(here  $f'$  denotes the derivative of  $f$ ).

- (a) Show that scalar multiplication **distributes scalars** over vector addition:  $c \odot (f \oplus g) = c \odot f \oplus c \odot g$ .
- (b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $f, g \in \mathcal{P}$ , and let  $c \in \mathbb{R}$ .

$$c \odot (f \oplus g) = c \odot (f' + g') = c(f' + g')' = cf'' + cg'' = cf' \oplus cg' = c \odot f \oplus c \odot g.$$

However, this is not a vector space, as there is no zero vector. Additionally,  $1 \odot f \neq f$  for any nonzero polynomial  $f$ .

□

<b>Standard V2.</b>	Mark:
---------------------	-------

Determine if  $\begin{bmatrix} 4 \\ -1 \\ 6 \\ -7 \end{bmatrix}$  belongs to the span of the set  $\left\{ \begin{bmatrix} 2 \\ 0 \\ -1 \\ 5 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 4 \\ 3 \end{bmatrix} \right\}$ .

**Solution:** Since

$$\text{RREF} \left( \left( \begin{array}{cc|c} 2 & 4 & 4 \\ 0 & -1 & -1 \\ -1 & 4 & 6 \\ 5 & 3 & -7 \end{array} \right) \right) = \left[ \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array} \right]$$

contains the contradiction  $0 = 1$ ,  $\begin{bmatrix} 4 \\ -1 \\ 6 \\ -7 \end{bmatrix}$  is not a linear combination of the three vectors.

□

<b>Standard V3.</b>	Mark:
---------------------	-------

Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \left( \begin{array}{cccc} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{array} \right) \right) = \left[ \begin{array}{cccc} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Let  $W$  be the set of all polynomials of even degree. Determine if  $W$  is a subspace of the vector space of all polynomials.

**Solution:**  $W$  is closed under scalar multiplication, but not under addition. For example,  $x - x^2$  and  $x^2$  are both in  $W$ , but  $(x - x^2) + (x^2) = x \notin W$ .

□

<b>Standard S1.</b>	Mark:
---------------------	-------

Determine if the set of polynomials  $\{x^2 + x, x^2 + 2x - 1, x^2 + 3x - 2\}$  is linearly dependent or linearly independent

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 3 \\ 1 & -1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

Since there is a nonpivot column, the set is linearly dependent.

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

<b>Standard S3.</b>	Mark:
---------------------	-------

Let  $W$  be the subspace of  $\mathcal{P}_3$  given by

$W = \text{span}(\{x^3 + x^2 + 2x + 1, 3x^3 + 3x^2 + 6x + 3, 3x^3 - x^2 + 3x - 2, 7x^3 - x^2 + 8x - 3\})$ . Find a basis for  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then a basis is  $\{x^3 + x^2 + 2x + 1, 3x^3 - x^2 + 3x - 2\}$ .

□

<b>Standard S4.</b>	Mark:
---------------------	-------

Let  $W = \text{span} \left( \left( \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix} \right) \right)$ . Find the dimension of  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This has two pivot columns, so  $W$  has dimension 2.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--