

Name: \_\_\_\_\_

# **MASTERY QUIZ DAY 19**

Math 237 – Linear Algebra

## **Version 5**

Fall 2017

Show all work and justify all of your answers. Answers without work or sufficient reasoning will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**A3.** Determine if the following linear maps are injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  given by  $S \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} 3x + 2y \\ x - y \\ x + 4y \end{bmatrix}$

(b)  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $T \left( \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} x + y + z \\ 2y + 3z \\ x - y - 2z \end{bmatrix}$

**Solution:**

(a)

$$\text{RREF} \left( \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 3 \\ 1 & -1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

Since there is a nonpivot column,  $T$  is not injective. Since there is a zero row,  $T$  is not surjective.

(b)

$$\text{RREF} \left( \begin{bmatrix} 3 & 2 \\ 1 & -1 \\ 1 & 4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Since all columns are pivot columns,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

□

**A4.** Let  $T : \mathcal{P}^3 \rightarrow \mathcal{P}^3$  be the linear transformation given by

$$T(ax^3 + bx^2 + cx + d) = (a + 3b + 3c + 7d)x^3 + (a + 3b - c - d)x^2 + (2a + 6b + 3c + 8d)x + (a + 3b - 2c - 3d)$$

Compute the kernel and image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the kernel is

$$\ker(T) = \{(-3a - b)x^3 + ax^2 + (-2b)x + b \mid a, b \in \mathbb{R}\} = \text{span}(\{3x^3 - x^2, x^3 + 2x - 1\})$$

and the image is

$$\text{Im}(T) = \text{span}(\{x^3 + x^2 + 2x + 1, 3x^3 + 3x^2 + 6x + 3, 3x^3 - x^2 + 3x - 2, 7x^3 - x^2 + 8x - 3\}) = \text{span}(\{x^3 + x^2 + 2x + 1, 3x^3 + 3x^2 + 6x + 3\})$$

□

**S2:**

**A3:**

**A4:**