

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 1**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Does  $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$ ?

**Solution:** Since

$$\text{RREF} \begin{bmatrix} 2 & 3 & 1 & -4 \\ -1 & 12 & 2 & 2 \\ 4 & -9 & 3 & -8 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

lacks a zero row, the vectors span  $\mathbb{R}^3$ .

□

**V4.** Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 3 & -1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ -1 & 8 \end{bmatrix}, \begin{bmatrix} -1 & 3 \\ 0 & 4 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^{2 \times 2}$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 3 & 2 & 1 & -1 \\ -1 & 0 & 4 & 3 \\ 2 & 2 & -1 & 0 \\ 3 & 4 & 8 & 4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 2**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Does  $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 3 \\ 5 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 5 \\ 1 \\ -3 \end{bmatrix} \right\} = \mathbb{R}^5$ ?

**Solution:** Since there are only three vectors, they cannot span  $\mathbb{R}^5$ .

□

**V4.** Let  $W$  be the set of all complex numbers that are purely real (i.e of the form  $a + 0i$ ) or purely imaginary (i.e. of the form  $0 + bi$ ). Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** No, because 1 is purely real and  $i$  is purely imaginary, but the linear combination  $1 + i$  is neither.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 3**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span  $\mathbb{R}^4$ .

□

**V4.** Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 3 & -1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ -1 & 8 \end{bmatrix}, \begin{bmatrix} -1 & 3 \\ 0 & 4 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^{2 \times 2}$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 3 & 2 & 1 & -1 \\ -1 & 0 & 4 & 3 \\ 2 & 2 & -1 & 0 \\ 3 & 4 & 8 & 4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

# MASTERY QUIZ DAY 17

Math 237 – Linear Algebra

## Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Determine if the vectors  $\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there are zero rows, they do not span. Alternatively, by inspection  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ , so the set is linearly dependent, so it spans a subspace of dimension at most 3, therefore it does not span  $\mathbb{R}^4$ .

□

**V4.** Determine if  $\left\{ \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^4$ .

**Solution:** It is closed under addition and scalar multiplication, so it is a subspace. Alternatively, it is the image of the linear transformation from  $\mathbb{R}^3 \rightarrow \mathbb{R}^4$  given by

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix}.$$

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 3 & -1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ -1 & 8 \end{bmatrix}, \begin{bmatrix} -1 & 3 \\ 0 & 4 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^{2 \times 2}$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 3 & 2 & 1 & -1 \\ -1 & 0 & 4 & 3 \\ 2 & 2 & -1 & 0 \\ 3 & 4 & 8 & 4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 17**

Math 237 – Linear Algebra

**Version 5**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Does  $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$ ?

**Solution:** Since

$$\text{RREF} \begin{bmatrix} 2 & 3 & 1 & -4 \\ -1 & 12 & 2 & 2 \\ 4 & -9 & 3 & -8 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

lacks a zero row, the vectors span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all  $\mathbb{R}^3$  vectors  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  satisfying  $x + y + z = 0$  (this forms a plane). Determine if  $W$  is a subspace of  $\mathbb{R}^3$ .

**Solution:** Yes, because  $z = -x - y$  and  $a \begin{bmatrix} x_1 \\ y_1 \\ -x_1 - y_1 \end{bmatrix} + b \begin{bmatrix} x_2 \\ y_2 \\ -x_2 - y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + bx_2 \\ ay_1 + by_2 \\ -(ax_1 + bx_2) - (ay_1 + by_2) \end{bmatrix}$ .

Alternately, yes because  $W$  is isomorphic to  $\mathbb{R}^2$ .

□

**S2.** Determine if the set  $\{2x^2 - x + 3, 2x^2 + 2, -x^2 + 4x + 1\}$  is a basis of  $\mathcal{P}^2$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 2 & 2 & -1 \\ -1 & 0 & 4 \\ 3 & 2 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

# MASTERY QUIZ DAY 17

Math 237 – Linear Algebra

## Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V3.** Does  $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$ ?

**Solution:** Since

$$\text{RREF} \begin{bmatrix} 2 & 3 & 1 & -4 \\ -1 & 12 & 2 & 2 \\ 4 & -9 & 3 & -8 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

lacks a zero row, the vectors span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all  $\mathbb{R}^3$  vectors  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  satisfying  $x + y + z = 0$  (this forms a plane). Determine if  $W$  is a subspace of  $\mathbb{R}^3$ .

**Solution:** Yes, because  $z = -x - y$  and  $a \begin{bmatrix} x_1 \\ y_1 \\ -x_1 - y_1 \end{bmatrix} + b \begin{bmatrix} x_2 \\ y_2 \\ -x_2 - y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + bx_2 \\ ay_1 + by_2 \\ -(ax_1 + bx_2) - (ay_1 + by_2) \end{bmatrix}$ .

Alternately, yes because  $W$  is isomorphic to  $\mathbb{R}^2$ .

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**V3:**

**V4:**

**S2:**