

Name: _____

MASTERY QUIZ DAY 8

Math 237 – Linear Algebra

Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write an augmented matrix corresponding to the following system of linear equations.

$$\begin{aligned}x_1 + 4x_3 &= 1 \\x_2 - x_3 &= 7 \\x_1 - x_2 + 3x_4 &= -1\end{aligned}$$

Solution:

$$\left[\begin{array}{cccc|c} 1 & 0 & 4 & 0 & 1 \\ 0 & 1 & -1 & 0 & 7 \\ 1 & -1 & 0 & 3 & -1 \end{array} \right]$$

□

E3. Solve the system of equations

$$\begin{aligned}x + 3y + 3z + 7w &= 0 \\x + 3y - z - w &= 0 \\2x + 6y + 3z + 8w &= 0 \\x + 3y - 2z - 3w &= 0\end{aligned}$$

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the solution set is

$$\left\{ \begin{bmatrix} -3a - b \\ a \\ -2b \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\} = \text{span} \left(\left\{ \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \\ -1 \end{bmatrix} \right\} \right)$$

□

E4. Find a basis for the solution set of the system ...

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

Determine if V is a vector space or not.

Solution:

1) Since real addition is associative, \oplus is associative.

2) Since real addition is commutative, \oplus is commutative.

3) $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .

5)

$$\begin{aligned}
 c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\
 &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\
 &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\
 &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\
 &= (cd) \odot (x_1, y_1)
 \end{aligned}$$

6) $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$

7)

$$\begin{aligned}
 c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\
 &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\
 &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\
 &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\
 &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)
 \end{aligned}$$

8)

$$\begin{aligned}
 (c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\
 &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\
 &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)
 \end{aligned}$$

□

E1:

E3:

E4:

V1: