

Name: \_\_\_\_\_

## MASTERY QUIZ DAY 12

Math 237 – Linear Algebra

### Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all points on the line  $x + y = 2$  with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

Determine if  $V$  is a vector space or not.

### Solution:

- 1) Since real addition is associative,  $\oplus$  is associative.
- 2) Since real addition is commutative,  $\oplus$  is commutative.
- 3)  $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$ , so  $(1, 1)$  is an additive identity element.
- 4)  $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$ , so  $(2 - x_1, 2 - y_1)$  is the additive inverse of  $(x_1, y_1)$ .
- 5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

6)  $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$

7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

□

**V3.** Determine if the vectors  $\begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, the vectors do not span  $\mathbb{R}^4$ .

□

**V4.** Determine if  $\left\{ \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^4$ .

**Solution:** It is closed under addition and scalar multiplication, so it is a subspace. Alternatively, it is the image of the linear transformation from  $\mathbb{R}^3 \rightarrow \mathbb{R}^4$  given by

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix}.$$

□

**V1:**

**V3:**

**V4:**