

Name:
J#:
Date:

Dr. Clontz

SEMIFINAL

Math 237 – Linear Algebra

Version 2

Fall 2017

Work any problems you wish on the provided answer sheets; take care to label the standard for each response. Show all work and justify all of your answers. Answers without work or sufficient reasoning will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{cccc|c} 3 & -1 & 0 & 1 & 5 \\ -1 & 9 & 1 & -7 & 0 \\ 1 & 0 & -1 & 0 & -3 \end{array} \right]$$

E2. Put the following matrix in reduced row echelon form.

$$\left[\begin{array}{ccc} 3 & -1 & 0 \\ -1 & 0 & -1 \\ -1 & 1 & 2 \\ 0 & 2 & 6 \end{array} \right]$$

E3. Find the solution set for the following system of linear equations.

$$2x_1 + 3x_2 - 5x_3 + 14x_4 = 8$$

$$x_1 + x_2 - x_3 + 5x_4 = 3$$

E4. Find a basis for the solution set of the system of equations

$$x + 3y + 3z + 7w = 0$$

$$x + 3y - z - w = 0$$

$$2x + 6y + 3z + 8w = 0$$

$$x + 3y - 2z - 3w = 0$$

V1. Let V be the set of all pairs of real numbers with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \odot (x_1, y_1) = (c^2 x_1, c^3 y_1)$$

(a) Show that scalar multiplication **distributes scalars** over vector addition:

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2).$$

(b) Determine if V is a vector space or not. Justify your answer.

V2. Determine if $\begin{bmatrix} 0 \\ -1 \\ 6 \\ -7 \end{bmatrix}$ belongs to the span of the set $\left\{ \begin{bmatrix} 2 \\ 0 \\ -1 \\ 5 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 4 \\ 3 \end{bmatrix} \right\}$.

V3. Does $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ -3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$?

V4. Let W be the set of all polynomials of the form $ax^3 + bx$. Determine if W is a subspace of \mathcal{P}^3 .

S1. Determine if the set of vectors $\left\{ \begin{bmatrix} -3 \\ 8 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 3 \end{bmatrix} \right\}$ is linearly dependent or linearly independent

S2. Determine if the set $\{x^3 - x, x^2 + x + 1, x^3 - x^2 + 2, 2x^2 - 1\}$ is a basis of \mathcal{P}^3 .

S3. Let $W = \text{span} \left(\left\{ \begin{bmatrix} 1 \\ -1 \\ 3 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ -7 \end{bmatrix} \right\} \right)$. Find a basis of W .

S4. Let W be the subspace of $\mathbb{R}^{2 \times 2}$ given by $W = \text{span} \left(\left\{ \begin{bmatrix} 2 & 0 \\ -2 & 0 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 3 & 6 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \right\} \right)$. Compute the dimension of W .

A1. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be the linear transformation given by

$$T \left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} -3x + y \\ -8x + 2y - z \\ 2y + 3z \\ 0 \end{bmatrix}.$$

Write the matrix for T with respect to the standard bases of \mathbb{R}^3 and \mathbb{R}^4 .

A2. Determine if $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} e^x \\ e^y \end{bmatrix}$ is a linear transformation.

A3. Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a) $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by the standard matrix $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.

(b) $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ given by the standard matrix $\begin{bmatrix} 2 & 3 & -1 & -2 \\ 0 & 1 & 3 & 1 \\ 2 & 1 & -7 & -4 \end{bmatrix}$

A4. Let $T : \mathbb{R}^{2 \times 3} \rightarrow \mathbb{R}^3$ be the linear map given by $T \left(\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix} \right) = \begin{bmatrix} a + x \\ b + y \\ c + z \end{bmatrix}$. Compute a basis for the kernel and a basis for the image of T .

M1. Let

$$A = \begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix} \qquad B = \begin{bmatrix} 3 & 1 & 0 \end{bmatrix} \qquad C = \begin{bmatrix} 3 & -1 & 4 \\ 1 & 0 & 2 \end{bmatrix}$$

Exactly one of the six products AB , AC , BA , BC , CA , CB can be computed. Determine which one, and compute it.

M2. Determine if the matrix $\begin{bmatrix} 2 & 1 & 0 & 3 \\ 1 & -1 & 3 & 1 \\ 3 & 2 & -1 & 7 \\ 4 & 1 & 2 & 0 \end{bmatrix}$ is invertible.

M3. Find the inverse of the matrix $\begin{bmatrix} 1 & -4 & 5 \\ -5 & 24 & -28 \\ 1 & -5 & 6 \end{bmatrix}$.

G1. Compute the determinant of the matrix

$$\begin{bmatrix} 1 & 3 & 2 & 4 \\ -2 & 3 & -1 & 1 \\ 5 & 0 & -4 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}.$$

G2. Compute the eigenvalues, along with their algebraic multiplicities, of the matrix $\begin{bmatrix} 9 & -3 & 2 \\ 19 & -6 & 5 \\ -11 & 4 & -2 \end{bmatrix}$.

G3. Compute the eigenspace of the eigenvalue -1 in the matrix $\begin{bmatrix} 4 & -2 & -1 \\ 15 & -7 & -3 \\ -5 & 2 & 0 \end{bmatrix}$.

G4. Compute the geometric multiplicity of the eigenvalue 2 in the matrix $\begin{bmatrix} -1 & 1 & 0 \\ -9 & 5 & 0 \\ 15 & -5 & 2 \end{bmatrix}$.

Standard: _____



Standard: _____



Standard: _____



Standard: _____



Standard: _____



Standard: _____

