

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all points on the parabola $y = x^2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 - x_2, y_1 + y_2 - 2x_1x_2)$$

$$c \odot (x_1, y_1) = (cx_1, c^2y_1)$$

(a) Show that scalar multiplication **distributes scalars** over vector addition:

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2).$$

(b) Determine if V is a vector space or not. Justify your answer.

Solution:

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1 - x_2, y_1 + y_2 - 2x_1x_2) = (c(x_1 - x_2), c^2(y_1 + y_2 - 2x_1x_2))$$

$$c \odot (x_1, y_1) \oplus c \odot (x_2, y_2) = (cx_1, c^2y_1) \oplus (cx_2, c^2y_2) = (cx_1 - cx_2, c^2y_1 + c^2y_2 - 2(cx_1)(cx_2))$$

Not a vector space as addition is not commutative.

□

V3. Determine if the vectors $\begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$, and $\begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}$ span \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, the vectors do not span \mathbb{R}^4 .

□

V4. Let W be the set of all \mathbb{R}^3 vectors $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ satisfying $x + y + z = 1$ (this forms a plane). Determine if W is a subspace of \mathbb{R}^3 .

Solution: No, because $\mathbf{0}$ does not belong to W .

□

S2. Determine if the set $\{x^3 - 3x^2 + 2x + 2, -x^3 + 4x^2 - x + 1, -x^3 + 2x + 1, 3x^2 + 3x + 9\}$ is a basis of \mathcal{P}^3 or not.

Solution:

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

V1:

V3:

V4:

S2: