

Name: \_\_\_\_\_

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all pairs of real numbers with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V$ ,  $c \in \mathbb{R}$ ,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \odot (x_1, y_1) = (0, cy_1)$$

(a) Show that scalar multiplication **distributes vectors** over scalar addition:

$$(c + d) \odot (x, y) = c \odot (x, y) \oplus d \odot (x, y).$$

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $(x_1, y_1) \in V$ , and let  $c, d \in \mathbb{R}$ . Then

$$(c + d) \odot (x_1, y_1) = (0, (c + d)y_1) = (0, cy_1) \oplus (0, dy_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

However,  $V$  is not a vector space, as  $1 \odot (x_1, y_1) = (0, y_1) \neq (x_1, y_1)$ .

□

**V3.** Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all  $\mathbb{R}^3$  vectors  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  satisfying  $x + y + z = 1$  (this forms a plane). Determine if  $W$  is a subspace of  $\mathbb{R}^3$ .

**Solution:** No, because  $\mathbf{0}$  does not belong to  $W$ .

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**