

Name: \_\_\_\_\_

### MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

#### Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all real numbers with the operations, for any  $x, y \in V$ ,  $c \in \mathbb{R}$ ,

$$x \oplus y = \sqrt{x^2 + y^2}$$

$$c \odot x = cx$$

(a) Show that the vector **addition**  $\oplus$  is **associative**:  $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $x, y, z \in \mathbb{R}$ . Then

$$\begin{aligned}(x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\&= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\&= \sqrt{x^2 + y^2 + z^2} \\&= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\&= x \oplus \sqrt{y^2 + z^2} \\&= x \oplus (y \oplus z)\end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

**V3.** Determine if the vectors  $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span  $\mathbb{R}^4$ .

□

**V4.** Let  $W$  be the set of all polynomials of even degree. Determine if  $W$  is a subspace of the vector space of all polynomials.

**Solution:**  $W$  is closed under scalar multiplication, but not under addition. For example,  $x - x^2$  and  $x^2$  are both in  $W$ , but  $(x - x^2) + (x^2) = x \notin W$ .

□

**S2.** Determine if the set  $\{x^3 - 3x^2 + 2x + 2, -x^3 + 4x^2 - x + 1, -x^3 + 2x + 1, 3x^2 + 3x + 9\}$  is a basis of  $\mathcal{P}^3$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all pairs of real numbers with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V$ ,  $c \in \mathbb{R}$ ,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \odot (x_1, y_1) = (0, cy_1)$$

(a) Show that scalar multiplication **distributes vectors** over scalar addition:

$$(c + d) \odot (x, y) = c \odot (x, y) \oplus d \odot (x, y).$$

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $(x_1, y_1) \in V$ , and let  $c, d \in \mathbb{R}$ . Then

$$(c + d) \odot (x_1, y_1) = (0, (c + d)y_1) = (0, cy_1) \oplus (0, dy_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

However,  $V$  is not a vector space, as  $1 \odot (x_1, y_1) = (0, y_1) \neq (x_1, y_1)$ .

□

**V3.** Does  $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$ ?

**Solution:** Since

$$\text{RREF} \begin{bmatrix} 2 & 3 & 1 & -4 \\ -1 & 12 & 2 & 2 \\ 4 & -9 & 3 & -8 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

lacks a zero row, the vectors span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all complex numbers that are purely real (i.e of the form  $a + 0i$ ) or purely imaginary (i.e. of the form  $0 + bi$ ). Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** No, because 1 is purely real and  $i$  is purely imaginary, but the linear combination  $1 + i$  is neither.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 0 \\ -1 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & 0 & -1 & 0 \\ 1 & 2 & 0 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

### MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

#### Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all real numbers together with the operations  $\oplus$  and  $\odot$  defined by, for any  $x, y \in V$  and  $c \in \mathbb{R}$ ,

$$\begin{aligned}x \oplus y &= x + y - 3 \\c \odot x &= cx - 3(c - 1)\end{aligned}$$

- (a) Show that **scalar multiplication** is **associative**:  $a \odot (b \odot x) = (ab) \odot x$ .
- (b) Determine if  $V$  is a vector space or not. Justify your answer

**Solution:** Let  $x, y \in V$ ,  $c, d \in \mathbb{R}$ . To show associativity:

$$\begin{aligned}c \odot (d \odot x) &= c \odot (dx - 3(d - 1)) \\&= c(dx - 3(d - 1)) - 3(c - 1) \\&= cdx - 3(cd - 1) \\&= (cd) \odot x\end{aligned}$$

We verify the remaining 7 properties to see that  $V$  is a vector space.

- 1) Real addition is associative, so  $\oplus$  is associative.
- 2)  $x \oplus 3 = x + 3 - 3 = x$ , so 3 is the additive identity.
- 3)  $x \oplus (6 - x) = x + (6 - x) - 3 = 3$ , so  $6 - x$  is the additive inverse of  $x$ .
- 4) Real addition is commutative, so  $\oplus$  is commutative.
- 5) Associativity shown above
- 6)  $1 \odot x = x - 3(1 - 1) = x$
- 7)

$$\begin{aligned}c \odot (x \oplus y) &= c \odot (x + y - 3) \\&= c(x + y - 3) - 3(c - 1) \\&= cx - 3(c - 1) + cy - 3(c - 1) - 3 \\&= (c \odot x) \oplus (c \odot y)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot x &= (c + d)x - 3(c + d - 1) \\&= cx - 3(c - 1) + dx - 3(d - 1) - 3 \\&= (c \odot x) \oplus (d \odot x)\end{aligned}$$

Therefore  $V$  is a vector space.

□

**V3.** Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all  $\mathbb{R}^3$  vectors  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  satisfying  $x + y + z = 1$  (this forms a plane). Determine if  $W$  is a subspace of  $\mathbb{R}^3$ .

**Solution:** No, because  $\mathbf{0}$  does not belong to  $W$ .

□

**S2.** Determine if the set  $\{x^3 - 3x^2 + 2x + 2, -x^3 + 4x^2 - x + 1, -x^3 + 2x + 1, 3x^2 + 3x + 9\}$  is a basis of  $\mathcal{P}^3$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**V1:** ☐

**V3:** ☐

**V4:** ☐

**S2:** ☐

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 14**

Math 237 – Linear Algebra

**Version 4**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all real numbers with the operations, for any  $x, y \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned}x \oplus y &= \sqrt{x^2 + y^2} \\ c \odot x &= cx\end{aligned}$$

(a) Show that the vector **addition**  $\oplus$  is **associative**:  $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $x, y, z \in \mathbb{R}$ . Then

$$\begin{aligned}(x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\ &= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\ &= \sqrt{x^2 + y^2 + z^2} \\ &= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\ &= x \oplus \sqrt{y^2 + z^2} \\ &= x \oplus (y \oplus z)\end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

**V3.** Determine if the vectors  $\begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row, the vectors do not span  $\mathbb{R}^4$ .

□

**V4.** Let  $W$  be the set of all  $\mathbb{R}^3$  vectors  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  satisfying  $x + y + z = 1$  (this forms a plane). Determine if  $W$  is a subspace of  $\mathbb{R}^3$ .

**Solution:** No, because  $\mathbf{0}$  does not belong to  $W$ .

□

**S2.** Determine if the set  $\{x^2 + x - 1, 3x^2 - x + 1, 2x - 2\}$  is a basis of  $\mathcal{P}_2$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 14**

Math 237 – Linear Algebra

**Version 5**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all pairs of real numbers with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2, y_1 + y_2) \\ c \odot (x_1, y_1) &= (0, cy_1)\end{aligned}$$

(a) Show that scalar multiplication **distributes vectors** over scalar addition:

$$(c + d) \odot (x, y) = c \odot (x, y) \oplus d \odot (x, y).$$

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $(x_1, y_1) \in V$ , and let  $c, d \in \mathbb{R}$ . Then

$$(c + d) \odot (x_1, y_1) = (0, (c + d)y_1) = (0, cy_1) \oplus (0, dy_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

However,  $V$  is not a vector space, as  $1 \odot (x_1, y_1) = (0, y_1) \neq (x_1, y_1)$ .

□

**V3.** Determine if the vectors  $\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there are zero rows, they do not span. Alternatively, by inspection  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ , so the set is linearly dependent, so it spans a subspace of dimension at most 3, therefore it does not span  $\mathbb{R}^4$ .

□

**V4.** Let  $W$  be the set of all polynomials of the form  $ax^3 + bx$ . Determine if  $W$  is a subspace of  $\mathcal{P}^3$ .

**Solution:** Yes because  $s(a_1x^3 + b_1x) + t(a_2x^3 + b_2x) = (sa_1 + ta_2)x^3 + (sb_1 + tb_2)x$  also belongs to  $W$ . Alternately, yes because  $W$  is isomorphic to  $\mathbb{R}^2$ .

□

**S2.** Determine if the set  $\{x^3 - x, x^2 + x + 1, x^3 - x^2 + 2, 2x^2 - 1\}$  is a basis of  $\mathcal{P}_3$



**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 \\ -1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix is not the identity matrix, it is not a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 14**

Math 237 – Linear Algebra

**Version 6**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**V1.** Let  $V$  be the set of all real numbers together with the operations  $\oplus$  and  $\odot$  defined by, for any  $x, y \in V$  and  $c \in \mathbb{R}$ ,

$$\begin{aligned}x \oplus y &= x + y - 3 \\c \odot x &= cx - 3(c - 1)\end{aligned}$$

- (a) Show that **scalar multiplication** is **associative**:  $a \odot (b \odot x) = (ab) \odot x$ .
- (b) Determine if  $V$  is a vector space or not. Justify your answer

**Solution:** Let  $x, y \in V$ ,  $c, d \in \mathbb{R}$ . To show associativity:

$$\begin{aligned}c \odot (d \odot x) &= c \odot (dx - 3(d - 1)) \\&= c(dx - 3(d - 1)) - 3(c - 1) \\&= cdx - 3(cd - 1) \\&= (cd) \odot x\end{aligned}$$

We verify the remaining 7 properties to see that  $V$  is a vector space.

- 1) Real addition is associative, so  $\oplus$  is associative.
- 2)  $x \oplus 3 = x + 3 - 3 = x$ , so 3 is the additive identity.
- 3)  $x \oplus (6 - x) = x + (6 - x) - 3 = 3$ , so  $6 - x$  is the additive inverse of  $x$ .
- 4) Real addition is commutative, so  $\oplus$  is commutative.
- 5) Associativity shown above
- 6)  $1 \odot x = x - 3(1 - 1) = x$
- 7)

$$\begin{aligned}c \odot (x \oplus y) &= c \odot (x + y - 3) \\&= c(x + y - 3) - 3(c - 1) \\&= cx - 3(c - 1) + cy - 3(c - 1) - 3 \\&= (c \odot x) \oplus (c \odot y)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot x &= (c + d)x - 3(c + d - 1) \\&= cx - 3(c - 1) + dx - 3(d - 1) - 3 \\&= (c \odot x) \oplus (d \odot x)\end{aligned}$$

Therefore  $V$  is a vector space.

□

**V3.** Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ .

□

**V4.** Let  $W$  be the set of all complex numbers  $a + bi$  satisfying  $a = 2b$ . Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** Yes, because  $c(2b_1 + b_1i) + d(2b_2 + b_2i) = 2(cb_1 + db_2) + (cb_1 + db_2)i$  belongs to  $W$ . Alternately, yes because  $W$  is isomorphic to  $\mathbb{R}$ .

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**V1:**

**V3:**

**V4:**

**S2:**