

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 28

Math 237 – Linear Algebra

### Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard M1.</b>	Mark:
---------------------	-------

Let

$$A = \begin{bmatrix} 0 & 0 & -1 & -1 \\ 1 & 3 & 7 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 0 & 1 & 7 & 7 \\ -1 & -2 & 0 & 4 \\ 0 & 0 & 1 & 5 \end{bmatrix}$$

$$C = \begin{bmatrix} 3 & 2 \\ 0 & 1 \\ -2 & -1 \end{bmatrix}$$

Exactly one of the six products  $AB$ ,  $AC$ ,  $BA$ ,  $BC$ ,  $CA$ ,  $CB$  can be computed. Determine which one, and compute it.

**Solution:**  $CA$  is the only one that can be computed, and

$$CA = \begin{bmatrix} 2 & 6 & 11 & 1 \\ 1 & 3 & 7 & 2 \\ -1 & -3 & -5 & 0 \end{bmatrix}$$

□

<b>Standard M2.</b>	Mark:
---------------------	-------

Determine if the matrix  $\begin{bmatrix} 3 & -1 & 0 & 4 \\ 2 & 1 & 1 & -1 \\ 0 & 1 & 1 & 3 \\ 1 & -2 & 0 & 0 \end{bmatrix}$  is invertible.

**Solution:** This matrix is row equivalent to the identity matrix, so it is invertible.

□

<b>Standard M3.</b>	Mark:
---------------------	-------

Find the inverse of the matrix  $\begin{bmatrix} 6 & 0 & 1 \\ -14 & 3 & -4 \\ -23 & 4 & -6 \end{bmatrix}$ .

**Solution:**

$$\begin{bmatrix} 6 & 0 & 1 \\ -14 & 3 & -4 \\ -23 & 4 & -6 \end{bmatrix}^{-1} = \begin{bmatrix} -2 & 4 & -3 \\ 8 & -13 & 10 \\ 13 & -24 & 18 \end{bmatrix}$$

□

<b>Standard G2.</b>	Mark:
---------------------	-------

Let  $A = \begin{bmatrix} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \end{bmatrix}$ . List the eigenvalues of  $A$  along with their algebraic multiplicities.

**Solution:**

$$\begin{aligned}
 \det(A - \lambda I) &= \det \begin{bmatrix} -3 - \lambda & 1 & 0 \\ -8 & 2 - \lambda & -1 \\ 0 & 2 & 3 - \lambda \end{bmatrix} \\
 &= (-3 - \lambda) \det \begin{bmatrix} 2 - \lambda & -1 \\ 2 & 3 - \lambda \end{bmatrix} - (1) \det \begin{bmatrix} -8 & -1 \\ 0 & 3 - \lambda \end{bmatrix} \\
 &= (-3 - \lambda) ((2 - \lambda)(3 - \lambda) + 2) - (-8(3 - \lambda)) \\
 &= (-3 - \lambda)(8 - 5\lambda + \lambda^2) + 24 - 8\lambda \\
 &= -\lambda^3 + 2\lambda^2 + 7\lambda - 24 + 24 - 8\lambda \\
 &= -\lambda^3 + 2\lambda^2 - \lambda \\
 &= -\lambda(\lambda^2 - 2\lambda + 1) \\
 &= -\lambda(\lambda - 1)^2
 \end{aligned}$$

So  $A$  has eigenvalues 0 (with multiplicity 1) and 1 (with algebraic multiplicity 2).

□

<b>Standard G3.</b>	Mark:
---------------------	-------

Compute the eigenspace associated to the eigenvalue 2 in the matrix  $\begin{bmatrix} -1 & 1 & 0 \\ -9 & 5 & 0 \\ 15 & -5 & 2 \end{bmatrix}$ .

**Solution:** The eigenspace is the solution space of the system  $(B - 2I)X = 0$ .

$$\text{RREF}(B - 2I) = \text{RREF} \left( \begin{bmatrix} -3 & 1 & 0 \\ -9 & 3 & 0 \\ 15 & -5 & 0 \end{bmatrix} \right) = \begin{bmatrix} 1 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

So the system simplifies to  $x - \frac{y}{3} = 0$ , or  $3x = y$ . Thus the eigenspace is

$$E_2 = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$$

□

<b>Additional Notes/Marks</b>	
-------------------------------	--