

| |
|-------|
| Name: |
| J#: |
| Date: |

Dr. Clontz

MASTERY QUIZ DAY 12

Math 237 – Linear Algebra

Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

| | |
|---------------------|-------|
| Standard V1. | Mark: |
|---------------------|-------|

Let V be the set of all real numbers together with the operations \oplus and \odot defined by, for any $x, y \in V$ and $c \in \mathbb{R}$,

$$\begin{aligned}x \oplus y &= x + y - 3 \\c \odot x &= cx - 3(c - 1)\end{aligned}$$

Determine if V is a vector space or not.

Solution: Let $x, y \in V$, $c, d \in \mathbb{R}$.

- 1) Real addition is associative, so \oplus is associative.
- 2) $x \oplus 3 = x + 3 - 3 = x$, so 3 is the additive identity.
- 3) $x \oplus (6 - x) = x + (6 - x) - 3 = 3$, so $6 - x$ is the additive inverse of x .
- 4) Real addition is commutative, so \oplus is commutative.
- 5)

$$\begin{aligned}c \odot (d \odot x) &= c \odot (dx - 3(d - 1)) \\&= c(dx - 3(d - 1)) - 3(c - 1) \\&= cdx - 3(cd - 1) \\&= (cd) \odot x\end{aligned}$$

$$6) 1 \odot x = x - 3(1 - 1) = x$$

7)

$$\begin{aligned}c \odot (x \oplus y) &= c \odot (x + y - 3) \\&= c(x + y - 3) - 3(c - 1) \\&= cx - 3(c - 1) + cy - 3(c - 1) - 3 \\&= (c \odot x) \oplus (c \odot y)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot x &= (c + d)x - 3(c + d - 1) \\&= cx - 3(c - 1) + dx - 3(d - 1) - 3 \\&= (c \odot x) \oplus (d \odot x)\end{aligned}$$

Therefore V is a vector space.

□

| | |
|---------------------|-------|
| Standard V3. | Mark: |
|---------------------|-------|

Determine if the vectors $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$, and $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$ span \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span \mathbb{R}^4 .

□

| | |
|---------------------|-------|
| Standard V4. | Mark: |
|---------------------|-------|

Determine if the set of all lattice points, i.e. $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$ is a subspace of \mathbb{R}^2 .

Solution: This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

| | |
|-------------------------------|--|
| Additional Notes/Marks | |
|-------------------------------|--|