

Name: \_\_\_\_\_

**MIDTERM EXAM**

Math 237 – Linear Algebra

**Version 6**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**E1.** Write an augmented matrix corresponding to the following system of linear equations.

$$\begin{aligned}x_1 + 3x_2 - 4x_3 + x_4 &= 5 \\ 3x_1 + 9x_2 + x_3 - 7x_4 &= 0 \\ x_1 - x_3 + x_4 &= 1\end{aligned}$$

**Solution:**

$$\left[ \begin{array}{cccc|c} 1 & 3 & -4 & 1 & 5 \\ 3 & 9 & 1 & -7 & 0 \\ 1 & 0 & -1 & 1 & 1 \end{array} \right]$$

□

**E2.** Find RREF  $A$ , where

$$A = \left[ \begin{array}{cccc|c} 2 & 2 & 1 & 2 & -1 \\ 1 & 1 & 2 & 4 & 5 \\ 3 & 3 & -1 & -2 & 1 \end{array} \right]$$

**Solution:**

$$\text{RREF } A = \left[ \begin{array}{cccc|c} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$$

□

**E3.** Find the solution set for the following system of linear equations.

$$\begin{aligned}2x_1 + 3x_2 - 5x_3 + 14x_4 &= 8 \\ x_1 + x_2 - x_3 + 5x_4 &= 3\end{aligned}$$

**Solution:** Let  $A = \left[ \begin{array}{cccc|c} 2 & 3 & -5 & 14 & 8 \\ 1 & 1 & -1 & 5 & 3 \end{array} \right]$ , so  $\text{RREF } A = \left[ \begin{array}{cccc|c} 1 & 0 & 2 & 1 & 1 \\ 0 & 1 & -3 & 4 & 2 \end{array} \right]$ . It follows that the solution set is given by  $\begin{bmatrix} 1 - 2a - b \\ 2 + 3a - 4b \\ a \\ b \end{bmatrix}$  for all real numbers  $a, b$ .

□

**E4.** Find a basis for the solution set to the homogeneous system of equations given by

$$\begin{aligned}2x_1 - 2x_2 + 6x_3 - x_4 &= 0 \\ 3x_1 + 6x_3 + x_4 &= 0 \\ -4x_1 + x_2 - 9x_3 + 2x_4 &= 0\end{aligned}$$

**Solution:** Let  $A = \left[ \begin{array}{cccc|c} 2 & -2 & 6 & -1 & 0 \\ 3 & 0 & 6 & 1 & 0 \\ -4 & 1 & -9 & 2 & 0 \end{array} \right]$ , so  $\text{RREF } A = \left[ \begin{array}{cccc|c} 1 & 0 & 2 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$ . It follows that the basis for the solution set is given by  $\left\{ \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right\}$ .

□

**V1.** Let  $V$  be the set of all points on the parabola  $y = x^2$  with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V, c \in \mathbb{R}$ ,

$$\begin{aligned} (x_1, y_1) \oplus (x_2, y_2) &= (x_1 - x_2, y_1 + y_2 - 2x_1x_2) \\ c \odot (x_1, y_1) &= (cx_1, c^2y_1) \end{aligned}$$

(a) Show that scalar multiplication **distributes scalars** over vector addition:  
 $c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:**

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1 - x_2, y_1 + y_2 - 2x_1x_2) = (c(x_1 - x_2), c^2(y_1 + y_2 - 2x_1x_2))$$

$$c \odot (x_1, y_1) \oplus c \odot (x_2, y_2) = (cx_1, c^2y_1) \oplus (cx_2, c^2y_2) = (cx_1 - cx_2, c^2y_1 + c^2y_2 - 2(cx_1)(cx_2))$$

Not a vector space as addition is not commutative.

□

**V2.** Determine if  $\begin{bmatrix} 0 \\ -1 \\ 2 \\ 6 \end{bmatrix}$  can be written as a linear combination of the vectors  $\begin{bmatrix} 3 \\ -1 \\ -1 \\ 0 \end{bmatrix}$  and  $\begin{bmatrix} -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$ .

**Solution:**

$$\text{RREF} \left( \left[ \begin{array}{cc|c} 3 & -1 & 0 \\ -1 & 0 & -1 \\ -1 & 1 & 2 \\ 0 & 2 & 6 \end{array} \right] \right) = \left[ \begin{array}{cc|c} 1 & 0 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Since this system has a solution,  $\begin{bmatrix} 0 \\ -1 \\ 2 \\ 6 \end{bmatrix}$  can be written as a linear combination of the vectors  $\begin{bmatrix} 3 \\ -1 \\ -1 \\ 0 \end{bmatrix}$  and

$\begin{bmatrix} -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$ , namely

$$\begin{bmatrix} 0 \\ -1 \\ 2 \\ 6 \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ -1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}.$$

□

**V3.** Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ . □

**V4.** Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace. □

**S1.** Determine if the set of matrices  $\left\{ \begin{bmatrix} 3 & -1 \\ 0 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 3 & -8 \\ 6 & 5 \end{bmatrix} \right\}$  is linearly dependent or linearly independent.

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 3 & 1 & 3 \\ -1 & 2 & -8 \\ 0 & -2 & 6 \\ 4 & 1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Since the reduced row echelon form has a nonpivot column, the vectors are linearly dependent. □

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis. □

**S3.** Let  $W = \text{span} \left( \left\{ \begin{bmatrix} -3 \\ -8 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 3 \end{bmatrix} \right\} \right)$ . Find a basis for  $W$ .

**Solution:** Let  $A = \begin{bmatrix} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \end{bmatrix}$ , and compute  $\text{RREF}(A) = \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$ . Since the first two columns are

pivot columns,  $\left\{ \begin{bmatrix} -3 \\ -8 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \right\}$  is a basis for  $W$ .

□

**S4.** Let  $W = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ -1 \\ 3 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ -7 \end{bmatrix} \right\} \right)$ . Compute the dimension of  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 2 & 3 & 1 \\ -1 & 0 & -1 & 1 \\ 3 & 1 & 4 & 1 \\ -3 & 1 & -2 & -7 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This has 3 pivot columns so  $\dim(W) = 3$ .

□

E1:

E2:

E3:

E4:

V1:

V2:

V3:

V4:

S1:

S2:

S3:

S4: