

Name: \_\_\_\_\_

**MASTERY QUIZ DAY 28**

Math 237 – Linear Algebra

**Version 6**

Fall 2017

Show all work and justify all of your answers. Answers without work or sufficient reasoning will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**M1.** Let

$$A = \begin{bmatrix} 0 & 0 & -1 & -1 \\ 1 & 3 & 7 & 2 \end{bmatrix} \quad B = \begin{bmatrix} 0 & 1 & 7 & 7 \\ -1 & -2 & 0 & 4 \\ 0 & 0 & 1 & 5 \end{bmatrix} \quad C = \begin{bmatrix} 3 & 2 \\ 0 & 1 \\ -2 & -1 \end{bmatrix}$$

Exactly one of the six products  $AB$ ,  $AC$ ,  $BA$ ,  $BC$ ,  $CA$ ,  $CB$  can be computed. Determine which one, and compute it.

**Solution:**  $CA$  is the only one that can be computed, and

$$CA = \begin{bmatrix} 2 & 6 & 11 & 1 \\ 1 & 3 & 7 & 2 \\ -1 & -3 & -5 & 0 \end{bmatrix}$$

□

**M2.** Determine if the matrix  $\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix}$  is invertible.

**Solution:** The second column is a multiple of the first, so it is not invertible.

□

**M3.** Find the inverse of the matrix  $\begin{bmatrix} 3 & -1 & 0 \\ 2 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$ .

**Solution:**

$$\begin{bmatrix} 3 & -1 & 0 \\ 2 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 0 & \frac{1}{2} & -\frac{1}{3} \\ -1 & \frac{2}{3} & -\frac{2}{5} \\ 1 & -\frac{3}{2} & \frac{5}{2} \end{bmatrix}$$

□

**G2.** Compute the eigenvalues, along with their algebraic multiplicities, of the matrix  $\begin{bmatrix} 8 & -3 & -1 \\ 21 & -8 & -3 \\ -7 & 3 & 2 \end{bmatrix}$ .

**Solution:**

$$\begin{aligned}\det(A - \lambda I) &= (8 - \lambda) \det \begin{bmatrix} -8 - \lambda & -3 \\ 3 & 2 - \lambda \end{bmatrix} - (-3) \det \begin{bmatrix} 21 & -3 \\ -7 & 2 - \lambda \end{bmatrix} + (-1) \det \begin{bmatrix} 21 & -8 - \lambda \\ -7 & 3 \end{bmatrix} \\ &= (8 - \lambda) (\lambda^2 + 6\lambda - 7) + 3(-21\lambda + 21) - (-7\lambda + 7) \\ &= (\lambda - 1) ((8 - \lambda)(\lambda + 7) - 63 + 7) \\ &= (\lambda - 1)(\lambda - \lambda^2) \\ &= -\lambda(\lambda - 1)^2\end{aligned}$$

So the eigenvalues are 0 (with algebraic multiplicity 1) and 1 (with algebraic multiplicity 2).

□

**G3.** Compute the eigenspace associated to the eigenvalue 2 in the matrix  $\begin{bmatrix} -1 & 1 & 0 \\ -9 & 5 & 0 \\ 15 & -5 & 2 \end{bmatrix}$ .

**Solution:** The eigenspace is the solution space of the system  $(B - 2I)X = 0$ .

$$\text{RREF}(B - 2I) = \text{RREF} \left( \begin{bmatrix} -3 & 1 & 0 \\ -9 & 3 & 0 \\ 15 & -5 & 0 \end{bmatrix} \right) = \begin{bmatrix} 1 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

So the system simplifies to  $x - \frac{y}{3} = 0$ , or  $3x = y$ . Thus the eigenspace is

$$E_2 = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$$

□

M1:  M2:  M3:  G2:  G3:  G1: