

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

(a) Show that this vector space has an additive identity element.

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$; then $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

Now we will show the other seven properties. Let $(x_1, y_1), (x_2, y_2) \in V$, and let $c, d \in \mathbb{R}$.

1) Since real addition is associative, \oplus is associative.

2) Since real addition is commutative, \oplus is commutative.

3) The additive identity is $(1, 1)$.

4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .

5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

6) $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$

7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

Therefore V is a vector space.

□

V3. Determine if the vectors $\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}$, $\begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}$, and $\begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix}$ span \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there are zero rows, they do not span. Alternatively, by inspection $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$, so the set is linearly dependent, so it spans a subspace of dimension at most 3, therefore it does not span \mathbb{R}^4 .

□

V4. Let W be the set of all polynomials of the form $ax^3 + bx$. Determine if W is a subspace of \mathcal{P}^3 .

Solution: Yes because $s(a_1x^3 + b_1x) + t(a_2x^3 + b_2x) = (sa_1 + ta_2)x^3 + (sb_1 + tb_2)x$ also belongs to W . Alternately, yes because W is isomorphic to \mathbb{R}^2 .

□

S2. Determine if the set $\{x^3 - x, x^2 + x + 1, x^3 - x^2 + 2, 2x^2 - 1\}$ is a basis of \mathcal{P}_3

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 \\ -1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix is not the identity matrix, it is not a basis.

□

V1:

V3:

V4:

S2:

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all pairs of real numbers with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \odot (x_1, y_1) = (c^2 x_1, c^3 y_1)$$

(a) Show that this scalar multiplication \odot distributes over vector addition \oplus .

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1), (x_2, y_2) \in V$ and let $c \in \mathbb{R}$.

$$\begin{aligned} c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + x_2, y_1 + y_2) \\ &= (c^2(x_1 + x_2), c^3(y_1 + y_2)) \\ &= (c^2 x_1, c^3 y_1) \oplus (c^2 x_2, c^3 y_2) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2) \end{aligned}$$

However, V is not a vector space, as the other distributive law fails:

$$(c + d) \odot (x_1, y_1) = ((c + d)^2 x_1, (c + d)^3 y_1) \neq ((c^2 + d^2)x_1, (c^3 + d^3)y_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

□

V3. Does $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 12 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ -3 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ -8 \end{bmatrix} \right\} = \mathbb{R}^3$?

Solution: Since

$$\text{RREF} \begin{bmatrix} 2 & 3 & 1 & -4 \\ -1 & 12 & 4 & 2 \\ 4 & -9 & -3 & -8 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 1/3 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

has a zero row, the vectors fail to span \mathbb{R}^3 .

□

V4. Let W be the set of all complex numbers that are purely real (i.e. of the form $a + 0i$) or purely imaginary (i.e. of the form $0 + bi$). Determine if W is a subspace of \mathbb{C} .

Solution: No, because 1 is purely real and i is purely imaginary, but the linear combination $1 + i$ is neither.

□

S2. Determine if the set $\{x^2 + x - 1, 3x^2 - x + 1, 2x - 2\}$ is a basis of \mathcal{P}_2

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

V1:

V3:

V4:

S2:

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

- (a) Show that this vector space has an additive identity element.
(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$; then $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

Now we will show the other seven properties. Let $(x_1, y_1), (x_2, y_2) \in V$, and let $c, d \in \mathbb{R}$.

- 1) Since real addition is associative, \oplus is associative.
2) Since real addition is commutative, \oplus is commutative.
3) The additive identity is $(1, 1)$.
4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .
5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

- 6) $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$
7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

- 8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

Therefore V is a vector space.

□

V3. Determine if the vectors $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$, $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$, $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$, and $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$ span \mathbb{R}^3 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span \mathbb{R}^3 .

□

V4. Determine if $\left\{ \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^4 .

Solution: It is closed under addition and scalar multiplication, so it is a subspace. Alternatively, it is the image of the linear transformation from $\mathbb{R}^3 \rightarrow \mathbb{R}^4$ given by

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix}.$$

□

S2. Determine if the set $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^3

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

V1: ☐

V3: ☐

V4: ☐

S2: ☐

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

(a) Show that this vector space has an additive identity element.

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$; then $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

Now we will show the other seven properties. Let $(x_1, y_1), (x_2, y_2) \in V$, and let $c, d \in \mathbb{R}$.

1) Since real addition is associative, \oplus is associative.

2) Since real addition is commutative, \oplus is commutative.

3) The additive identity is $(1, 1)$.

4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .

5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

6) $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$

7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

Therefore V is a vector space.

□

V3. Determine if the vectors $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$, $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$, $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$, and $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$ span \mathbb{R}^3 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span \mathbb{R}^3 .

□

V4. Let W be the set of all \mathbb{R}^3 vectors $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ satisfying $x + y + z = 0$ (this forms a plane). Determine if W is a subspace of \mathbb{R}^3 .

Solution: Yes, because $z = -x - y$ and $a \begin{bmatrix} x_1 \\ y_1 \\ -x_1 - y_1 \end{bmatrix} + b \begin{bmatrix} x_2 \\ y_2 \\ -x_2 - y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + bx_2 \\ ay_1 + by_2 \\ -(ax_1 + bx_2) - (ay_1 + by_2) \end{bmatrix}$.
Alternately, yes because W is isomorphic to \mathbb{R}^2 .

□

S2. Determine if the set $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^3

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

V1:

V3:

V4:

S2:

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

- (a) Show that this vector space has an additive identity element.
(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$; then $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

Now we will show the other seven properties. Let $(x_1, y_1), (x_2, y_2) \in V$, and let $c, d \in \mathbb{R}$.

- 1) Since real addition is associative, \oplus is associative.
- 2) Since real addition is commutative, \oplus is commutative.
- 3) The additive identity is $(1, 1)$.
- 4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .
- 5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

- 6) $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$
7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

Therefore V is a vector space.

□

V3. Determine if the vectors $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 5 \\ -1 \\ -2 \end{bmatrix}$, $\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$, and $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$ span \mathbb{R}^3

Solution:

$$\text{RREF} \left(\begin{bmatrix} -3 & 5 & 2 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & -2 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix has only two pivot columns, the vectors do not span \mathbb{R}^3 .

□

V4. Determine if the set of all lattice points, i.e. $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$ is a subspace of \mathbb{R}^2 .

Solution: This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

S2. Determine if the set $\{x^3 - x, x^2 + x + 1, x^3 - x^2 + 2, 2x^2 - 1\}$ is a basis of \mathcal{P}_3

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 \\ -1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix is not the identity matrix, it is not a basis.

□

V1:

V3:

V4:

S2:

Name: _____

MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

V1. Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V$, $c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

(a) Show that this vector space has an additive identity element.

(b) Determine if V is a vector space or not. Justify your answer.

Solution: Let $(x_1, y_1) \in V$; then $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.

Now we will show the other seven properties. Let $(x_1, y_1), (x_2, y_2) \in V$, and let $c, d \in \mathbb{R}$.

1) Since real addition is associative, \oplus is associative.

2) Since real addition is commutative, \oplus is commutative.

3) The additive identity is $(1, 1)$.

4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .

5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

6) $1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$

7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cx_2 - 2c + 1, cy_1 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

Therefore V is a vector space.

□

V3. Does $\text{span} \left\{ \begin{bmatrix} 2 \\ -1 \\ 4 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 3 \\ 5 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 5 \\ 1 \\ -3 \end{bmatrix} \right\} = \mathbb{R}^5$?

Solution: Since there are only three vectors, they cannot span \mathbb{R}^5 .

□

V4. Determine if the set of all lattice points, i.e. $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$ is a subspace of \mathbb{R}^2 .

Solution: This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

S2. Determine if the set $\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 0 \\ 0 \\ -1 \end{bmatrix} \right\}$ is a basis of \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & 0 & -1 & 0 \\ 1 & 2 & 0 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

V1:

V3:

V4:

S2: