

| |
|-------|
| Name: |
| J#: |
| Date: |

Dr. Clontz

MASTERY QUIZ DAY 12

Math 237 – Linear Algebra

Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

| | |
|---------------------|-------|
| Standard V1. | Mark: |
|---------------------|-------|

Let V be the set of all points on the line $x + y = 2$ with the operations, for any $(x_1, y_1), (x_2, y_2) \in V, c \in \mathbb{R}$,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 2))\end{aligned}$$

Determine if V is a vector space or not.

Solution:

- 1) Since real addition is associative, \oplus is associative.
- 2) Since real addition is commutative, \oplus is commutative.
- 3) $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$, so $(1, 1)$ is an additive identity element.
- 4) $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$, so $(2 - x_1, 2 - y_1)$ is the additive inverse of (x_1, y_1) .
- 5)

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1))) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

$$6) 1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$$

7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cy_1 - 2c + 1, cx_2 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}(c + d) \odot (x_1, y_1) &= ((c + d)x_1 - (c + d - 1), (c + d)y_1 - (c + d - 1)) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= c \odot (x_1, y_1) \oplus d \odot (x_1, y_1)\end{aligned}$$

□

| | |
|---------------------|-------|
| Standard V3. | Mark: |
|---------------------|-------|

Determine if the vectors $\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}$, $\begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}$, and $\begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix}$ span \mathbb{R}^4 .

Solution:

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there are zero rows, they do not span. Alternatively, by inspection $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$, so the set is linearly dependent, so it spans a subspace of dimension at most 3, therefore it does not span \mathbb{R}^4 .

□

| | |
|---------------------|-------|
| Standard V4. | Mark: |
|---------------------|-------|

Let W be the set of all 2 by 2 matrices which are not invertible. Determine if W is a subspace of $M_{2,2}$.

Solution: W is closed under scalar multiplication, but not under addition.

□