

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

### Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard V1.</b>	Mark:
---------------------	-------

Let  $V$  be the set of all pairs of real numbers with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V$ ,  $c \in \mathbb{R}$ ,

$$(x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \odot (x_1, y_1) = (0, cy_1)$$

(a) Show that scalar multiplication **distributes vectors** over scalar addition:

$$(c + d) \odot (x, y) = c \odot (x, y) \oplus d \odot (x, y).$$

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $(x_1, y_1) \in V$ , and let  $c, d \in \mathbb{R}$ . Then

$$(c + d) \odot (x_1, y_1) = (0, (c + d)y_1) = (0, cy_1) \oplus (0, dy_1) = c \odot (x_1, y_1) \oplus d \odot (x_1, y_1).$$

However,  $V$  is not a vector space, as  $1 \odot (x_1, y_1) = (0, y_1) \neq (x_1, y_1)$ .

□

<b>Standard V3.</b>	Mark:
---------------------	-------

Determine if the vectors  $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span  $\mathbb{R}^4$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Let  $W$  be the set of all complex numbers  $a + bi$  satisfying  $a = 2b$ . Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** Yes, because  $c(2b_1 + b_1i) + d(2b_2 + b_2i) = 2(cb_1 + db_2) + (cb_1 + db_2)i$  belongs to  $W$ . Alternately, yes because  $W$  is isomorphic to  $\mathbb{R}$ .

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\{x^3 - 3x^2 + 2x + 2, -x^3 + 4x^2 - x + 1, -x^3 + 2x + 1, 3x^2 + 3x + 9\}$  is a basis of  $\mathcal{P}^3$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

### Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard V1.</b>	Mark:
---------------------	-------

Let  $V$  be the set of all polynomials with the operations, for any  $f, g \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned} f \oplus g &= f' + g' \\ c \odot f &= cf' \end{aligned}$$

(here  $f'$  denotes the derivative of  $f$ ).

- (a) Show that scalar multiplication **distributes scalars** over vector addition:  $c \odot (f \oplus g) = c \odot f \oplus c \odot g$ .
- (b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $f, g \in \mathcal{P}$ , and let  $c \in \mathbb{R}$ .

$$c \odot (f \oplus g) = c \odot (f' + g') = c(f' + g')' = cf'' + cg'' = cf' \oplus cg' = c \odot f \oplus c \odot g.$$

However, this is not a vector space, as there is no zero vector. Additionally,  $1 \odot f \neq f$  for any nonzero polynomial  $f$ .

□

<b>Standard V3.</b>	Mark:
---------------------	-------

Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Let  $W$  be the set of all polynomials of even degree. Determine if  $W$  is a subspace of the vector space of all polynomials.

**Solution:**  $W$  is closed under scalar multiplication, but not under addition. For example,  $x - x^2$  and  $x^2$  are both in  $W$ , but  $(x - x^2) + (x^2) = x \notin W$ .

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\{x^3 - 3x^2 + 2x + 2, -x^3 + 4x^2 - x + 1, -x^3 + 2x + 1, 3x^2 + 3x + 9\}$  is a basis of  $\mathcal{P}^3$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

Standard V1.	Mark:
--------------	-------

Let  $V$  be the set of all real numbers with the operations, for any  $x, y \in V$ ,  $c \in \mathbb{R}$ ,

$$x \oplus y = \sqrt{x^2 + y^2}$$

$$c \odot x = cx$$

(a) Show that the vector **addition**  $\oplus$  is **associative**:  $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $x, y, z \in \mathbb{R}$ . Then

$$\begin{aligned} (x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\ &= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\ &= \sqrt{x^2 + y^2 + z^2} \\ &= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\ &= x \oplus \sqrt{y^2 + z^2} \\ &= x \oplus (y \oplus z) \end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

Standard V3.	Mark:
--------------	-------

Determine if the vectors  $\begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 1 \\ 0 \\ -3 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 3 \\ 0 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} -1 \\ 1 \\ -1 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 1 & 3 & 1 \\ 2 & 0 & 0 & -1 \\ 1 & -3 & -2 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Since every row contains a pivot, the vectors span  $\mathbb{R}^4$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Determine if the set of all lattice points, i.e.  $\{(x, y) \mid x \text{ and } y \text{ are integers}\}$  is a subspace of  $\mathbb{R}^2$ .

**Solution:** This set is closed under addition, but not under scalar multiplication so it is not a subspace.

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

Standard V1.	Mark:
--------------	-------

Let  $V$  be the set of all points on the line  $x + y = 2$  with the operations, for any  $(x_1, y_1), (x_2, y_2) \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned}(x_1, y_1) \oplus (x_2, y_2) &= (x_1 + x_2 - 1, y_1 + y_2 - 1) \\ c \odot (x_1, y_1) &= (cx_1 - (c - 1), cy_1 - (c - 1))\end{aligned}$$

- Show that this vector space has an **additive identity** element  $\mathbf{0}$  satisfying  $(x, y) \oplus \mathbf{0} = (x, y)$ .
- Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $(x_1, y_1) \in V$ ; then  $(x_1, y_1) \oplus (1, 1) = (x_1, y_1)$ , so  $(1, 1)$  is an additive identity element.

Now we will show the other seven properties. Let  $(x_1, y_1), (x_2, y_2) \in V$ , and let  $c, d \in \mathbb{R}$ .

- Since real addition is associative,  $\oplus$  is associative.
- Since real addition is commutative,  $\oplus$  is commutative.
- The additive identity is  $(1, 1)$ .
- $(x_1, y_1) \oplus (2 - x_1, 2 - y_1) = (1, 1)$ , so  $(2 - x_1, 2 - y_1)$  is the additive inverse of  $(x_1, y_1)$ .
- 

$$\begin{aligned}c \odot (d \odot (x_1, y_1)) &= c \odot (dx_1 - (d - 1), dy_1 - (d - 1)) \\ &= (c(dx_1 - (d - 1)) - (c - 1), c(dy_1 - (d - 1)) - (c - 1)) \\ &= (cdx_1 - cd + c - (c - 1), cdy_1 - cd + c - (c - 1)) \\ &= (cdx_1 - (cd - 1), cdy_1 - (cd - 1)) \\ &= (cd) \odot (x_1, y_1)\end{aligned}$$

$$6) \ 1 \odot (x_1, y_1) = (x_1 - (1 - 1), y_1 - (1 - 1)) = (x_1, y_1)$$

7)

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (x_1 + y_1 - 1, x_2 + y_2 - 1) \\ &= (c(x_1 + y_1 - 1) - (c - 1), c(x_2 + y_2 - 1) - (c - 1)) \\ &= (cx_1 + cy_1 - 2c + 1, cx_2 + cy_2 - 2c + 1) \\ &= (cx_1 - (c - 1), cy_1 - (c - 1)) \oplus (cx_2 - (c - 1), cy_2 - (c - 1)) \\ &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)\end{aligned}$$

8)

$$\begin{aligned}
 (c+d) \odot (x_1, y_1) &= ((c+d)x_1 - (c+d-1), (c+d)y_1 - (c+d-1)) \\
 &= (cx_1 - (c-1), cy_1 - (c-1)) \oplus (dx_1 - (d-1), dy_1 - (d-1)) \\
 &= c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)
 \end{aligned}$$

Therefore  $V$  is a vector space.

□

<b>Standard V3.</b>	Mark:
---------------------	-------

Determine if the vectors  $\begin{bmatrix} 8 \\ 21 \\ -7 \end{bmatrix}$ ,  $\begin{bmatrix} -3 \\ -8 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 11 \\ -5 \end{bmatrix}$  span  $\mathbb{R}^3$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 21 & -8 & -3 & 11 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the rank is less than 3, they do not span  $\mathbb{R}^3$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Let  $W$  be the set of all complex numbers that are purely real (i.e of the form  $a + 0i$ ) or purely imaginary (i.e. of the form  $0 + bi$ ). Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** No, because 1 is purely real and  $i$  is purely imaginary, but the linear combination  $1 + i$  is neither.

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 0 \\ -1 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & 0 & -1 & 0 \\ 1 & 2 & 0 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--



Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard V1.</b>	Mark:
---------------------	-------

Let  $V$  be the set of all real numbers with the operations, for any  $x, y \in V$ ,  $c \in \mathbb{R}$ ,

$$x \oplus y = \sqrt{x^2 + y^2}$$

$$c \odot x = cx$$

- (a) Show that the vector **addition**  $\oplus$  is **associative**:  $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ .
- (b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $x, y, z \in \mathbb{R}$ . Then

$$\begin{aligned}
 (x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\
 &= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\
 &= \sqrt{x^2 + y^2 + z^2} \\
 &= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\
 &= x \oplus \sqrt{y^2 + z^2} \\
 &= x \oplus (y \oplus z)
 \end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

<b>Standard V3.</b>	Mark:
---------------------	-------

Determine if the vectors  $\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there are zero rows, they do not span. Alternatively, by inspection  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ , so the set is linearly dependent, so it spans a subspace of dimension at most 3, therefore it does not span  $\mathbb{R}^4$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Let  $W$  be the set of all polynomials of the form  $ax^3 + bx$ . Determine if  $W$  is a subspace of  $\mathcal{P}^3$ .

**Solution:** Yes because  $s(a_1x^3 + b_1x) + t(a_2x^3 + b_2x) = (sa_1 + ta_2)x^3 + (sb_1 + tb_2)x$  also belongs to  $W$ . Alternately, yes because  $W$  is isomorphic to  $\mathbb{R}^2$ .

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\left\{ \begin{bmatrix} 1 & -3 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 3 \\ 3 & 9 \end{bmatrix} \right\}$  is a basis of  $M_{2,2}$  or not.

**Solution:**

$$\text{RREF} \begin{bmatrix} 1 & -1 & -1 & 0 \\ -3 & 4 & 0 & 3 \\ 2 & -1 & 2 & 3 \\ 2 & 1 & 1 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 14

Math 237 – Linear Algebra

## Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

Standard V1.	Mark:
--------------	-------

Let  $V$  be the set of all real numbers with the operations, for any  $x, y \in V$ ,  $c \in \mathbb{R}$ ,

$$x \oplus y = \sqrt{x^2 + y^2}$$

$$c \odot x = cx$$

(a) Show that the vector **addition**  $\oplus$  is **associative**:  $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $x, y, z \in \mathbb{R}$ . Then

$$\begin{aligned} (x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\ &= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\ &= \sqrt{x^2 + y^2 + z^2} \\ &= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\ &= x \oplus \sqrt{y^2 + z^2} \\ &= x \oplus (y \oplus z) \end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

Standard V3.	Mark:
--------------	-------

Determine if the vectors  $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 5 \\ -1 \\ -2 \end{bmatrix}$ ,  $\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$ , and  $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} -3 & 5 & 2 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & -2 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix has only two pivot columns, the vectors do not span  $\mathbb{R}^3$ .

□

<b>Standard V4.</b>	Mark:
---------------------	-------

Let  $W$  be the set of all complex numbers that are purely real (i.e of the form  $a + 0i$ ) or purely imaginary (i.e. of the form  $0 + bi$ ). Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**Solution:** No, because 1 is purely real and  $i$  is purely imaginary, but the linear combination  $1 + i$  is neither.

□

<b>Standard S2.</b>	Mark:
---------------------	-------

Determine if the set  $\{x^2 + x - 1, 3x^2 - x + 1, 2x^2 - 2\}$  is a basis of  $\mathcal{P}^2$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 2 \\ 1 & -1 & 0 \\ -1 & 1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Since the resulting matrix is the identity matrix, it is a basis.

□

<b>Additional Notes/Marks</b>	
-------------------------------	--