

Name: \_\_\_\_\_

**SEMIFINAL**

Math 237 – Linear Algebra

**Version 3**

Fall 2017

**Choose up to 6 problems to work.** Work each problem on one of the attached pages; write the standard in the lower left corner. Show all work and justify all of your answers. Answers without work or sufficient reasoning will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**E1.** Write an augmented matrix corresponding to the following system of linear equations.

$$\begin{aligned}x_1 + 3x_2 - 4x_3 + x_4 &= 5 \\3x_1 + 9x_2 + x_3 - 7x_4 &= 0 \\x_1 - x_3 + x_4 &= 1\end{aligned}$$

**E2.** Put the following matrix in reduced row echelon form.

$$\left[ \begin{array}{ccc|c} -3 & 1 & 0 & 2 \\ -8 & 2 & -1 & 6 \\ 0 & 2 & 3 & -2 \end{array} \right]$$

**E3.** Solve the system of linear equations.

$$\begin{aligned}2x + y - z + w &= 5 \\3x - y - 2w &= 0 \\-x + 5z + 3w &= -1\end{aligned}$$

**E4.** Find a basis for the solution set of the system of equations

$$\begin{aligned}x + 2y + 3z + w &= 0 \\3x - y + z + w &= 0 \\2x - 3y - 2z &= 0 \\-x + 2z + 5w &= 0\end{aligned}$$

**V1.** Let  $V$  be the set of all polynomials with the operations, for any  $f, g \in V$ ,  $c \in \mathbb{R}$ ,

$$\begin{aligned}f \oplus g &= f' + g' \\c \odot f &= cf'\end{aligned}$$

(here  $f'$  denotes the derivative of  $f$ ).

(a) Show that scalar multiplication **distributes scalars** over vector addition:  $c \odot (f \oplus g) = c \odot f \oplus c \odot g$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**V2.** Determine if  $\begin{bmatrix} 0 \\ -1 \\ 2 \\ 6 \end{bmatrix}$  can be written as a linear combination of the vectors  $\begin{bmatrix} 3 \\ -1 \\ -1 \\ 0 \end{bmatrix}$  and  $\begin{bmatrix} -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$ .

**V3.** Determine if the vectors  $\begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}$ , and  $\begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix}$  span  $\mathbb{R}^4$ .

**V4.** Let  $W$  be the set of all complex numbers  $a + bi$  satisfying  $a = 2b$ . Determine if  $W$  is a subspace of  $\mathbb{C}$ .

**S1.** Determine if the vectors  $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix}$  are linearly dependent or linearly independent

**S2.** Determine if the set  $\{x^3 - x, x^2 + x + 1, x^3 - x^2 + 2, 2x^2 - 1\}$  is a basis of  $\mathcal{P}^3$ .

**S3.** Let  $W = \text{span} \left( \left\{ \begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$ . Find a basis of  $W$ .

**S4.** Let  $W = \text{span} \left( \left\{ \begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$ . Compute the dimension of  $W$ .

**A1.** Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  be the linear transformation given by

$$T \left( \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \right) = \begin{bmatrix} x_1 + 3x_3 \\ 3x_2 - 5x_3 \end{bmatrix}.$$

Write the matrix for  $T$  with respect to the standard bases of  $\mathbb{R}^4$  and  $\mathbb{R}^2$ .

**A2.** Determine if  $D : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$  given by  $D \left( \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = a - 3c$  is a linear transformation or not.

**A3.** Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by the standard matrix  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the standard matrix  $\begin{bmatrix} 2 & 3 & -1 & -2 \\ 0 & 1 & 4 & 1 \\ 2 & 1 & -7 & -4 \end{bmatrix}$

**A4.** Let  $T : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} a & b \\ x & y \end{bmatrix} \right) = \begin{bmatrix} a + x \\ 0 \\ b + y \end{bmatrix}$ . Compute a basis for the kernel and a basis for the image of  $T$ .

**M1.** Let

$$A = \begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix} \qquad B = \begin{bmatrix} 3 & 1 & 0 \end{bmatrix} \qquad C = \begin{bmatrix} 3 & -1 & 4 \\ 1 & 0 & 2 \end{bmatrix}$$

Exactly one of the six products  $AB$ ,  $AC$ ,  $BA$ ,  $BC$ ,  $CA$ ,  $CB$  can be computed. Determine which one, and compute it.

**M2.** Determine if the matrix  $\begin{bmatrix} 3 & -1 & 0 \\ 2 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$  is invertible.

**M3.** Find the inverse of the matrix  $\begin{bmatrix} 2 & -1 & -3 \\ -14 & 9 & 24 \\ 3 & -2 & -5 \end{bmatrix}$ .

**G1.** Compute the determinant of the matrix  $\begin{bmatrix} 3 & -1 & 0 & 4 \\ 2 & 1 & 1 & -1 \\ 0 & 1 & 1 & 3 \\ 1 & -2 & 0 & 0 \end{bmatrix}$ .

**G2.** Compute the eigenvalues, along with their algebraic multiplicities, of the matrix  $\begin{bmatrix} 2 & -3 & 2 \\ 8 & -9 & 5 \\ 8 & -7 & 3 \end{bmatrix}$ .

**G3.** Compute the eigenspace of the eigenvalue  $-1$  in the matrix  $\begin{bmatrix} 4 & -2 & -1 \\ 15 & -7 & -3 \\ -5 & 2 & 0 \end{bmatrix}$ .

**G4.** Compute the geometric multiplicity of the eigenvalue  $-1$  in the matrix  $\begin{bmatrix} 4 & -2 & -1 \\ 15 & -7 & -3 \\ -5 & 2 & 0 \end{bmatrix}$ .

**Standard:** \_\_\_\_\_



**Standard:** \_\_\_\_\_



**Standard:** \_\_\_\_\_



**Standard:** \_\_\_\_\_



**Standard:** \_\_\_\_\_





**Standard:** \_\_\_\_\_

