

Name: _____

MASTERY QUIZ DAY 28

Math 237 – Linear Algebra

Version 6

Fall 2017

Show all work and justify all of your answers. Answers without work or sufficient reasoning will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

M1. Let

$$A = \begin{bmatrix} 3 \\ 5 \\ -1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & -1 & 3 & -3 \\ 2 & 1 & -1 & 2 \end{bmatrix} \quad C = \begin{bmatrix} 2 & -1 \\ 0 & 4 \\ 3 & 1 \end{bmatrix}$$

Exactly one of the six products AB , AC , BA , BC , CA , CB can be computed. Determine which one, and compute it.

Solution: CB is the only one that can be computed, and

$$CB = \begin{bmatrix} 0 & -3 & 7 & -8 \\ 8 & 4 & -4 & 8 \\ 5 & -2 & 8 & -7 \end{bmatrix}$$

□

M2. Determine if the matrix $\begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix}$ is invertible.

Solution: The second column is a multiple of the first, so it is not invertible.

□

M3. Find the inverse of the matrix $\begin{bmatrix} 4 & -1 & -8 \\ 2 & 1 & 3 \\ 1 & 1 & 4 \end{bmatrix}$.

Solution: $\left[\begin{array}{ccc|ccc} 4 & -1 & -8 & 1 & 0 & 0 \\ 2 & 1 & 3 & 0 & 1 & 0 \\ 1 & 1 & 4 & 0 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -4 & 5 \\ 0 & 1 & 0 & -5 & 24 & -28 \\ 0 & 0 & 1 & 1 & -5 & 6 \end{array} \right]$. Thus the inverse is $\begin{bmatrix} 1 & -4 & 5 \\ -5 & 24 & -28 \\ 1 & -5 & 6 \end{bmatrix}$.

□

G2. Let $A = \begin{bmatrix} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \end{bmatrix}$. List the eigenvalues of A along with their algebraic multiplicities.

Solution:

$$\begin{aligned}\det(A - \lambda I) &= \det \begin{bmatrix} -3 - \lambda & 1 & 0 \\ -8 & 2 - \lambda & -1 \\ 0 & 2 & 3 - \lambda \end{bmatrix} \\ &= (-3 - \lambda) \det \begin{bmatrix} 2 - \lambda & -1 \\ 2 & 3 - \lambda \end{bmatrix} - (1) \det \begin{bmatrix} -8 & -1 \\ 0 & 3 - \lambda \end{bmatrix} \\ &= (-3 - \lambda) ((2 - \lambda)(3 - \lambda) + 2) - (-8(3 - \lambda)) \\ &= (-3 - \lambda)(8 - 5\lambda + \lambda^2) + 24 - 8\lambda \\ &= -\lambda^3 + 2\lambda^2 + 7\lambda - 24 + 24 - 8\lambda \\ &= -\lambda^3 + 2\lambda^2 - \lambda \\ &= -\lambda(\lambda^2 - 2\lambda + 1) \\ &= -\lambda(\lambda - 1)^2\end{aligned}$$

So A has eigenvalues 0 (with multiplicity 1) and 1 (with algebraic multiplicity 2).

□

G3. Compute the eigenspace of the eigenvalue -1 in the matrix $\begin{bmatrix} 4 & -2 & -1 \\ 15 & -7 & -3 \\ -5 & 2 & 0 \end{bmatrix}$.

Solution:

$$\text{RREF}(A + I) = \begin{bmatrix} 1 & -\frac{2}{5} & -\frac{1}{5} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

So the eigenspace is spanned by $\begin{bmatrix} 2 \\ 5 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 0 \\ 5 \end{bmatrix}$.

□

M1: M2: M3: G2: G3: G1: