

Name: \_\_\_\_\_

**MIDTERM EXAM**

Math 237 – Linear Algebra

**Version 3**

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

**E1.** Write an augmented matrix corresponding to the following system of linear equations.

$$\begin{aligned}x_1 + 3x_2 - 4x_3 + x_4 &= 5 \\ 3x_1 + 9x_2 + x_3 - 7x_4 &= 0 \\ x_1 - x_3 + x_4 &= 1\end{aligned}$$

**Solution:**

$$\left[ \begin{array}{cccc|c} 1 & 3 & -4 & 1 & 5 \\ 3 & 9 & 1 & -7 & 0 \\ 1 & 0 & -1 & 1 & 1 \end{array} \right]$$

□

**E2.** Find RREF  $A$ , where

$$A = \left[ \begin{array}{ccc|c} 2 & -1 & 5 & 4 \\ -1 & 0 & -2 & -1 \\ 1 & 3 & -1 & -5 \end{array} \right]$$

**Solution:**

$$\text{RREF } A = \left[ \begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

□

**E3.** Find the solution set for the following system of linear equations.

$$\begin{aligned}2x_1 + 3x_2 - 5x_3 + 14x_4 &= 8 \\ x_1 + x_2 - x_3 + 5x_4 &= 3\end{aligned}$$

**Solution:** Let  $A = \left[ \begin{array}{cccc|c} 2 & 3 & -5 & 14 & 8 \\ 1 & 1 & -1 & 5 & 3 \end{array} \right]$ , so  $\text{RREF } A = \left[ \begin{array}{cccc|c} 1 & 0 & 2 & 1 & 1 \\ 0 & 1 & -3 & 4 & 2 \end{array} \right]$ . It follows that the solution set is given by  $\begin{bmatrix} 1 - 2a - b \\ 2 + 3a - 4b \\ a \\ b \end{bmatrix}$  for all real numbers  $a, b$ .

□

**E4.** Find a basis for the solution set of the system of equations

$$\begin{aligned}x + 3y + 3z + 7w &= 0 \\ x + 3y - z - w &= 0 \\ 2x + 6y + 3z + 8w &= 0 \\ x + 3y - 2z - 3w &= 0\end{aligned}$$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the solution set is

$$\left\{ \begin{bmatrix} -3a - b \\ a \\ -2b \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

So a basis for the solution set is

$$\left\{ \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \\ -1 \end{bmatrix} \right\}$$

□

**V1.** Let  $V$  be the set of all real numbers with the operations, for any  $x, y \in V$ ,  $c \in \mathbb{R}$ ,

$$x \oplus y = \sqrt{x^2 + y^2}$$

$$c \odot x = cx$$

(a) Show that the vector **addition**  $\oplus$  is **associative**:  $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ .

(b) Determine if  $V$  is a vector space or not. Justify your answer.

**Solution:** Let  $x, y, z \in \mathbb{R}$ . Then

$$\begin{aligned} (x \oplus y) \oplus z &= \sqrt{x^2 + y^2} \oplus z \\ &= \sqrt{(\sqrt{x^2 + y^2})^2 + z^2} \\ &= \sqrt{x^2 + y^2 + z^2} \\ &= \sqrt{x^2 + (\sqrt{y^2 + z^2})^2} \\ &= x \oplus \sqrt{y^2 + z^2} \\ &= x \oplus (y \oplus z) \end{aligned}$$

However, this is not a vector space, as there is no zero vector.

□

**V2.** Determine if  $\begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix}$  belongs to the span of the set  $\left\{ \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ -6 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\}$ .

**Solution:** Since

$$\text{RREF} \left( \left[ \begin{array}{ccc|c} 1 & 2 & 0 & 3 \\ 2 & 4 & 0 & -2 \\ -3 & -6 & 0 & 4 \end{array} \right] \right) = \left[ \begin{array}{ccc|c} 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

contains the contradiction  $0 = 1$ ,  $\begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix}$  is not a linear combination of the three vectors.

□

**V3.** Determine if the vectors  $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} 5 \\ -1 \\ -2 \end{bmatrix}$ ,  $\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$ , and  $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$  span  $\mathbb{R}^3$

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} -3 & 5 & 2 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & -2 & -1 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since the resulting matrix has only two pivot columns, the vectors do not span  $\mathbb{R}^3$ .

□

**V4.** Determine if  $\left\{ \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix} \mid x, y, z \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^4$ .

**Solution:** It is closed under addition and scalar multiplication, so it is a subspace. Alternatively, it is the image of the linear transformation from  $\mathbb{R}^3 \rightarrow \mathbb{R}^4$  given by

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} x \\ y \\ 0 \\ z \end{bmatrix}.$$

□

**S1.** Determine if the set of vectors  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} \right\}$  is linearly dependent or linearly independent

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 3 \\ 1 & -1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

Since there is a nonpivot column, the set is linearly dependent.

□

**S2.** Determine if the set  $\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 0 \\ -1 \end{bmatrix} \right\}$  is a basis of  $\mathbb{R}^4$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & -1 & 0 & 2 \\ 1 & 0 & -1 & 0 \\ 1 & 2 & 0 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this is not the identity matrix, the set is not a basis.

□

**S3.** Let  $W$  be the subspace of  $\mathcal{P}_3$  given by  
 $W = \text{span}(\{x^3 + x^2 + 2x + 1, 3x^3 + 3x^2 + 6x + 3, 3x^3 - x^2 + 3x - 2, 7x^3 - x^2 + 8x - 3\})$ . Find a basis for  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then a basis is  $\{x^3 + x^2 + 2x + 1, 3x^3 - x^2 + 3x - 2\}$ .

□

**S4.** Let  $W = \text{span} \left( \left( \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix} \right) \right)$ . Find the dimension of  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This has two pivot columns, so  $W$  has dimension 2.

□

**E1:**

**E2:**

**E3:**

**E4:**

**V1:**

**V2:**

**V3:**

**V4:**

**S1:**

**S2:**

**S3:**

**S4:**