

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 15

Math 237 – Linear Algebra

## Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

Standard V2.	Mark:
--------------	-------

Determine if  $\begin{bmatrix} 1 \\ 4 \\ 3 \end{bmatrix}$  is a linear combination of the vectors  $\begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$ ,  $\begin{bmatrix} 1 \\ -1 \\ 4 \end{bmatrix}$ , and  $\begin{bmatrix} 5 \\ 1 \\ -6 \end{bmatrix}$ .

**Solution:**

$$\text{RREF} \left( \left[ \begin{array}{ccc|c} 3 & 1 & 5 & 1 \\ 0 & -1 & 1 & 4 \\ -1 & 4 & -6 & 3 \end{array} \right] \right) = \left[ \begin{array}{ccc|c} 1 & 0 & 2 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

So  $\begin{bmatrix} 1 \\ 4 \\ 3 \end{bmatrix}$  is not a linear combination of the three vectors.

□

Standard S1.	Mark:
--------------	-------

Determine if the set of polynomials  $\{x^3 - 8x, x^3 + 2x^2 + 2, -x^2 + 3\}$  is linearly dependent or linearly independent

**Solution:**

$$\text{RREF} \left( \left[ \begin{array}{ccc} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \\ 0 & 0 & 0 \end{array} \right] \right) = \left[ \begin{array}{ccc} 1 & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

This has a non pivot column, therefore the set is linearly dependent.

□

Standard S3.	Mark:
--------------	-------

Let  $W = \text{span} \left( \left\{ \begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$ . Find a basis of  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ -2 & 3 & 1 & 0 \\ 0 & 6 & 1 & 1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then  $\left\{ \begin{bmatrix} 2 \\ 0 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 3 \\ 6 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}$  is a basis of  $W$ .

□

<b>Standard S4.</b>	Mark:
---------------------	-------

Let  $W = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ -1 \\ 3 \\ -3 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ -7 \end{bmatrix} \right\} \right)$ . Compute the dimension of  $W$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 2 & 3 & 1 \\ -1 & 0 & -1 & 1 \\ 3 & 1 & 4 & 1 \\ -3 & 1 & -2 & -7 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This has 3 pivot columns so  $\dim(W) = 3$ .

□

<b>Additional Notes/Marks</b>	
-------------------------------	--