

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 21

Math 237 – Linear Algebra

### Version 1

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if the following linear maps are injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  given by  $S \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} 3x + 2y \\ x - y \\ x + 4y \end{bmatrix}$

(b)  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $T \left( \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} x + y + z \\ 2y + 3z \\ x - y - 2z \end{bmatrix}$

**Solution:**

(a)

$$\text{RREF} \left( \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 3 \\ 1 & -1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

Since there is a nonpivot column,  $T$  is not injective. Since there is a zero row,  $T$  is not surjective.

(b)

$$\text{RREF} \left( \begin{bmatrix} 3 & 2 \\ 1 & -1 \\ 1 & 4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Since all columns are pivot columns,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^4$  be the linear transformation given by

$$T \left( \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} x + 3y + 3z + 7w \\ x + 3y - z - w \\ 2x + 6y + 3z + 8w \\ x + 3y - 2z - 3w \end{bmatrix}$$

Compute the kernel and image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the kernel is

$$\ker(T) = \left\{ \begin{bmatrix} -3a - b \\ a \\ -2b \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\} = \text{span} \left( \left\{ \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \\ -1 \end{bmatrix} \right\} \right)$$

and the image is

$$\text{Im}(T) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix} \right\} \right) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix} \right\} \right)$$

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 21

Math 237 – Linear Algebra

### Version 2

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if the following linear maps are injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  given by  $S \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} 3x + 2y \\ x - y \\ x + 4y \end{bmatrix}$

(b)  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $T \left( \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} x + y + z \\ 2y + 3z \\ x - y - 2z \end{bmatrix}$

**Solution:**

(a)

$$\text{RREF} \left( \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 3 \\ 1 & -1 & -2 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

Since there is a nonpivot column,  $T$  is not injective. Since there is a zero row,  $T$  is not surjective.

(b)

$$\text{RREF} \left( \begin{bmatrix} 3 & 2 \\ 1 & -1 \\ 1 & 4 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Since all columns are pivot columns,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} 8x - 3y - z + 4w \\ y + 3z - 4w \\ -7x + 3y + 2z - 5w \end{bmatrix}$ . Compute the kernel and image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 0 & 1 & 3 & -4 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Im}(T) = \text{span} \left( \left\{ \begin{bmatrix} 8 \\ 0 \\ -7 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 3 \end{bmatrix} \right\} \right)$$

$$\text{ker}(T) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 3 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$$

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 21

Math 237 – Linear Algebra

### Version 3

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 7 & -1 & 5 \end{bmatrix}$

### Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a no zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 7 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$

Since there is not a zero row,  $T$  is surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the linear transformation given by

$$T \left( \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right) = \begin{bmatrix} -3x + y \\ -8x + 2y - z \\ 2y + 3z \end{bmatrix}$$

Compute the kernel and image of  $T$ .

**Solution:** Let  $A = \begin{bmatrix} -3 & 1 & 0 \\ -8 & 2 & -1 \\ 0 & 2 & 3 \end{bmatrix}$ , and compute  $\text{RREF}(A) = \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$ . Then the image is the span of the (pivot) columns, so

$$\text{Im } T = \text{span} \left( \left\{ \begin{bmatrix} -3 \\ -8 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \right\} \right)$$

The kernel is the solution set of  $AX = 0$ , so

$$\ker T = \left\{ \begin{bmatrix} c \\ 3c \\ -2c \end{bmatrix} \mid c \in \mathbb{R} \right\} = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} \right\} \right)$$

□

Additional Notes/Marks	
------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 21

Math 237 – Linear Algebra

## Version 4

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix}$

## Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{4}{5} & -\frac{2}{5} \\ 0 & 1 & \frac{1}{5} & \frac{3}{5} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row,  $T$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^4$  be the linear transformation given by

$$T \left( \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} x + 3y + 3z + 7w \\ x + 3y - z - w \\ 2x + 6y + 3z + 8w \\ x + 3y - 2z - 3w \end{bmatrix}$$

Compute the kernel and image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the kernel is

$$\ker(T) = \left\{ \begin{bmatrix} -3a - b \\ a \\ -2b \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\} = \text{span} \left( \left\{ \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \\ -1 \end{bmatrix} \right\} \right)$$

and the image is

$$\text{Im}(T) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix} \right\} \right) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix} \right\} \right)$$

□

<b>Additional Notes/Marks</b>	
-------------------------------	--



Name:
J#:
Date:

Dr. Clontz

## MASTERY QUIZ DAY 21

Math 237 – Linear Algebra

### Version 5

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix}$

### Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{4}{5} & -\frac{2}{5} \\ 0 & 1 & \frac{1}{5} & \frac{3}{5} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row,  $T$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be the linear map given by  $T \left( \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} 8x - 3y - z + 4w \\ y + 3z - 4w \\ -7x + 3y + 2z - 5w \end{bmatrix}$ . Compute the kernel and image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 8 & -3 & -1 & 4 \\ 0 & 1 & 3 & -4 \\ -7 & 3 & 2 & -5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Im}(T) = \text{span} \left( \left\{ \begin{bmatrix} 8 \\ 0 \\ -7 \end{bmatrix}, \begin{bmatrix} -3 \\ 1 \\ 3 \end{bmatrix} \right\} \right)$$

$$\text{ker}(T) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 3 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ 0 \\ 1 \end{bmatrix} \right\} \right)$$

□

<b>Additional Notes/Marks</b>	
-------------------------------	--

Name:
J#:
Date:

Dr. Clontz

# MASTERY QUIZ DAY 21

Math 237 – Linear Algebra

## Version 6

Fall 2017

Show all work. Answers without work will not receive credit. You may use a calculator, but you must show all relevant work to receive credit for a standard.

<b>Standard A3.</b>	Mark:
---------------------	-------

Determine if each of the following linear transformations is injective (one-to-one) and/or surjective (onto).

(a)  $S : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  given by the matrix  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix}$ .

(b)  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  given by the matrix  $\begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix}$

## Solution:

(a)  $\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 0 & 1 \\ 3 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Since each column is a pivot column,  $S$  is injective. Since there is a zero row,  $S$  is not surjective.

(b) Since  $\dim \mathbb{R}^4 > \dim \mathbb{R}^3$ ,  $T$  is not injective.

$$\text{RREF} \left( \begin{bmatrix} 2 & 3 & -1 & 1 \\ -1 & 1 & 1 & 1 \\ 4 & 11 & -1 & 5 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & -\frac{4}{5} & -\frac{2}{5} \\ 0 & 1 & \frac{1}{5} & \frac{3}{5} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since there is a zero row,  $T$  is not surjective.

□

<b>Standard A4.</b>	Mark:
---------------------	-------

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^4$  be the linear transformation given by

$$T \left( \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} x + 3y + 3z + 7w \\ x + 3y - z - w \\ 2x + 6y + 3z + 8w \\ x + 3y - 2z - 3w \end{bmatrix}$$

Compute the kernel and image of  $T$ .

**Solution:**

$$\text{RREF} \left( \begin{bmatrix} 1 & 3 & 3 & 7 \\ 1 & 3 & -1 & -1 \\ 2 & 6 & 3 & 8 \\ 1 & 3 & -2 & -3 \end{bmatrix} \right) = \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Then the kernel is

$$\ker(T) = \left\{ \begin{bmatrix} -3a - b \\ a \\ -2b \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\} = \text{span} \left( \left\{ \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \\ -1 \end{bmatrix} \right\} \right)$$

and the image is

$$\text{Im}(T) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 6 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ -1 \\ 8 \\ -3 \end{bmatrix} \right\} \right) = \text{span} \left( \left\{ \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ 3 \\ -2 \end{bmatrix} \right\} \right)$$

□

Additional Notes/Marks	
------------------------	--