

Gamma-Poisson Model

General Principles

To model the relationship between a count outcome variable and one or more independent variables with overdispersion , we can use the *Negative Binomial model*.

Considerations

Caution

- We have the same considerations as for the [Poisson model](#).
- Overdispersion is handled because the Gamma-Poisson model assumes that each Poisson count observation has its own rate. This is an additional parameter specified in the model (in the code, it is `log_days`).

Example

Below is an example code snippet demonstrating a Bayesian Gamma-Poisson model using the Bayesian Inference (BI) package.

Python

```
from BI import bi, jnp
```

```
# Setup device -----
```

```
m = bi(platform='cpu') # Import
```

```
# Import Data & Data Manipulation -----
```

```
# Import
```

```

from importlib.resources import files
data_path = m.load.sim_gamma_poisson(only_path=True)
m.data(data_path, sep=',')

# Define model -----
def model(log_days, monastery, y):
    a = m.dist.normal(0, 1, name = 'a', shape=(1,))
    b = m.dist.normal(0, 1, name = 'b', shape=(1,))
    phi = m.dist.exponential(1, name = 'phi', shape=(1,))
    mu = jnp.exp(log_days + a + b * monastery)
    Lambda = m.dist.gamma(rate = mu*phi, concentration = phi, name = 'Lambda')
    m.dist.poisson(rate = Lambda, obs=y)
# Run MCMC -----
m.fit(model, progress_bar=False) # Optimize model parameters through MCMC sampling

# Summary -----
m.summary() # Get posterior distributions

```

jax.local_device_count 16

	mean	sd	hdi_5.5%	hdi_94.5%	mcse_mean	mcse_sd	ess_bulk	ess_tail	r_hat
Lambda[0]	1.46	0.34	0.91	1.98	0.00	0.01	7092.60	2636.76	1.00
Lambda[1]	1.47	0.35	0.90	1.99	0.00	0.01	6563.80	2953.08	1.00
Lambda[2]	1.62	0.37	1.06	2.20	0.00	0.01	6043.02	2447.33	1.00
Lambda[3]	1.55	0.35	0.96	2.07	0.00	0.01	7858.22	2777.30	1.00
Lambda[4]	1.46	0.35	0.91	2.00	0.00	0.01	7697.28	2700.66	1.00
...	...	...	...	...	...	...	...	...	...
Lambda[3398]	3.14	0.75	1.90	4.26	0.01	0.02	7399.87	2573.72	1.00
Lambda[3399]	2.97	0.71	1.91	4.14	0.01	0.01	6194.16	2676.34	1.00
a[0]	-0.41	0.02	-0.43	-0.38	0.00	0.00	537.82	896.11	1.01
b[0]	-2.75	0.03	-2.80	-2.70	0.00	0.00	859.02	1588.78	1.00
phi[0]	17.01	2.47	12.93	20.60	0.26	0.16	96.47	147.81	1.05

R

```

library(BayesianInference)

# Setup platform-----

```

```

m=importBI(platform='cpu')

# Import data -----
m$data(m$load$sim_gamma_poisson(only_path=T), sep = '')
m$data_to_model(list('log_days', 'monastery', 'y' )) # Send to model (convert to jax array)

# Define model -----
model <- function(log_days, monastery, y){
  # Parameter prior distributions
  alpha = bi.dist.normal(0, 1, name='alpha', shape=c(1))
  beta = bi.dist.normal(0, 1, name='beta', shape=c(1))
  phi = bi.dist.exponential(1, name='phi', shape=c(1))
  mu = jnp$exp(log_days + alpha + beta * monastery)
  Lambda = bi.dist.gamma(rate = mu*phi, concentration = phi, name = 'Lambda')
  # Likelihood
  bi.dist.poisson(rate=Lambda, obs=y)
}

# Run MCMC -----
m$fit(model) # Optimize model parameters through MCMC sampling

# Summary -----
m$summary() # Get posterior distributions

```

Julia

```

using BayesianInference

# Setup device-----
m = importBI(platform="cpu")

# Import Data & Data Manipulation -----
# Import
data_path = m.load.sim_gamma_poisson(only_path = true)
m.data(data_path, sep=',')

# Define model -----
@BI function model(log_days, monastery, y)
  a = m.dist.normal(0, 1, name = "a", shape=(1,))
  b = m.dist.normal(0, 1, name = "b", shape=(1,))
  phi = m.dist.exponential(1, name = "phi", shape=(1,))
  mu = jnp.exp(log_days + a + b * monastery)

```

```

    Lambda = m.dist.gamma(rate = mu*phi, concentration = phi, name = "Lambda")
    m.dist.poisson(rate = Lambda, obs=y)
end

# Run mcmc -----
m.fit(model) # Optimize model parameters through MCMC sampling

# Summary -----
m.summary() # Get posterior distributions

```

Mathematical Details

Bayesian model

In the Bayesian formulation, we define each parameter with priors . We can express the Bayesian regression model accounting for prior distributions as follows:

$$Y_i \sim \text{Poisson}(\lambda_i)$$

$$\lambda_i \sim \text{Gamma}(\mu_i \phi, \phi)$$

$$\log(\mu_i) = \text{rates}_i + \alpha + \beta X_i$$

$$\alpha \sim \text{Normal}(0, 1)$$

$$\beta \sim \text{Normal}(0, 1)$$

$$\phi \sim \text{Exponential}(1)$$

Where:

- Y_i is the dependent variable for observation i .
- λ_i is the rate parameter of the Poisson distribution for observation i , assuming that each Poisson count observation has its own $rate_i$.
- μ_i is the mean rate parameter.

-
- ϕ controls the level of overdispersion in the rates.
- α is the intercept term.
- β is the regression coefficient.
- X_i is the value of the predictor variable for observation i .

Notes

Note

- We can apply multiple variables similarly as in [chapter 2](#).
- We can apply interaction terms similarly as in [chapter 3](#).
- We can apply categorical variables similarly as in [chapter 4](#).

Reference(s)