

Gamma-Poisson Model

General Principles

To model the relationship between a count outcome variable and one or more independent variables with overdispersion , we can use the *Negative Binomial model*.

Considerations

Caution

- We have the same considerations as for the [Poisson model](#).
- Overdispersion is handled because the Gamma-Poisson model assumes that each Poisson count observation has its own rate. This is an additional parameter specified in the model (in the code, it is `log_days`).

Example

Below is an example code snippet demonstrating a Bayesian Gamma-Poisson model using the Bayesian Inference (BI) package.

Python

```
from BI import bi, jnp
```

```
# Setup device -----  
m = bi(platform='cpu') # Import
```

```
# Import Data & Data Manipulation -----  
# Import
```

```

from importlib.resources import files
data_path = m.load.sim_gamma_poisson(only_path=True)
m.data(data_path, sep=',')

# Define model -----
def model(log_days, monastery, y):
    a = m.dist.normal(0, 1, name = 'a', shape=(1,))
    b = m.dist.normal(0, 1, name = 'b', shape=(1,))
    phi = m.dist.exponential(1, name = 'phi', shape=(1,))
    mu = jnp.exp(log_days + a + b * monastery)
    Lambda = m.dist.gamma(rate = mu*phi, concentration = phi, name = 'Lambda')
    m.dist.poisson(rate = Lambda, obs=y)
# Run MCMC -----
m.fit(model, progress_bar=False) # Optimize model parameters through MCMC sampling

# Summary -----
m.summary() # Get posterior distributions

```

jax.local_device_count 16

arviz - WARNING - Shape validation failed: input_shape: (1, 500), minimum_shape: (chains=2, c

	mean	sd	hdi_5.5%	hdi_94.5%	mcse_mean	mcse_sd	ess_bulk	ess_tail	r_hat
Lambda[0]	1.56	0.41	0.92	2.17	0.01	0.03	1260.23	333.77	NaN
Lambda[1]	1.51	0.35	0.98	2.04	0.01	0.01	761.43	472.24	NaN
Lambda[2]	1.58	0.39	0.92	2.11	0.01	0.02	711.05	393.66	NaN
Lambda[3]	1.42	0.34	0.92	1.93	0.01	0.02	958.15	283.08	NaN
Lambda[4]	1.47	0.34	0.95	2.01	0.01	0.01	816.98	438.47	NaN
...	...	...	...	...	...	...	...	...	...
Lambda[3398]	3.50	0.80	2.24	4.66	0.02	0.04	1154.11	415.89	NaN
Lambda[3399]	3.67	0.83	2.48	5.04	0.03	0.06	643.82	165.42	NaN
a[0]	-0.42	0.01	-0.45	-0.40	0.00	0.00	69.68	122.11	NaN
b[0]	-2.75	0.03	-2.80	-2.70	0.00	0.00	81.87	261.49	NaN
phi[0]	17.30	2.10	13.86	20.23	0.44	0.29	22.93	58.25	NaN

R

```

library(BayesianInference)

# Setup platform-----
m=importBI(platform='cpu')

# Import data -----
m$data(m$load$sim_gamma_poisson(only_path=T), sep = '')
m$data_to_model(list('log_days', 'monastery', 'y' )) # Send to model (convert to jax array)

# Define model -----
model <- function(log_days, monastery, y){
  # Parameter prior distributions
  alpha = bi.dist.normal(0, 1, name='alpha', shape=c(1))
  beta = bi.dist.normal(0, 1, name='beta', shape=c(1))
  phi = bi.dist.exponential(1, name='phi', shape=c(1))
  mu = jnp$exp(log_days + alpha + beta * monastery)
  Lambda = bi.dist.gamma(rate = mu*phi, concentration = phi, name = 'Lambda')
  # Likelihood
  bi.dist.poisson(rate=Lambda, obs=y)
}

# Run MCMC -----
m$fit(model) # Optimize model parameters through MCMC sampling

# Summary -----
m$summary() # Get posterior distributions

```

Julia

```

using BayesianInference

# Setup device-----
m = importBI(platform="cpu")

# Import Data & Data Manipulation -----
# Import
data_path = m.load.sim_gamma_poisson(only_path = true)
m.data(data_path, sep=',')

# Define model -----
@BI function model(log_days, monastery, y)

```

```

a = m.dist.normal(0, 1, name = "a", shape=(1,))
b = m.dist.normal(0, 1, name = "b", shape=(1,))
phi = m.dist.exponential(1, name = "phi", shape=(1,))
mu = jnp.exp(log_days + a + b * monastery)
Lambda = m.dist.gamma(rate = mu*phi, concentration = phi, name = "Lambda")
m.dist.poisson(rate = Lambda, obs=y)
end

# Run mcmc -----
m.fit(model) # Optimize model parameters through MCMC sampling

# Summary -----
m.summary() # Get posterior distributions

```

Mathematical Details

Bayesian model

In the Bayesian formulation, we define each parameter with priors . We can express the Bayesian regression model accounting for prior distributions as follows:

$$Y_i \sim \text{Poisson}(\lambda_i)$$

$$\lambda_i \sim \text{Gamma}(\mu_i \phi, \phi)$$

$$\log(\mu_i) = \text{rates}_i + \alpha + \beta X_i$$

$$\alpha \sim \text{Normal}(0, 1)$$

$$\beta \sim \text{Normal}(0, 1)$$

$$\phi \sim \text{Exponential}(1)$$

Where:

- Y_i is the dependent variable for observation i .

- λ_i is the rate parameter of the Poisson distribution for observation i , assuming that each Poisson count observation has its own *rate* _{i} .
- μ_i is the mean rate parameter.
-
- ϕ controls the level of overdispersion in the rates.
- α is the intercept term.
- β is the regression coefficient.
- X_i is the value of the predictor variable for observation i .

Notes

Note

- We can apply multiple variables similarly as in [chapter 2](#).
- We can apply interaction terms similarly as in [chapter 3](#).
- We can apply categorical variables similarly as in [chapter 4](#).

Reference(s)