

Tutorial sheet - 2

SPRING 2020

MATHEMATICS-II (MA10002)(Linear Algebra)

1. Determine which of the following form a basis of the respective vector spaces:

- (a) $\{4t^2 - 2t + 3, 6t^2 - t + 4, 8t^2 - 8t + 7\}$ of $\mathbb{P}_2(\mathbb{R})$,
- (b) Let V be a real vector space with $\{\alpha, \beta, \gamma\}$ as a basis. Check whether $\{\alpha + \beta + \gamma, \beta + \gamma, \gamma\}$ is also a basis of V
- (c)

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \right\}$$

for V , where V is the vector space of all 2×2 real matrices.

2. Determine the basis and dimension of the following subspaces

- (a) The subspace V , of all 2×2 real symmetric matrices.
- (b) $U = \{(x, y, z, w) \in \mathbb{R}^4 : x+2y-z=0, 2x+y+w=0\}$ of \mathbb{R}^4 .
- (c) Let $U = \{p \in \mathbb{P}_4(\mathbb{R}) : \int_{-1}^1 p(t)dt = 0\}$.

3. If $U = L(\{(1, 2, 1), (2, 1, 3)\})$, $W = L(\{(1, 0, 0), (0, 0, 1)\})$, show that U and W are subspaces of \mathbb{R}^3 . Find the dimension of $W_1, W_2, W_1 + W_2, W_1 \cap W_2$.

4. Check the following mappings are linear transformation or not:

- (a) $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$, defined by $T(x, y, z) = (x^2, |y| + z)$, $\forall (x, y, z) \in \mathbb{R}^3$.
- (b) $T: \mathbb{P}_3(\mathbb{R}) \rightarrow \mathbb{P}_4(\mathbb{R})$, defined by $T(p(x)) = (1-x)p'(0) - xp(x)$.

5. Give an example of a function $\phi: \mathbb{C} \rightarrow \mathbb{C}$, such that $\phi(w+z) = \phi(w) + \phi(z)$, $\forall w, z \in \mathbb{C}$. But ϕ is not linear over \mathbb{C} .

6. Find the null space and range space of the following linear transformations. Also find their respective dimensions and verify the rank-nullity theorem.

- (a) $T: \mathbb{P}_2(\mathbb{R}) \rightarrow \mathbb{P}_3(\mathbb{R})$ defined by $T(f(x)) = 2f'(x) + \int_0^x 3f(t)dt$.
- (b) $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$, defined by $T(x, y, z) = (\frac{x-y-z}{2}, \frac{z}{2})$
- (c) $T: M_{2 \times 2}(F) \rightarrow M_{2 \times 2}(F)$ defined by $T(A) = \frac{A - A^T}{2}$, $\forall A \in M_{2 \times 2}(F)$.

7. (a) Determine the linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ such that $T(1, 1) = (1, 0, 2)$, $T(2, 3) = (1, -1, 4)$.

- (b) Determine the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ which maps the basis vectors $\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ of \mathbb{R}^3 to the vectors $\{(1, 1), (2, 3), (3, 2)\}$ respectively.
 - i) Find $T(1, 1, 0)$, $T(6, 0, -1)$,
 - ii) Find $\text{Ker}(T)$ & $\text{Im}(T)$.
 - iii) Prove that T is not one-to-one but onto.

8. Find the matrix of the linear transformations w.r.t the given ordered bases:

(a) $D : \mathbb{P}_4(\mathbb{R}) \rightarrow \mathbb{P}_4(\mathbb{R})$ defined by $D(p(x)) = 3 \frac{d^3}{dx^3}(p(x))$, w.r.t. the ordered basis $\{1, x, x^2, x^3, x^4\}$ for both $\mathbb{P}_4(\mathbb{R})$.

(b) $T : P_3(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ by

$$T(f(x)) = \begin{bmatrix} 2f''(0) & f(3) \\ 0 & f'(2) \end{bmatrix}$$

w.r.t. the ordered basis $\{1, x, x^2, x^3\}$ and $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$.

9. Prove that there does not exist a linear map $T : \mathbb{R}^5 \rightarrow \mathbb{R}^5$ such that $R(T) = N(T)$.

10. Solve the following system of equations by Gauss-elimination method:

$$\begin{array}{ll} (a) & \begin{array}{l} 9x + 3y + 4z = 7 \\ 4x + 3y + 4z = 8 \\ x + y + z = 3 \end{array} \\ (b) & \begin{array}{l} x + 2y + 3z + 2w = -1 \\ -x - 2y - 2z + w = 2 \\ 2x + 4y + 8z + 12w = 4 \end{array} \end{array}$$

11. Find the rank of the matrix A using definition where

$$(i) A = \begin{bmatrix} 1 & 0 & 3 \\ 4 & -1 & 5 \\ 2 & 0 & 6 \end{bmatrix} \quad (ii) \begin{bmatrix} 2 & 3 & -1 & 1 \\ 3 & 0 & 4 & 2 \\ 6 & 9 & -3 & 3 \end{bmatrix}$$

12. Determine the rank of the following matrices by reducing to row echelon form.

$$(a) \begin{bmatrix} 1 & 2 & 1 & 0 \\ 2 & 4 & 8 & 6 \\ 3 & 6 & 6 & 3 \end{bmatrix} \quad (b) \begin{bmatrix} 0 & 0 & 2 & 2 & 0 \\ 1 & 3 & 2 & 4 & 1 \\ 2 & 6 & 2 & 6 & 2 \\ 3 & 9 & 1 & 10 & 6 \end{bmatrix}$$

13. Find all x such that the rank of the matrix $\begin{bmatrix} 1 & x & x \\ x & 1 & x \\ x & x & 1 \end{bmatrix}$ is less than 3.

14. Determine whether the following matrices are invertible or not, if it is, then compute the inverse :

$$(a) \begin{bmatrix} 1 & 1 & 2 \\ 2 & 4 & 4 \\ 3 & 3 & 7 \end{bmatrix} \quad (b) \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -1 \\ 1 & 5 & 4 \end{bmatrix}$$

15. Find the value of k for which the system of equations has non-trivial solution.

$$\begin{array}{l} x + 2y + z = 0 \\ 2x + y + 3z = 0 \\ x + ky + 3z = 0 \end{array}$$

16. Solve the system of equations in integers

$$\begin{array}{l} x + 2y + z = 1 \\ 3x + y + 2z = 3 \\ x + 7y + 2z = 1 \end{array}$$

17. Solve if possible

$$\begin{aligned}x + 2y + z - 3w &= 1 \\2x + 4y + 3z + w &= 3 \\3x + 6y + 4z - 2w &= 5\end{aligned}$$

18. Determine the condition for which the system

$$\begin{aligned}x + y + z &= b \\2x + y + 3z &= b + 1 \\5x + 2y + az &= b^2\end{aligned}$$

admits of (i) only one solution, (ii) no solution, (iii) infinitely many solutions.