
IRIS
(day)
IRIS
(cell type)
A
Early chondro-
cytes (1) are
similar to Meso-
dermal progen-
itors (Tbx6+) (2)
but arise later
1
2
B
Epithelial cells show
branching as the
placodal area gives
rise to pre-epidermal
keratinocytes (left
branch) and gut and
olfactory epithelial
cells (right branch)
C
Different types of white
blood cells, such as
Kupffer cells (right)
and border-associated
macrophages (2nd from
right) are distinct and
arise at different times
D
Though most CNS
neurons show
branching, thalamic
neuronal precursors
have a distinct origin
Day
8
9
11
13
15
17
19
UMAP
(cell type)
UMAP
(day)
8 10 14 19
Day
CNS_neurons
Definitive_erythroid
Hepatocytes
Endothelium
Interm._neur._prog.
Primitive_erythroid
Mesoderm
Neural_crest_PNS_neurons
Neuroectoderm_and_glia
Intestine
Epithelial_cells
Muscle_cells
Mast_cells
White_blood_cells
Lung_and_airway
T_cells
Adipocytes
Cardiomyocytes
B_cells
Neural_crest_PNS_glia
Ependymal_cells
Eye_and_other
Oligodendrocytes
Olfactory_sensory_neurons
Megakaryocytes
Testis_and_adrenal