

forthright48

Learning Never Ends

Home CPPS 101 About Me

Saturday, July 11, 2015

Number of Divisors of an Integer

Problem

Given an integer N , find its number of divisors.

For example, 12 has the following divisors 1, 2, 3, 4, 6 and 12. So its number of divisors is 6.

Number Of Divisors Function: $NOD(N)$ or $\sigma_0(N)$

Number of Divisors of N is also called $\sigma_0(N)$. That's just a fancy way of asking for the number of divisors of N . You can read more on it from [Wiki](#). I will refer to Number of Divisors of N as either $\sigma_0(N)$ or $NOD(N)$. They both mean same.

Brute Force Method $O(N)$

Anyways, what is the easiest way to find Number of Divisors (NOD)? We can try to divide N with all numbers from $1 - N$ and keep count of how many of those numbers divide N . This will definitely work, but obviously we need to do better.

Optimized to $O(\sqrt{N})$

In "[Primality Test - Naive Methods](#)", we established that if we can write $N = A \times B$, then one of $\{A, B\}$ must be $\leq \sqrt{N}$.

So using that same idea we can find NOD by simply looping till \sqrt{N} and check if that particular number divides N . If it doesn't then it is not a divisor and if it does then we found a $\{A, B\}$ pair of divisors. If $A \neq B$, then we add 2 to result, otherwise we add 1.

Examples

Suppose $N = 24$. We need to loop till $\lfloor \sqrt{24} \rfloor = 4$. We get the following pairs of divisors, $\{1, 24\}, \{2, 12\}, \{3, 8\}, \{4, 6\}$ So our answer is 8.

Let's try again with $N = 36$. We need to loop till $\lfloor \sqrt{36} \rfloor = 6$. We get the following pairs of divisors, $\{1, 36\}, \{2, 18\}, \{3, 12\}, \{4, 9\}, \{6, 6\}$ So our answer is 9. Notice that the last pair does not satisfy the condition $A \neq B$. So we add one for that pair.

This brings out an interesting observation. NOD of N is always even except for when N is a perfect square. Because whenever N is perfect square, \sqrt{N} would be its divisor and it will form a pair with itself.

Code Using Loop till \sqrt{N}

```

1  int NOD ( int n ) {
2      int res = 0;
3      int sqn = sqrt ( n );
4
5      for ( int i = 1; i < sqn; i++ ) {
6          if ( n % i == 0 ) res += 2; //Found a divisor pair {A,B}
7      }
8
9      //Need to check if sqn divides n
10     if ( n % sqn == 0 ) {
11         if ( sqn * sqn == n ) res++; //If n is perfect square
12         else res += 2; //Not perfect square
13     }
14
15     return res;

```

Follow by Email

Email address...

Labels

[Analysis](#) [Arithmetic Function](#)
[Binary](#) [Bitwise](#) [Combinat](#)
[Contest](#) [CPPS](#) [D&C](#) [Divi](#)
[Factorization](#) [GCD](#) [Graph](#)
[Logarithm](#) [Math](#) [Modul](#)
[Number Theor](#)
[Primality](#) [Test](#) [Prime P](#)
[Squaring](#) [Sequence](#) [Sieve](#) [Tree](#) [UVa](#)

Blog Archive

► 2018 (2)
 ► 2017 (2)
 ▼ 2015 (35)
 ► Sep (7)
 ► Aug (13)
 ▼ Jul (15)
 [Simple Hyperbolic Equation](#)
 [Linear Diophantine](#)
 [Extended Euclidean](#)
 [Introduction to Modular Arithmetic](#)
 [Divisor Summatory](#)
 [Sum of Divisors of](#)
 [Prime Number The](#)
 [Upper Bound for N Divisors](#)
 [Highly Composite I](#)
 [Number of Divisors](#)
 [Prime Factorization](#)
 [Sieve of Eratosthe](#)
 [Generating Prim](#)
 [Primality Test - Nai](#)
 [Lowest Common I](#)
 [Number](#)
 [Euclidean Algorith](#)
 [Common Divisor](#)

Follow me on Twitter

We loop from 1 to $\sqrt{N} - 1$ at line 5. Then at line 10, we handle \sqrt{N} separately.

Using Prime Factorization

It is possible to find NOD from prime factorization of N .

Suppose $N = 12$. Its prime factorization is $2^2 \times 3$. Is it possible to divide 12 with 5? No. Cause the prime factorization of 12 does not contain 5. We can divide N with primes that appear in factorization of N .

Next, can we divide 12 with 2^3 ? No. The power of 2 in prime factorization of 12 is 2^2 . So we cannot divide 12 with 2^3 , since 2^2 is not divisible by 2^3 .

So, if we are trying to divide N , that has prime factorization $N = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_x^{a_x}$, then the divisors of N , D , will have prime factorization of form $D = p_1^{b_1} \times p_2^{b_2} \times \dots \times p_x^{b_x}$, where $b_i \leq a_i$.

For example, divisor of $12 = 2^2 \times 3$ will have prime factorization, $D = 2^x \times 3^y$, where $x \leq 2, y \leq 1$. When $x = 1$ and $y = 1$, $D = 2^1 \times 3^1 = 6$ which is a divisor of 12. If we use a different combination of $[x, y]$, then we get a different divisor. So how many different combination can we use here? We know that, $0 \leq x \leq 2$ and $0 \leq y \leq 1$. So for x we can use $\{0, 1, 2\}$ and for y we can use $\{0, 1\}$. So we can use $3 \times 2 = 6$ combinations of $\{x, y\}$. And that is our NOD.

So, if $N = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_x^{a_x}$, then $D = p_1^{b_1} \times p_2^{b_2} \times \dots \times p_x^{b_x}$. We get new divisors for using different combination for $\{b_1, b_2, \dots, b_x\}$. How many different combinations can we make? $(a_1 + 1) \times (a_2 + 1) \times \dots \times (a_x + 1)$. Therefore, $\sigma_0(N) = (a_1 + 1) \times (a_2 + 1) \times \dots \times (a_x + 1)$.

So basically, in order to find NOD, all we need to do is factorize N , then take each of its prime factors power, increase them by 1 and finally multiply all of them. Easy right?

Examples

$N = 12 = 2^2 \times 3$. $NOD(N) = (2 + 1) \times (1 + 1) = 6$.

$N = 15 = 3 \times 5$. $NOD(N) = (1 + 1) \times (1 + 1) = 4$.

$N = 252 = 2^2 \times 3^2 \times 7$. $NOD(N) = (2 + 1) \times (2 + 1) \times (1 + 1) = 18$.

Try out this yourself using other small values of N .

Code Using Prime Factorization

The code for NOD is not very different from the one we used for $factorize()$ in [Prime Factorization of an Integer](#). We just need to modify it slightly to suit our needs.

```

1  int NOD ( int n ) {
2      int sqrtn = sqrt ( n );
3      int res = 1;
4      for ( int i = 0; i < prime.size() && prime[i] <= sqrtn; i++ ) {
5          if ( n % prime[i] == 0 ) {
6              int p = 0; /*Counter for power of prime*/
7              while ( n % prime[i] == 0 ) {
8                  n /= prime[i];
9                  p++;
10             }
11             sqrtn = sqrt ( n );
12             p++; /*Increase it by one at end*/
13             res *= p; /*Multiply with answer*/
14         }
15     }
16     if ( n != 1 ) {
17         res *= 2; /*Remaining prime has power p^1. So multiply with 2*/
18     }
19     return res;
20 }
```

I have highlighted the lines that are different from $factorize()$.

Reference

1. Divisor Function - https://en.wikipedia.org/wiki/Divisor_function

Posted by [Mohammad Samiul Islam](#)



Labels: [Divisors](#), [Factorization](#), [Number Theory](#)

1 comment:

Tweets by @forthright48



Mohammad Samiul Islam
@forthright48

Is it a pokemon or bigdata?
[pixelastic.github.io/pokemon-bigdata](https://github.com/pixelastic/pokemon-bigdata)
... #pokemon #bigdata



Pokemon
Is it a Pokemon or bigdata?
pixelastic.github.io/pokemon-bigdata



Mohammad Samiul Islam
@forthright48

I don't know why I have been default bash shell all these
[ohmyz.sh](https://github.com/ohmyzsh/ohmyzsh) Glad I found @c



oh my zsh
Oh-My-Zsh
[ohmyz.sh](https://github.com/ohmyzsh/ohmyzsh)




Mohammad Samiul Islam
@forthright48

[Embed](#)

Followers

Followers (16)



[Follow](#)

Total Pageviews

7 3 1 1 7

**Misbah Ahmad** October 28, 2016 at 2:30 PM

Such a great Article.
Thanks for it.

[Reply](#)

Enter your comment...



Comment as:

BIDDUT SARF ▼

[Sign out](#)[Publish](#)[Preview](#)☐ Notify me**Leave comments for Queries, Bugs and Hugs.**[Newer Post](#)[Home](#)[Older Post](#)Subscribe to: [Post Comments \(Atom\)](#)**Popular Posts**[SPOJ LCMSUM - LCM Sum](#)

Problem Link - SPOJ LCMSUM Given n , calculate the sum $LCM(1, n) + LCM(2, n) + \dots + LCM(n, n)$, where $LCM(i, n)$ denotes the ...

[Euclidean Algorithm - Greatest Common Divisor](#)

Problem Given two number A and B, find the greatest number that divides both A and B. What we are trying to find here is the Greatest Comm...

[Extended Euclidean Algorithm](#)

Extended Euclidean Algorithm is an extension of Euclidean Algorithm which finds two things for integer a and b : It finds the value of...

[Chinese Remainder Theorem Part 1 - Coprime Moduli](#)

Second part of the series can be found on: Chinese Remainder Theorem Part 2 - Non Coprime Moduli Wow. It has been two years since I pub...

[Prufer Code: Linear Representation of a Labeled Tree](#)

I guess this is going to be my first post (apart from the contest analysis') which is not about Number Theory! It's not about graph ...

[Segmented Sieve of Eratosthenes](#)

Problem Given two integers A and B , find number of primes inside the range of A and B inclusive. Here, $1 \leq A \leq B \leq 10^6$...

[Sieve of Eratosthenes - Generating Primes](#)

Problem Given an integer N, generate all primes less than or equal to N. Sieve of Eratosthenes - Explanation Sieve of Eratosthenes ...

[Number of Digits of Factorial](#)

Problem Given an integer N , find number of digits in $N!$. For example, for $N = 3$, number of digits in $3! = 3! = 3 \times 2 \times 1 = 6$...

[Chinese Remainder Theorem Part 2 - Non Coprime Moduli](#)

As promised on the last post, today we are going to discuss the "Strong Form" of Chinese Remainder Theorem, i.e, what do we do whe...

[Euler Totient or Phi Function](#)

I have been meaning to write a post on Euler Phi for a while now, but I have been struggling with its proof. I heard it required Chinese Rem...