

Model transformations

BIOL2022 - L04c

Dr Januar Harianto The University of Sydney



Learning objectives

You should:

- ☐ Understand why model transformations are necessary.
- ☐ Differentiate between transforming the data and formulating a new model.
- ☐ Apply common transformations (log, square root) to the response variable.
- ☐ Interpret the results of a log-transformed model.

Why do we perform model transformations?


- When data does not meet LINE assumptions, we can attempt transformations to improve the model fit before considering more complex models.
- Transformations linearise (or at least, attempt to) the relationship between the response and predictor variables.
- *Not cheating!* **We are improving the model fit, not changing the data arbitrarily.**

The idea behind transformations

Given a simple model between two variables, y and x :

$$y \sim x$$

Where the relationship is not linear, we may end up with a model that does not fit the data well:



Two ways to transform

We can either formulate a new model that better fits the data, or transform the data to better fit the model. **Both methods are (practically) equivalent.**




Figure 1

It is not easy to formulate a new model

It turns out that you need a lot of domain knowledge to formulate a new model.

To fit the model to this particular dataset, we need to formulate:

$$y = a \times 2^x + b$$

Transforming the data is easier. Basically, we can transform the response variable and approximate the model:

$$\log_2(y) \sim x$$

It is not perfect and may even introduce issues, but it can be a good starting point. It is also easier to do when dealing with complex multi-factorial models.

How do we transform data?

Irregardless of the complexity of the model, apply the transformation to the response variable:

$$y \sim x_1 + x_2 + \dots + x_n \rightarrow f(y) \sim x_1 + x_2 + \dots + x_n$$

Depending on the relationship between the response and predictor variables, we can apply different transformations:

- **Logarithmic transformation:** $f(y) = \log(y)$: right skewed data
- **Square root transformation:** $f(y) = \sqrt{y}$: count data with many small values
- **Reciprocal transformation:** $f(y) = \frac{1}{y}$: when other transforms do not work

In most cases, a logarithmic transformation is a good starting point. It is also easier to interpret the results.

Interpretation of a log-transformed model

Given a model:

$$\log(y) = \beta_0 + \beta_1 x$$

where we are transforming the response variable y using the **natural** logarithm, $\log(y)$, then a one-unit increase in x is associated with a change of $(e^{\beta_1} - 1) \times 100\%$ in y .

We can also use estimated marginal means to interpret the results in R, where back-transforms can be automatically calculated to the model.

Example




New York City skyline enveloped in heavy smog, May 1973. Photo by Chester Higgins/NARA (CC BY-NC 2.0)


Air quality in New York City, 1973

Is air quality (ozone concentration) in New York City influenced by solar radiation? The model is:

$$\text{ozone} \sim \text{solar radiation}$$



Assumptions



Did the model assumptions hold?

- **Linearity:** the relationship between ozone and solar radiation is not linear, evident fan-shape in the residual vs fitted plot.
- **Normality:** the residuals in the qq-plot are “u-shaped”, indicating a positively skewed distribution.
- **Equal variance:** the residuals are not homoscedastic: increasing variance with increasing fitted values seen in the scale-location plot, although it is not severe (not more than 2 standard deviations).

Transforming the data

Given the non-linear relationship between ozone and solar radiation, we can apply a logarithmic transformation to the response variable:

$$\log(\text{ozone}) \sim \text{solar radiation}$$





Figure 2: Residual diagnostics of the log-transformed model

Comparison

Untransformed model

Transformed model



Is the model a better fit?

Yes! Fanning in the residual vs fitted plot is reduced, and the “u-shaped” distribution in the qq-plot is no longer evident. Scale-location plot shows a more consistent variance across the fitted values.

```
Call:
lm(formula = log(Ozone) ~ Solar.R, data = airquality)

Residuals:
    Min       1Q   Median       3Q      Max
-2.64991 -0.56329  0.02199  0.55373  1.47755

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  2.6152491   0.1666990   15.688  < 2e-16 ***
Solar.R       0.0043326   0.0008097    5.351 4.88e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.7741 on 109 degrees of freedom
(42 observations deleted due to missingness)
Multiple R-squared:  0.208, Adjusted R-squared:  0.2008
F-statistic: 28.63 on 1 and 109 DF,  p-value: 4.885e-07
```

How do we interpret the results?

So, is transformation necessary?

Let us compare the summaries of both models. This is sometimes called a [sensitivity analysis](#).

Untransformed model

Log-transformed model

```
1 summary(fit)
```

```
Call:
lm(formula = Ozone ~ Solar.R, data = airquality)

Residuals:
    Min       1Q   Median       3Q      Max
-48.292 -21.361  -8.864   16.373  119.136

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  18.59873     6.74790   2.756 0.006856 **
Solar.R       0.12717     0.03278   3.880 0.000179 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 31.33 on 109 degrees of freedom
(42 observations deleted due to missingness)
Multiple R-squared:  0.1213,    Adjusted R-squared:  0.1133
F-statistic: 15.05 on 1 and 109 DF,  p-value: 0.0001793
```

Interpretation trade-offs

- The **untransformed model** is easier to interpret: for every 1 W/m^2 increase in solar radiation, ozone concentration is predicted to increase by $\sim 0.13 \text{ ppb}$.
- However, this model does not fit the data well and violates several assumptions, so we cannot be confident in this prediction.
- The **log-transformed model** fits the data better, but is more difficult to interpret.
- A 1 W/m^2 increase in solar radiation is associated with a 0.43% increase in ozone concentration.
- We can also back-transform the results to the original scale, but this gives us the *median* change in ozone concentration, not the *mean*.

Verdict: If the goal is to understand the relationship between variables and **make predictions**, the transformed model is better. If the goal is simple interpretation and the model violations are not severe, the untransformed model may be acceptable: **most biologists prioritise interpretability**.

A sensitivity analysis is quick and easy to perform, so it is worth doing as soon as you are unsure of your model's assumptions (happens often).

Questions to consider

- When should you consider transforming your data versus fitting a more complex model (e.g., a generalised linear model)?
- How do you choose the appropriate transformation for your data?
- What are the challenges in interpreting the coefficients of a log-transformed model, and how can back-transformation help?
- Can transformations fix all violations of model assumptions? When might they not be enough?

Thanks!

This presentation is based on the SOLES Quarto reveal.js template and is licensed under a Creative Commons Attribution 4.0 International License.