

Modelling multiple differences: a linear model with two or more categorical X variables

BIOL2022 - L04b

Dr Januar Harianto The University of Sydney



Learning objectives

You should be able to:

- ☐ Frame ANOVA as a specific application of the General Linear Model (GLM) for categorical predictors.
- ☐ Formulate a GLM to test for differences between two or more group means.
- ☐ Explain how an ANOVA summary partitions the variance of a GLM with categorical predictors.
- ☐ Interpret main effects and interaction terms for categorical predictors within a GLM framework.
- ☐ Use post-hoc tests to probe significant main effects and interactions in a GLM.

Analysis of variance (ANOVA)

Using categorical predictors in a GLM

It is better to solve the right problem approximately than to solve the *wrong* problem exactly.

— John Tukey (1915-2000)

The ANOVA model


ANOVA is a type of linear regression that uses **categorical** predictors.

How it works (GLM perspective)

1. Fit a linear model using dummy variables for the categories.
2. Split the total variation in the response variable into:
 - Variation *between* groups (explained by the model).
 - Variation *within* groups (unexplained, or residual).
3. Compare these to test if the differences between groups are greater than the variation within groups.


Variation between groups

Can we explain the variation in body mass by species (i.e. are the means different)?




Variation within groups

Can we explain the variation in body mass by an overall mean (i.e. **are they all the same**)?



Another way to look at it

Are the distributions of body mass different enough to be considered separate?



Modelling the relationship

For a GLM:

$$\text{body mass} = \beta_0 + \beta_1 \cdot \text{species} + \epsilon$$


Once we use an ANOVA summary to process the model, we get:

$$\text{body mass} = \text{mean} + \text{species effect (i.e. difference)} + \epsilon$$

That is:

- β_0 is the **overall mean** of body mass.
- β_1 is the **difference** in body mass between species.
- We can add the mean to each species effect to get the **estimated body mass** for each species.

Example



Are species and sex significant predictors of body mass in penguins?

First, the general linear model

body mass $\sim$ species + sex

The interaction term

Because we are interested in both predictor variables, we include an **interaction term**:


body mass $\sim$ species $\times$ sex

Four steps (as always), with a post-hoc

1. Visualise the relationship.
2. Fit the regression model.
3. Check and revise the model if needed (assumptions, outliers, issues).
4. Interpret the model ✨ and post-hoc ✨.

1 of 4: Visualise the relationship

body mass $\sim$ species $\times$ sex




Side note: Beyond bar and box plots

2 of 4: Fit the model

This is performed using software so we can assume this is done correctly (for now). The model is still:

$$\text{body mass} \sim \text{species} \times \text{sex}$$

3 of 4: Check assumptions



Diagnostic plots for the linear model with species and sex as predictors of body mass.

4: Interpret

In practice, if we are interested in comparing means, the **ANOVA summary** table is the most useful.

```
Analysis of Variance Table
```

```
Response: body_mass_g
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
species	2	145190219	72595110	758.358	< 2.2e-16 ***
sex	1	37090262	37090262	387.460	< 2.2e-16 ***
species:sex	2	1676557	838278	8.757	0.0001973 ***
Residuals	327	31302628	95727		

```
---
```



```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

There are **significant interactions**, so we no longer interpret the main effects. **Interpret the interaction term instead.**

Interpreting interactions

We want to check two things:

- Whether the relationship between body mass and species is dependent on **sex**, and/or
- Whether the relationship between body mass and sex is dependent on **species**.



Interpreting interactions

- Species affects the relationship between body mass and sex. Males are generally heavier than females, but the difference is less pronounced in Chinstrap penguins as seen by the flatter slope compared to the other two species.
- Sex also affects the relationship between body mass and species. Adelie males are generally heavier than Chinstrap males, but in females it is the opposite: Adelie females are generally lighter than Chinstrap females.

What if we look at the non-ANOVA summary?

```
Call:
lm(formula = body_mass_g ~ species * sex, data = penguins)

Residuals:
    Min       1Q   Median       3Q      Max
-827.21 -213.97   11.03   206.51  861.03

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)    3368.84     36.21   93.030 < 2e-16 ***
speciesChinstrap    158.37     64.24    2.465  0.01420 *
speciesGentoo    1310.91     54.42   24.088 < 2e-16 ***
sexmale          674.66     51.21   13.174 < 2e-16 ***
speciesChinstrap:sexmale -262.89     90.85   -2.894  0.00406 **
speciesGentoo:sexmale   130.44     76.44    1.706  0.08886 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Post-hoc

How and why do we perform post-hoc tests?

- If there is a significant effect of a categorical predictor with more than two levels, we need to perform post-hoc tests to determine which specific groups differ from each other.
- Post-hoc tests control for the increased risk of Type I error that arises from making multiple comparisons, ensuring that the overall significance level is maintained.

Questions to consider

- When is it more appropriate to summarise a GLM with an ANOVA table versus regression coefficients?
- How do the standard GLM assumptions (LINE) apply when working with categorical predictors?
- How does the GLM formulation and interpretation change when including one vs. two categorical predictors, especially regarding main effects and interactions?
- What does a significant interaction term in a GLM tell us about the relationship between categorical predictors?
- Why are post-hoc tests necessary for interpreting GLMs with significant categorical predictors with more than two levels?

Thanks!

This presentation is based on the [SOLES Quarto reveal.js template](#) and is licensed under a [Creative Commons Attribution 4.0 International License](#).