MCMC Part 2

Julia Wrobel

Overview

Today, we cover:

- Gibb's sampling

Resources:

- Peng Advanced Stat Computing Chapter 7

Single component Metropolis-Hastings

The standard Metropolis algorithm updates the entire parameter vector at once with a single step.

- Hence, a p-dimensional parameter vector must have a p-dimensional proposal distribution q.
- However, it is sometimes simpler to update individual components of the parameter vector one at a time, in an algorithm known as single component Metropolis-Hastings (SCMH).

Let $x^t = (x_1^t, x_2^t, \dots, x_p^t)$ be a p-dimensional vector of parameter values at iteration t.

Single component M-H

The SCMH algorithm updates each parameter in x one at a time. All that is needed to update the ith component of x^t is a proposal distribution $q(x_i^*|x_i^t, x_{-i}^t)$. SCMH updates the ith component via the following steps:

(1) Sample $x_i^* \sim q(x_i^*|x_i^t, x_{-i}^t)$ as a proposal for component i.
(2) Let

$$\alpha(x_i^*|x^t) = \min\left\{\frac{\pi(x_i^*|x_{-i}^t)q(x_i^t|x_i^*, x_{-i}^t)}{\pi(x_i^t|x_{-i}^t)q(x_i^*|x_i^t, x_{-i}^t)}, 1\right\}$$

(3) Accept x_i^* with probability $\alpha(x_i^*|x^t)$
(4) Repeat steps 1-3 p times until each component is updated

Gibb's sampling

- Special case of single component M-H where you always accept the candidate
- For sampling from multivariate distributions (2 or more parameters)
  - Relies on ability to calculate full conditionals

Useful when:

- Sampling from $\pi(\theta_1, \theta_2, \theta_3)$ is hard
- Sampling from $\pi(\theta_1|\theta_2, \theta_3)$, $\pi(\theta_2|\theta_1, \theta_3)$, $\pi(\theta_3|\theta_1, \theta_2)$ is easy

Gibb's sampling algorithm

Steps for each iteration:

1. Sample $\theta_1^{n+1} \sim \pi(\theta_1 | \theta_2^n, \theta_3^n)$
2. Sample $\theta_2^{n+1} \sim \pi(\theta_2 | \theta_1^{n+1}, \theta_3^n)$
3. Sample $\theta_3^{n+1} \sim \pi(\theta_3 | \theta_1^{n+1}, \theta_2^{n+1})$

Do this for the desired number of samples, discard burn-in, and voila! You now have a sample from the joint distribution $\pi(\theta_1, \theta_2, \theta_3)$.

Example: Bivariate normal distribution

Suppose we want to simulate $x = (x_1, x_2)$ from a bivariate Normal distribution with mean $\mu = (\mu_1, \mu_2)$ and covariance matrix

$$\Sigma = \begin{pmatrix} \sigma_1^2 & \rho \\ \rho & \sigma_2^2 \end{pmatrix}$$

we can make use of Gibbs sampling to simulate a Markov chain that will converge to a bivariate Normal.

Bivariate normal distribution

Goal is to simulate from joint distribution $\pi(x_1, x_2|\mu, \Sigma)$.

Complete conditionals are:

- $\pi(x_1|x_2, \mu, \Sigma) = N(\mu_1 + \rho\frac{\sigma_1}{\sigma_2}(x_2 - \mu_2), \sigma_1^2(1-\rho))$
- $\pi(x_2|x_1, \mu, \Sigma) = N(\mu_2 + \rho\frac{\sigma_2}{\sigma_1}(x_1 - \mu_1), \sigma_2^2(1-\rho))$

So, we sample

1. $x_1^{t+1} \sim N(\mu_1 + \rho\frac{\sigma_1}{\sigma_2}(x_2^t - \mu_2), \sigma_1^2(1-\rho))$
2. $x_2^{t+1} \sim N(\mu_2 + \rho\frac{\sigma_2}{\sigma_1}(x_1^{t+1} - \mu_1), \sigma_2^2(1-\rho))$

Bivariate normal distribution

```
gibbs_bivariate = function(x0, n = 1000,
                           mu = c(1, 5), rho = 0.5,
                           sigma1 = 1, sigma2 = 2){
  x1 = x0[1]
  x2 = x0[2]
  mu1 = mu[1]
  mu2 = mu[2]

  x = matrix(NA, nrow = n, ncol = 2)
  for(i in 1:n){
    x[i,] = c(x1,x2)

    x1 = rnorm(1, mean = mu1 + rho * sigma1/sigma2 * (x2-mu2),
               sd = sigma1 * sqrt(1-rho))
    x2 = rnorm(1, mean = mu2 + rho * sigma2/sigma1 * (x1-mu1),
               sd = sigma2 * sqrt(1-rho))
```

Bivariate normal distribution

```r
set.seed(21433)
gibbs_x = gibbs_bivariate(c(0,0), n = 10000)

colMeans(gibbs_x[1:100,])
```

```
       x1        x2
0.7881165 4.6632079
```

```r
colMeans(gibbs_x)
```

```
      x1       x2
1.012080 5.013594
```

Bivariate normal distribution



Bivariate normal distribution



Proof: Gibbs sampling as M-H

Gibbs is a special case of single component M-H that always accepts the proposal. Define the following:

- $\pi(\theta_1, \theta_2)$: joint distribution we want to sample from
- $q(\theta_1^* | \theta_1^t, \theta_2^t)$: proposal distribution for new value of θ_1
- Acceptance ratio:

$$\alpha(\theta_1^* | \theta^t) = \min \left\{ \frac{\pi(\theta_1^*, \theta_2^t) q(\theta_1^t | \theta_1^*, \theta_2^t)}{\pi(\theta_1^t, \theta_2^t) q(\theta_1^* | \theta_1^t, \theta_2^t)}, 1 \right\}$$

- Choose full conditional as proposal, i.e.:

Proof: Gibbs sampling as M-H

Bayesian linear regression

$$y = X\beta + \epsilon; \epsilon \sim N(0, \sigma_y^2 I_n)$$

- Need to make (prior) distributional assumptions about β, σ_y^2
  - $\beta \sim N(0, \sigma_b^2 I_p)$
  - $\sigma_y^2 \sim IG(A, B)$
- Joint posterior of interest is

$$\pi(\beta, \sigma_y^2 | y, x, A, B, \sigma_b^2)$$

- Need full conditionals
  - $\pi(\beta | \sigma_y^2, y, x, A, B, \sigma_b^2)$
  - $\pi(\sigma_y^2 | \beta, y, x, A, B, \sigma_b^2)$

Full conditionals

$$\beta \mid \sigma_y^2, y \sim \mathcal{N}(\beta_{\text{post}}, \Sigma_{\text{post}})$$

$$\Sigma_{\text{post}} = \left(\frac{1}{\sigma_y^2} X^\top X + \frac{1}{\sigma_\beta^2} I_p \right)^{-1}$$

$$\beta_{\text{post}} = \left(X^\top X + \frac{\sigma_y^2}{\sigma_\beta^2} I_p \right)^{-1} (X^\top y)$$

$$\sigma_y^2 \mid \beta, y \sim IG\left(A + n/2, B + \frac{1}{2} ||y - x\beta||^2 \right)$$

Variations on Gibb's sampling

- **Blocked Gibb's sampling**: update blocks (or groups) of parameters jointly rather than one at a time
  - Helps improve mixing when variables are highly correlated
  - We are already doing this for Bayesian linear regression
- **Collapsed Gibb's sampling**: integrate/marginalize out some variables analytically before sampling
  - Reduces dimensionality and can improve convergence
- **Metropolis-within-Gibbs**: Uses Metropolis-Hastings steps for full conditionals that are not easily sampled from.
  - Commonly used in Bayesian hierarchical models

Gibbs sampler for Bayesian regression

I started implementing this myself… for lab we will go the rest of the way there and compare it to the results of `stan_glm` from the Bayesian Data Analysis lecture.

- My partial Gibb's sampler is in the file `gibbs_blr.R` and the lab file is `lab_gibbs.Rmd`.