PSYCH308A - Data Analysis 4 (DA4)

Brady C. Jackson

2024/10/04

Contents

Question 01 (Q01): Answer to Q01:					 	1 2
Question 02 (Q02): Answer to Q02:					 	2 2
Question 03 (Q03): Answer to Q03:					 	2 2
	Q04 through Q09: (RQ1): Q04 - Q06:					
Question 10 (Q10): Answer to Q10:					 	16 16
Question 11 (Q11): Answer to Q11:					 	16 16
Question 12 (Q12): Answer to Q12:					 	16 16
<pre># Load packages. Set # masking messages in library(psych) library(jmv) # library(ggplot2) library(dplyr) library(magrittr) library(tidyr) # library(stringr) #</pre>	for descriptive for pivot_longer	see	e th	е		

Question 01 (Q01):

1. Interpret a p-value of .042. (This question is **not** asking for the decision this p-value results in, rather what does this value mean?)

Answer to Q01:

A p-value of .042 implies that the finding of some statistical test (z-test, t-test, etc.) that the probability of obtaining the test statistic found or something more extreme was .042 (a.k.a. 4.2%) assuming the null hypothesis was true

Question 02 (Q02):

2. *In no more than two sentences*, what is the relationship between sample size, effect size, and power?

Answer to Q02:

Power is positively correlated with both sample size and effect size in a manner that convolutes the two (sample and effect). That is to say, the power of a finding will increase if either sample size increases or effect size increases, and if one (e.g. sample size) is necessarily small, a higher power can be achieved by increasing the other (e.g. effect size).

Question 03 (Q03):

3. A researcher records the number of words recalled by students presented with a list of words for 1 minute. In one group, students were presented with the list of words in color; in a second group, the same words were presented in black and white. An equal number of students were in each group. The researcher reports the following: Participants recalled significantly more words when the words were presented in color (M = 12.4 words) versus black and white (M = 10.9 words), t(48) = 2.01, p = .035, d = 0.18. Based on the previous statement, what is the sample size in each group?

Answer to Q03:

Given the wording of the prompt, I'm assuming this test was conducted as an independent sample t-test ("In one group ... in a second group" instead of phrasing like, "...the group of students looked at colored words and then later looked at black and white words...").

The degrees of freedom for the t-test ran is given as 48 (t(48)). As this is an independent t-test, two parameters had to be estimated, the mean of the population from which group 1 was drawn, and the mean of the population from which group 2 was drawn. Therefore, we have the relationship

$$\begin{split} df &= [Things\ we\ know\ (n_{total})] - [Estimated\ Parameters\ (n_{P,est})] \\ df &= n_{total}\ -\ n_{P,est} \\ 48 &= n_{total}\ -\ 2 \\ 48 &+ 2 = n_{total} \\ 50 &= n_{total} \end{split}$$

Therefore, there were a total of 50 students in the study, with 25 in each group given that the prompt states the same number of students were in each group.

Research Prompt for Q04 through Q09:

You are teaching your first Intro to Psychology course! After the midterm, you are disappointed with your students' overall test scores. You decide to implement two different required study techniques. There are 100 students in the class; 50 of them will be required to meet in groups to study right before the final (Group A) and the other 50 will be required to create flashcards to aid in memorization (Group B). You are interested in two primary research questions:

Research Question 1 (RQ1): Q04 - Q06:

Did student test scores improve significantly from the midterm to the final? Data: 308A.RQ1 Data.DA4.csv

Question 04 (Q04):

4. Visualize your data for this research question. Include your visualization here.

Answer to Q04:

Loading Data for Q04: The first step to visualizing the data is to load it. See code below:

```
# In order to visualize the data we must first load it. To do so we lool
# in the present working directory for CSV files, and take the one that
# has RQ1 in its filename. Append it to the current working directory
# to create the fullpath for loading
here <- getwd()
rq1_name <- list.files(here, pattern = ".*RQ1.*csv")

# Use the file.path function to create a platform appropriate fullpath / filename
# to the Research Question 1 data.
rq1_file <- file.path(here, rq1_name)

# Read the CSV data in as rq1_dat
rq1_dat <- read.csv(rq1_file, header = TRUE)

# Lower-case all column names for convenience
colnames(rq1_dat) <- tolower(colnames(rq1_dat))</pre>
```

Data Prep Q04 Ultimately we're going to answer Research Question 1 as a dependent T-test since we're evaluating change in student test scores over time (regardless) of study method. So we're going to prepare the data further with some math. See code below:

```
# Since we want to know if scores improved between the midterm and final we'll
# look at midterm scores, final scores, and the delta between the two.
# To do this, we need to add one more column to our math, the difference
# between the two. We need to maintain chirality (direction) as it will
# be informative if scores rise or decrease so we will do final - midterm
# without taking the absolute value. If scores decreased, we should see a
# positive mean of the diff. If they decreased we'll see a negative mean
# of the diff.
rq1_dat$fin_mid_diff <- rq1_dat$final - rq1_dat$midterm
```

Visualize Prepared Data Q04 - Descriptives and Bar Chart See the histograms below for visualization of the data pertinent to this


research question. Bar graphs of the midterm scores, the final scores, and the diff of final - midterm are also shown for better understanding


##

##

DESCRIPTIVES

Descriptives ## ## midterm final fin_mid_diff ## ## N 100 100 100 ## Missing 0 0 0 ## Mean 72.39620 80.45750 8.061300 ## Std. error mean 1.060126 0.4742058 1.064727 ## Median 72.16500 80.22500 8.545000 ## Standard deviation 10.60126 4.742058 10.64727 ## Variance 22.48712 113.3644 112.3867 ## Minimum 41.13000 63.80000 -26.11000 ## Maximum 95.65000 93.46000 38.56000 ## Skewness -0.1713860 0.08208768 0.1736520 ## Std. error skewness 0.2413798 0.2413798 0.2413798 ## Kurtosis 0.4235590 1.572569 0.9956757 ## Std. error kurtosis 0.4783311 0.4783311 0.4783311






```
# Now we're going to create a bar chart of the midterm scores, final scores,
# and diffs .. though the diffs data willbe oddly scaled (much smaller) than the
# raw scores
# In order to plot the scores as bar charts, we need to permute the data so all
# scores are in one vector and there's n indy axis vector of the same size that
# defines whether the score is a midterm score, final score, or difference.
# We use pivot_longer to concatenate the three dcore vectors to each other
# (interleaved), with the column names saved in the new "name" vector.
rq1_long <- rq1_dat[2:4] %>%
            pivot_longer(cols = c(midterm, final, fin_mid_diff) )
# We're going to lay all three datasets out on a single bar graph
# even though that may jackup the scaling to be bad for the diff.
bar_rq1 <- ggplot( rq1_long, aes(name, value) )</pre>
# Annotate the ggplot object to make a bar graph of test scodes.
bar_rq1 + stat_summary( fun = mean,
                        geom = "bar",
                        position = "dodge",
                        fill="slateblue1"
                      ) +
          stat_summary( fun.data = mean_cl_normal,
                        geom = "errorbar",
                        position = position_dodge(width = 0.90),
                        width = 0.2
```

```
) +
labs(x = "Test Type", y = "Score (pts))") +
ggtitle('Test Scores and Difference')
```

Test Scores and Difference



Discussion of Visualization for RQ1 It is notable that the error bar on the final scores is so much tighter than

that of the midterm scores. This is reflected in the final scores histogram which looks less "spread out" (smaller variance) than the same of the midterm scores. The larger spread in the differences of scores is also accounted for by the larger spread in the midterms as the tighter final scores minus the more spread out midterm scores will result in more spread out differences.

Question 05 (Q05):

5. Did student test scores improve significantly from the midterm to the final? Using RStudio to analyze, conduct a hypothesis test to evaluate this question. Organize your answer according to the 4 steps of hypothesis testing.

Answer to Q05:

Question 05 T-Test Code: The code below runs the t-test by feeding the final and midterm scores, not the

manually written diff above, to the t-test algorithm for paired samples. The

organization of the results of this t-test, according to the 4 steps of NHST are in the next section.

We use the ttestPS to conduct a paired samples t-test from the jamovi (jmv)

```
# package. We want to include the effect size, confidence intervals, the
# means of our inputs, the standard errors, and the descriptive statistics in
# out output.
# NOTE: we supply "final" scores as i1 instead of i2 because that seems to
        qive the same directionality in mean difference in what we found
        in our descriptive stats above.
rq1_test_results <- jmv::ttestPS( data = rq1_dat,
                                  pairs = list(list(i1='final', i2='midterm')),
                                  effectSize = TRUE,
                                  ci = TRUE,
                                  meanDiff = TRUE,
                                  desc = TRUE
# Dump the test results to output
rq1_test_results
##
##
    PAIRED SAMPLES T-TEST
##
   Paired Samples T-Test
##
##
##
                                          statistic
                                                        df
                                                                                   Mean difference
                                                                    р
##
##
      final
               midterm
                           Student's t
                                           7.571237
                                                       99.00000
                                                                    < .0000001
                                                                                          8.061300
##
##
      Note. H
               <sub>Measure 1 - Measure 2</sub>
##
##
##
    Descriptives
##
##
                 N
                        Mean
                                     Median
                                                 SD
                                                               SE
##
                 100
                                     80.22500
                                                               0.4742058
##
      final
                         80.45750
                                                   4.742058
      midterm
                        72.39620
                                     72.16500
                                                               1.0601257
##
                 100
                                                 10.601257
##
# save the test results in a dataframe object for use later
rq1_test_df <- rq1_test_results$ttest$asDF
```

SE d

Question 05 Discussion To perform the significance test we use the SCEC acronym to remember our steps:

Specify, Criteria, Estimate, Conclude:

- 1. **Specify** the Hypotheses:
 - a. Null Hypothesis: Student test scores did **not** significantly change between the midterm and the final.
 - b. Alternative Hypothesis: Student test scores dd change significantly between the midterm and the final.

```
\begin{array}{lll} H_0: \; \mu_{diff} \; = \; 0 \\ H_1: \; \mu_{diff} \; \neq \; 0 \end{array}
```

2. **Criteria:** We will check for significance using a two-tailed t-test with alpha set to .05. Our degrees of freedom will be equal to the number of samples we can take differences between, n_{diff} (a.k.a. the number of things we know), minus the number of things we're estimating, in this case the mean difference in scores from the population, of which there is 1.

```
So: df = n_{diff} - 1

df = 100 - 1

df = 99

... which gives us a t_{crit} of 1.984

\alpha = .05

df = 99

t_{crit} = 1.984
```

3. **Estimate:** Given the test conducted in the R-code above, we found the following t-test results. Please note that all values printed below are generated in text, on the fly, using the embedded r/tex-code shown after the printout:

```
\begin{array}{l} \bar{x}_{diff} = 8.0613 \\ ESE_{diff} = 1.064727 \\ t_{calc} = 7.5712366 \\ \mathrm{p} < .000001 \\ d = 0.7571237 \\ \eta^2 = 0.3666984 \end{array}
```

Verbatim Code:

```
$\bar{x}_{diff}\ =\ $ `\r rq1_test_df$'md[stud]'`
$ESE_{diff}\ =\ $ `\r rq1_test_df$'sed[stud]'`
$t_{calc}\ =\ $ `\r rq1_test_df$'stat[stud]'`
`\r if( rq1_test_df$'p[stud]' >= 0.000001 ){ paste("p =", str_sub(sprintf("%.3f", rq1_test_df$'p[stud]' > \r rq1_test_df$'es[stud]'`
$\eta^2\ =\ $ `\r rq1_test_df$'stat[stud]'^2 / (rq1_test_df$'stat[stud]'^2 + rq1_test_df$'df[stud]
```

4. Conclude: Given that $t_{calc} > t_{crit}$ we reject H_0 . The test scores did change significantly between the midterm and the final. Given that \bar{x}_{diff} is positive when the midterm scores are subtracted from the final scores, this indicates that the test scores significantly improved between the midterm and the final.

Question 06 (Q06):

6. Report your findings in APA format. (Hint: make sure to answer the research question!)

Answer to Q06: We conducted a dependent t-test to examine whether test scores improved between the midterm and final tests for a class of 100 students given required study techniques. We found that final test scores, after implementing the study techniques, were significantly improved (final, $\bar{x}=80.46,\ SD=4.74$) compared to midterm test scores before the study techniques were implemented (midterm, $\bar{x}=72.40,\ SD=10.60$), $t(99)=7.57,\ p<.001,\ \eta^2=.37$. This was a large effect; the introduction of study techniques accounted for 37% of variance in the test score differences.

Research Question 2 (RQ2): Q07 - Q09:

Does the study technique used predict scores on the final exam?

Data: 308A.RQ2 Data.DA4.csv

Question 07 (Q07):

7. Visualize your data for this research question. Include your visualization here.

Answer to Q07:

Loading Data for Q07: The first step to visualizing the data is to load it. See code below:

```
# In order to visualize the data we must first load it. To do so we look
# in the present working directory for CSV files, and take the one that
# has RQ2 in its filename. Append it to the current working directory
# to create the fullpath for loading
here <- getwd()
rq2_name <- list.files(here, pattern = ".*RQ2.*csv")

# Use the file.path function to create a platform appropriate fullpath / filename
# to the Research Question 2 data.
rq2_file <- file.path(here, rq2_name)

# Read the CSV data in as rq1_dat
rq2_dat <- read.csv(rq2_file, header = TRUE)

# Lower-case all column names for convenience
colnames(rq2_dat) <- tolower(colnames(rq2_dat))</pre>
```

Data Prep Q07 Ultimately we're going to answer Research Question 2 as an independent t-test since we're evaluating differences in test scores between the study technique employed by Group A and that by Group B. Since we're not looking at matched samples between groups (i.e. the same students are not in both groups) there is no math to do here. We just need to make sure the group category is set to the correct datatype (factor). We also rename the categories as "A" and "B" are not particularly descriptive. See code below:

```
# Recast the variables in the "group" vector to be factors instead of freeform
# character strings
rq2_dat$group <- as.factor(rq2_dat$group)

# Rename group "A" to A_group_study and group "B" to "B_flashcards"
levels(rq2_dat$group) <- sub("A", "A_group_study", levels(rq2_dat$group))
levels(rq2_dat$group) <- sub("B", "B_flashcards", levels(rq2_dat$group))</pre>
```

Visualize Prepared Data Q07 - Descriptives and Bar Chart See the histograms below for visualization of the data pertinent to this research question (RQ2). Bar graphs of the final scores for Group A and Group B


are also shown for better understanding.

```
# We're going to create a descriptives object so we can checkout the histogram # and descriptive stats of the final scores split by Group. Since we have the
```


```
# descriptives available for all final scores combined from RQ1 above, we
# won't recreate that here.
rq2_desc <- jmv::descriptives( rq2_dat,
                                vars = c('final'),
                                splitBy = c('group'),
                                hist = TRUE,
                                dens = TRUE,
                                sd = TRUE,
                                variance = TRUE,
                                se = TRUE,
                                skew = TRUE,
                                kurt = TRUE
                              )
# Render the rq2 descriptives object as output.
rq2_desc
##
```


```
##
    DESCRIPTIVES
##
##
    Descriptives
##
##
                              group
                                                final
##
##
      N
                              A_group_study
                                                         50
##
                              B_flashcards
                                                         50
##
                                                          0
      Missing
                              A_group_study
##
                              B_flashcards
                                                          0
                              A_group_study
##
      Mean
                                                   81.25500
##
                              B_flashcards
                                                   79.66000
##
      Std. error mean
                              A_group_study
                                                  0.8349668
##
                              B_flashcards
                                                  0.4307366
##
      Median
                              A_group_study
                                                   80.94000
##
                              B_flashcards
                                                   79.78500
##
      Standard deviation
                              A_group_study
                                                   5.904107
##
                              B_flashcards
                                                   3.045768
##
      Variance
                              A_group_study
                                                   34.85848
##
                              B_flashcards
                                                   9.276702
##
      Minimum
                              A_group_study
                                                   63.80000
##
                              B_flashcards
                                                   69.54000
##
      Maximum
                              A_group_study
                                                   93.46000
##
                              B_flashcards
                                                  85.75000
##
      Skewness
                              A_group_study
                                                -0.1407758
##
                              B_flashcards
                                                -0.5920366
##
      Std. error skewness
                              A_group_study
                                                  0.3366007
##
                              B_flashcards
                                                 0.3366007
##
      Kurtosis
                              A_group_study
                                                  0.5314386
##
                              B_flashcards
                                                   1.417350
##
      Std. error kurtosis
                              A_group_study
                                                  0.6619084
##
                              B_flashcards
                                                  0.6619084
```

##



```
# Now we're going to create a bar chart of the final scores sorted by group.
# No pivoting is needed here as data is already properly formatted.
# First we create the bar chart ggplot object via associating the group with
# the scores as an aesthetic object (aes)
bar_rq2 <- ggplot( rq2_dat, aes(group, final) )</pre>
# Now we annotate the bar chart appropriately. Bars will be colored orchid.
# error bars will be added.
bar_rq2 + stat_summary( fun = mean,
                        geom = "bar",
                        position = "dodge",
                        fill="orchid1"
                      ) +
          stat_summary( fun.data = mean_cl_normal,
                        geom = "errorbar",
                        position = position_dodge(width = 0.90),
                        width = 0.2
                      ) +
          labs(x = "Study Method of Group", y = "Score (pts))") +
          ggtitle('Final Test Scores by Study Method')
```





Discussion of Visualization for RQ2 The study technique employed by Group B, using flashcards for memorization,

produced less spread in test scores for that group, however, the mean final test scores were remarkably similar to those of Group A. Despite this, the study technique employed by Group A, studying in groups right before the final exam had more spread out test scores (higher variance / standard deviation). I'll be surprised if the difference in means between these two groups is significant.

Question 08 (Q08):

8. Does the study technique used predict scores on the final exam? Using RStudio to analyze, conduct a hypothesis test to evaluate this question. Organize your answer according to the 4 steps of hypothesis testing.

Answer to Q08:

Question 08 T-Test Code: The code below runs the t-test by feeding the final test scores to the t-test algorithm for independent samples. The organization of the results of this t-test, according to the 4 steps of NHST are in the next section.

```
# We use the ttestIS to conduct an independent t-test from the jamovi (jmv)
# package. We want to include the effect size, confidence intervals, the
# means of our inputs, the standard errors, and the descriptive statistics in
# out output.

rq2_test_results <- jmv::ttestIS( data = rq2_dat,
```

```
vars = 'final',
                                   group = 'group',
                                   effectSize = TRUE,
                                   ci = TRUE,
                                  meanDiff = TRUE,
                                  desc = TRUE
# Dump the test results to output
rq2_test_results
##
##
    INDEPENDENT SAMPLES T-TEST
##
##
    Independent Samples T-Test
##
##
                                                                        Mean difference
                                                                                             SE difference
                               Statistic
                                              df
                                                           p
##
##
      final
               Student's t
                               1.697670
                                             98.00000
                                                          0.0927436
                                                                                1.595000
                                                                                                0.9395231
##
##
                <sub>A_group_study</sub>
                                             <sub>B_flashcards</sub>
       Levene's test is significant (p < .05), suggesting a violation of the assumption of equal variar
##
##
##
##
    Group Descriptives
##
               Group
                                                    Median
##
                                        Mean
##
##
      final
               A_group_study
                                 50
                                        81.25500
                                                    80.94000
                                                                 5.904107
                                                                              0.8349668
               B flashcards
                                        79.66000
                                                    79.78500
##
                                 50
                                                                 3.045768
                                                                              0.4307366
##
# save the test results in a dataframe object for use later
rq2_test_df <- rq2_test_results$ttest$asDF
```

Question 08 Discussion To perform the significance test we use the SCEC acronym to remember our steps:

Specify, Criteria, Estimate, Conclude:

- 1. **Specify** the Hypotheses:
 - a. *Null Hypothesis:* There is **no** difference between the final test scores of students using the Group A study technique vs. students using the Group B study technique.
 - b. Alternative Hypothesis: There mean of Group A's final scores is different than the mean of Group B's final scores.

$$\begin{array}{lll} H_0: \ \mu_A \ = \ \mu_B \\ H_1: \ \mu_A \ \neq \ \mu_B \end{array}$$

2. Criteria: We will check for significance using a two-tailed t-test with alpha set to .05. Our degrees of freedom will be equal to the total number of students between both groups, $n_A + n_B$ (a.k.a. the number of things we know), minus the number of things we're estimating, in this case the mean of the final test scores for Group A and the mean of the final test scores for Group B, or two items.

```
So: df = n_{students} - 2
```

```
\begin{array}{ll} df &=& 100 & -2 \\ df &=& 98 \\ \text{... which gives us a } t_{crit} \text{ of } 1.984 \\ \alpha &=& .05 \\ df &=& 98 \\ t_{crit} &=& 1.984 \end{array}
```

3. **Estimate:** Given the test conducted in the R-code above, we found the following t-test results. Please note that all values printed below are generated in text, on the fly, using the embedded r/tex-code shown after the printout:

```
\begin{array}{l} \bar{x}_A \, - \, \bar{x}_B \, = \, 1.595 \\ ESED \, = \, 0.9395231 \\ t_{calc} \, = \, 1.6976698 \\ \mathrm{p} \, = \, .093 \\ d \, = \, 0.339534 \\ \eta^2 \, = \, 0.0285688 \end{array}
```

Verbatim Code:

```
$\bar{x}_{A}\ -\ \bar{x}_{B}\ =\ $ `\r rq2_test_df$'md[stud]'`
$ESED\ =\ $ `\r rq2_test_df$'sed[stud]'`
$t_{calc}\ =\ $ `\r rq2_test_df$'stat[stud]'`
`\r if( rq2_test_df$'p[stud]' >= 0.000001 ){ paste("p =", str_sub(sprintf("%.3f", rq2_test_df$'p[stud]' \s\ \r rq2_test_df$'es[stud]'`
$\eta^2\ =\ $ `\r rq2_test_df$'stat[stud]'^2 / (rq2_test_df$'stat[stud]'^2 + rq2_test_df$'df[stud]
```

4. Conclude: Given that $t_{calc} < t_{crit}$ we fail to reject H_0 . The mean final test scores did not vary significantly between Group A and Group B. We failed to find a significant difference in study technique impact on final test scores.

Question 09 (Q09):

9. ...

- a. Report your findings in APA format. (Hint: make sure to answer the research question!)
 - Answer (a): We conducted an independent t-test to determine whether studying as a group just prior to a course final (Group A), or studying using flashcards to improve memorization (Group B), was better at improving final test scores amongst 100 students, with 50 students randomly assigned to each group. The students studying as a group ($\bar{x}_A = 81.23$) did not perform significantly better or worse than those studying with flashcards ($\bar{x}_B = 79.66$), t(98) = 1.984, p = .093, $\eta^2 = .03$. This was a small effect, the difference in study technique accounted for slightly less than 3% of variance the final scores of the students.
- b. The Dean of the university was also interested in your results, as this may help to raise scores in other departments. Unfortunately, she does not understand statistical language. Please interpret your findings for the Dean. Did scores improve? Which technique is better?

Answer (b): We required students to try one of two study techniques between the midterm and the final of our psychology course to determine which technique, if any, was better at improving test scores. We found that both techniques, studying in groups just before the final, and using flashcards to improve memorization, significantly improved test scores. The study techniques accounted for 37% of the improvement measured in test scores. However, we also found that neither study technique was notably better than the other. So students could use either or both techniques to improve scores.

Question 10 (Q10):

- 10. A developmental psychologist is interested in the effect of a positive psychology intervention on the well-being of aging adults. She administers the intervention, collects well-being scores from a sample of 100 participants, and tests whether their well-being differs significantly from the national average. Using G*Power, she determines the power for her test is .80.
 - a. Interpret this value
 - b. What suggestion would you give her if she wants a higher probability of detecting a true effect?

Answer	to	$010 \cdot$
Allswei	υO	ω_{TO} .

This is my answer

Question 11 (Q11):

11. What would it mean if your analysis returned the following values? Consider the meaning of t - not the decision associated with it.

a.
$$t(24) = 0.35$$

b.
$$t(24) = 1.00$$

c.
$$t(24) = 3.2$$

Answer to Q11:

This is my answer

Question 12 (Q12):

12. Draw and annotate all the properties of the null and alternative curves: power, beta, alpha, type 1 error, type 2 error.

Answer to Q12:

See attached hand-drawn image below: