

Pairwise Measures: Four Possibilities for Truth Assignment

- □ Four possibilities based on the agreement between cluster label and partition label
 - □ TP: true positive—Two points \mathbf{x}_i and \mathbf{x}_j belong to the same partition T, and they also in the same cluster C

$$TP = |\{(\mathbf{x}_i, \mathbf{x}_i) : y_i = y_i \text{ and } \hat{y}_i = \hat{y}_i\}|$$

where y_i : the true partition label, and \hat{y}_i : the cluster label for point \mathbf{x}_i

- □ *FN*: false negative: $FN = |\{(\mathbf{x}_i, \mathbf{x}_j) : y_i = y_j \text{ and } \hat{y}_i \neq \hat{y}_j\}|$
- □ *FP: false positive* $FP = |\{(\mathbf{x}_i, \mathbf{x}_j) : y_i \neq y_j \text{ and } \hat{y}_i = \hat{y}_j\}|$
- □ *TN*: true negative $TN = |\{(\mathbf{x}_i, \mathbf{x}_j) : y_i \neq y_j \text{ and } \hat{y}_i \neq \hat{y}_j\}|$
- Calculate the four measures:

$$TP = \sum_{i=1}^{r} \sum_{j=1}^{k} {n_{ij} \choose 2} = \frac{1}{2} \left(\left(\sum_{i=1}^{r} \sum_{j=1}^{k} n_{ij}^{2} \right) - n \right) \quad FN = \sum_{j=1}^{k} {m_{j} \choose 2} - TP$$


$$FP = \sum_{i=1}^{r} {n_{i} \choose 2} - TP \quad TN = N - (TP + FN + FP) = \frac{1}{2} \left(n^{2} - \sum_{i=1}^{r} n_{i}^{2} - \sum_{j=1}^{k} m_{j}^{2} + \sum_{i=1}^{r} \sum_{j=1}^{k} n_{ij}^{2} \right)$$

Pairwise Measures: Jaccard Coefficient and Rand Statistic

- ☐ Jaccard coefficient: Fraction of true positive point pairs, but after ignoring the true negatives (thus asymmetric)
 - □ Jaccard = TP/(TP + FN + FP) [i.e., denominator ignores TN]
 - □ Perfect clustering: Jaccard = 1
- Rand Statistic:
 - \square Rand = (TP + TN)/N
 - Symmetric; perfect clustering: Rand = 1
- **□** Fowlkes-Mallow Measure:
 - Geometric mean of precision and recall

$$FM = \sqrt{prec \times recall} = \frac{TP}{\sqrt{(TP + FN)(TP + FP)}}$$

□ Using the above formulas, one can calculate all the measures for the green table (leave as an exercise)



C\T	T ₁	T ₂	T ₃	Sum
C_1	0	20	30	50
C_2	0	20	5	25
C_3	25	0	0	25
m_j	25	40	35	100