

# BLM2502

## Theory of Computation

# BLM2502 Theory of Computation

## Course Outline

Week	Content
1	Introduction to Course
2	Computability Theory, Complexity Theory, Automata Theory, Set Theory, Relations, Proofs, Pigeonhole Principle
3	Regular Expressions
4	Finite Automata
5	<b>Deterministic and Nondeterministic Finite Automata</b>
6	<b>Epsilon Transition, Equivalence of Automata</b>
7	Pumping Theorem
8	<b>April 10 - 14 week is the first midterm week</b>
9	Context Free Grammars
10	Parse Tree, Ambiguity,
11	Pumping Theorem
12	Turing Machines, Recognition and Computation, Church-Turing Hypothesis
13	Turing Machines, Recognition and Computation, Church-Turing Hypothesis
14	<b>May 22 - 27 week is the second midterm week</b>
15	Review
16	Final Exam date will be announced

Week I - Introduction

# Non-regular languages

(Pumping Lemma)

## Non-regular languages

$$\{a^n b^n : n \geq 0\}$$

$$\{vv^R : v \in \{a,b\}^*\}$$

## Regular languages

$$a^*b$$

$$b^*c + a$$

$$b + c(a + b)^*$$

*etc...*

How can we prove that a language  $L$  is not regular?

Prove that there is no DFA or NFA or RE that accepts  $L$

**Difficulty:** this is not easy to prove  
(since there is an infinite number of them)

**Solution:** use the Pumping Lemma !!!




# The Pigeonhole Principle




4 pigeons



3 pigeonholes



A pigeonhole must  
contain at least two pigeons





$n$  pigeons





.....




$m$  pigeonholes

$n > m$



.....





# The Pigeonhole Principle

$n$  pigeons


$m$  pigeonholes

$n > m$

There is a pigeonhole  
with at least 2 pigeons



.....




# The Pigeonhole Principle

## and


## DFAs

Consider a DFA with 4 states



Consider the walk of a “long” string:  $aaaaab$   
(length at least 4)


A state is repeated in the walk of  $aaaaab$



The state is repeated as a result of the pigeonhole principle

Walk of  $aaaaab$

Pigeons:  
(walk states)



Are more than

Nests:  
(Automaton states)




Repeated  
state


Consider the walk of a “long” string:  $aabb$   
(length at least 4)

Due to the pigeonhole principle:

A state is repeated in the walk of  $aabb$





The state is repeated as a result of the pigeonhole principle






**In General:** If  $|w| \geq \# \text{states of DFA}$ ,  
by the pigeonhole principle,  
a state is repeated in the walk  $w$

Walk of  $w = \sigma_1 \sigma_2 \cdots \sigma_k$




$$|w| \geq \# \text{states of DFA} = m$$



# The Pumping Lemma

Take an **infinite** regular language  $L$   
(contains an infinite number of strings)

There exists a DFA that accepts  $L$




$m$   
states

Take string  $w \in L$  with  $|w| \geq m$

(number of  
states of DFA)


then, at least one state is repeated  
in the walk of  $w$



There could be many states repeated


Take  $q$  to be the first state repeated

One dimensional projection of walk  $w$ :




We can write  $w = xyz$

One dimensional projection of walk  $w$ :




In DFA:  $w = x y z$

contains only  
first occurrence of  $q$





Observation:  $\text{length } |xy| \leq m$  number of states of DFA




Unique States

Since, in  $xy$  no state is repeated (except  $q$ )


Observation:  $\text{length } |y| \geq 1$

Since there is at least one transition in loop




We do not care about the form of string  $z$ .

$z$  may actually overlap with the paths of  $x$  and  $y$




Additional string: The string  $xz$   
is accepted

Do not follow loop




Additional string: The string  $x y y z$  is accepted

Follow loop  
2 times



Additional string: The string  $x y y y z$  is accepted

Follow loop  
3 times



In General:


The string

$x y^i z$

is accepted


$i = 0, 1, 2, \dots$

Follow loop  
 $i$  times





Therefore:  $x y^i z \in L \quad i = 0, 1, 2, \dots$

Language accepted by the DFA








In other words, we described:



The Pumping Lemma !!!



# The Pumping Lemma:

- Given a infinite regular language  $L$
- there exists an integer  $m$  (critical length)
- for any string  $w \in L$  with length  $|w| \geq m$
- we can write  $w = x y z$
- with  $|x y| \leq m$  and  $|y| \geq 1$
- such that:  $x y^i z \in L \quad i = 0, 1, 2, \dots$

In the book:

Critical length  $m$  = Pumping length  $p$

# Applications of the Pumping Lemma

## Observation:

Every language of finite size has to be regular

(we can easily construct an NFA  
that accepts every string in the language)

Therefore, every non-regular language  
has to be of infinite size

(contains an infinite number of strings)

Suppose you want to prove that  
An infinite language  $L$  is not regular

1. Assume the opposite:  $L$  is regular
2. The pumping lemma should hold for  $L$
3. Use the pumping lemma to obtain a contradiction
4. Therefore,  $L$  is not regular

# Explanation of Step 3: How to get a contradiction

1. Let  $m$  be the critical length for  $L$
2. Choose a particular string  $w \in L$  which satisfies the length condition  $|w| \geq m$
3. Write  $w = xyz$
4. Show that  $w' = xy^i z \notin L$  for some  $i \neq 1$
5. This gives a contradiction, since from pumping lemma  $w' = xy^i z \in L$

Note: It suffices to show that  
only one string  $w \in L$   
gives a contradiction

You don't need to obtain  
contradiction for every  $w \in L$



# Example of Pumping Lemma application

**Theorem:** The language  $L = \{a^n b^n : n \geq 0\}$   
is not regular

**Proof:** Use the Pumping Lemma

$$L = \{a^n b^n : n \geq 0\}$$

Assume for contradiction  
that  $L$  is a regular language

Since  $L$  is infinite  
we can apply the Pumping Lemma

$$L = \{a^n b^n : n \geq 0\}$$

Let  $m$  be the critical length for  $L$

Pick a string  $w$  such that:  $w \in L$

and length  $|w| \geq m$

We pick  $w = a^m b^m$

From the Pumping Lemma:

we can write  $w = a^m b^m = x y z$

with lengths  $|x y| \leq m, \quad |y| \geq 1$

$$w = xyz = a^m b^m = \underbrace{a \dots a}_{x} \underbrace{a \dots a}_{y} \underbrace{a \dots a b \dots b}_{z}$$

The diagram illustrates the decomposition of the string  $w = a^m b^m$  into  $xyz$ . The string is represented as  $a \dots a a \dots a a \dots a b \dots b$ . A green bracket above the first two groups of  $a$ 's is labeled  $m$ , and another green bracket above the last group of  $a$ 's and the  $b$ 's is labeled  $m$ . Red brackets below the string partition it into three segments:  $x$  (the first group of  $a$ 's),  $y$  (the second group of  $a$ 's), and  $z$  (the third group of  $a$ 's followed by the  $b$ 's).

Thus:  $y = a^k, \quad 1 \leq k \leq m$

$$x y z = a^m b^m$$

$$y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^i z \in L$   
 $i = 0, 1, 2, \dots$

Thus:  $x y^2 z \in L$

$$x y z = a^m b^m \quad y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^2 z \in L$

$$xy^2z = \overbrace{a \dots a a \dots a a \dots a a \dots a}^{m+k} \overbrace{b \dots b}^m \in L$$

$\underbrace{\hspace{1.5cm}}_x \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{2.5cm}}_z$

**Thus:**  $a^{m+k} b^m \in L$

$$a^{m+k}b^m \in L \quad k \geq 1$$

---

**BUT:**  $L = \{a^n b^n : n \geq 0\}$



$$a^{m+k}b^m \notin L$$

**CONTRADICTION!!!**

Therefore: Our assumption that  $L$   
is a regular language is not true

**Conclusion:**  $L$  is not a regular language

END OF PROOF



Non-regular language  $\{a^n b^n : n \geq 0\}$

Regular languages

$L(a^* b^*)$

# More Applications of the Pumping Lemma

# The Pumping Lemma:

- Given a infinite regular language  $L$
- there exists an integer  $m$  (critical length)
- for any string  $w \in L$  with length  $|w| \geq m$
- we can write  $w = x y z$
- with  $|x y| \leq m$  and  $|y| \geq 1$
- such that:  $x y^i z \in L \quad i = 0, 1, 2, \dots$

Non-regular languages

$$L = \{vv^R : v \in \Sigma^*\}$$



Regular languages

**Theorem:** The language

$$L = \{vv^R : v \in \Sigma^*\} \quad \Sigma = \{a, b\}$$

is not regular

**Proof:** Use the Pumping Lemma

$$L = \{vv^R : v \in \Sigma^*\}$$

Assume for contradiction  
that  $L$  is a regular language

Since  $L$  is infinite  
we can apply the Pumping Lemma

$$L = \{vv^R : v \in \Sigma^*\}$$

Let  $m$  be the critical length for  $L$

Pick a string  $w$  such that:  $w \in L$

and length  $|w| \geq m$

We pick  $w = a^m b^m b^m a^m$

From the Pumping Lemma:

we can write:  $w = a^m b^m b^m a^m = x y z$

with lengths:  $|x y| \leq m, \quad |y| \geq 1$

$$w = xyz = \underbrace{a \dots a}_{x} \underbrace{a \dots a}_{y} \underbrace{a \dots a \dots b \dots b \dots b \dots a \dots a}_{z}$$

$\begin{matrix} m & m & m & m \\ \text{---} & \text{---} & \text{---} & \text{---} \\ & & & \end{matrix}$

Thus:  $y = a^k, \quad 1 \leq k \leq m$



$$x y z = a^m b^m b^m a^m \quad y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^i z \in L$   
 $i = 0, 1, 2, \dots$

Thus:  $x y^2 z \in L$

$$x y z = a^m b^m b^m a^m \quad y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^2 z \in L$

$$xy^2z = \overbrace{a \dots a}^{m+k} \overbrace{a \dots a}^m \overbrace{a \dots a}^m \overbrace{a \dots a}^m \in L$$

$\underbrace{\hspace{1.5cm}}_x \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{4cm}}_z$

**Thus:**  $a^{m+k} b^m b^m a^m \in L$

$$a^{m+k}b^mb^ma^m \in L \quad k \geq 1$$

---

**BUT:**  $L = \{vv^R : v \in \Sigma^*\}$



$$a^{m+k}b^mb^ma^m \notin L$$

**CONTRADICTION!!!**

Therefore: Our assumption that  $L$   
is a regular language is not true

**Conclusion:**  $L$  is not a regular language

END OF PROOF

# Non-regular languages

$$L = \{a^n b^l c^{n+l} : n, l \geq 0\}$$



Regular languages

**Theorem:** The language

$$L = \{a^n b^l c^{n+l} : n, l \geq 0\}$$

is not regular

**Proof:** Use the Pumping Lemma

$$L = \{a^n b^l c^{n+l} : n, l \geq 0\}$$

Assume for contradiction  
that  $L$  is a regular language

Since  $L$  is infinite  
we can apply the Pumping Lemma

$$L = \{a^n b^l c^{n+l} : n, l \geq 0\}$$

Let  $m$  be the critical length of  $L$

Pick a string  $w$  such that:  $w \in L$  and  
length  $|w| \geq m$

We pick  $w = a^m b^m c^{2m}$



From the Pumping Lemma:

We can write  $w = a^m b^m c^{2m} = x y z$

With lengths  $|x y| \leq m, |y| \geq 1$

$$w = xyz = \overbrace{a \dots a}^m \overbrace{a \dots a}^m \overbrace{ab \dots bc \dots cc \dots c}^{2m}$$
$$\underbrace{\hspace{1.5cm}}_x \underbrace{\hspace{1.5cm}}_y \underbrace{\hspace{4.5cm}}_z$$

Thus:  $y = a^k, 1 \leq k \leq m$

$$x y z = a^m b^m c^{2m}$$

$$y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^i z \in L$   
 $i = 0, 1, 2, \dots$

Thus:  $x y^0 z = xz \in L$

$$x y z = a^m b^m c^{2m} \quad y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $xz \in L$

$$xz = \overbrace{a \dots a}^{m-k} \overbrace{a \dots a}^m \overbrace{b \dots b}^m \overbrace{c \dots c}^{2m} \in L$$

$$\underbrace{\hspace{1.5cm}}_x \underbrace{\hspace{4.5cm}}_z$$

Thus:  $a^{m-k} b^m c^{2m} \in L$

$$a^{m-k} b^m c^{2m} \in L \quad k \geq 1$$

---

**BUT:**  $L = \{a^n b^l c^{n+l} : n, l \geq 0\}$



$$a^{m-k} b^m c^{2m} \notin L$$

**CONTRADICTION!!!**

Therefore: Our assumption that  $L$   
is a regular language is not true

**Conclusion:**  $L$  is not a regular language

END OF PROOF

Non-regular languages

$$L = \{a^{n!} : n \geq 0\}$$



Regular languages

**Theorem:** The language  $L = \{a^{n!} : n \geq 0\}$   
is not regular

$$n! = 1 \cdot 2 \cdots (n-1) \cdot n$$

**Proof:** Use the Pumping Lemma

$$L = \{a^{n!} : n \geq 0\}$$

Assume for contradiction  
that  $L$  is a regular language

Since  $L$  is infinite  
we can apply the Pumping Lemma



$$L = \{a^{n!} : n \geq 0\}$$

Let  $m$  be the critical length of  $L$

Pick a string  $w$  such that:  $w \in L$

$$\text{length } |w| \geq m$$

We pick  $w = a^{m!}$

# From the Pumping Lemma:

We can write  $w = a^{m!} = x y z$

With lengths  $|x y| \leq m, \quad |y| \geq 1$

$$w = xyz = a^{m!} = \overbrace{a \dots a}^m \overbrace{a \dots a}^{m!-m}$$
$$\underbrace{\hspace{1.5cm}}_x \underbrace{\hspace{1.5cm}}_y \underbrace{\hspace{4cm}}_z$$

**Thus:**  $y = a^k, \quad 1 \leq k \leq m$

$$x y z = a^{m!}$$

$$y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^i z \in L$   
 $i = 0, 1, 2, \dots$

Thus:  $x y^2 z \in L$

$$x y z = a^{m!}$$

$$y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^2 z \in L$

$$xy^2z = \overbrace{a \dots a a \dots a a \dots a a \dots a}^{m+k} \overbrace{a \dots a}^{m!-m} \in L$$


$\underbrace{\hspace{1.5cm}}_x \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{4cm}}_z$

**Thus:**  $a^{m!+k} \in L$

$$a^{m!+k} \in L \qquad 1 \leq k \leq m$$

---

Since:  $L = \{a^{n!} : n \geq 0\}$



There must exist  $p$  such that:

$$m!+k = p!$$

However:  $m!+k \leq m!+m$  for  $m > 1$

$$\leq m!+m!$$

$$< m!m + m!$$

$$= m!(m+1)$$

$$= (m+1)!$$



$$m!+k < (m+1)!$$



$$m!+k \neq p! \quad \text{for any } p$$

$$a^{m!+k} \in L \quad 1 \leq k \leq m$$

---

**BUT:**  $L = \{a^{n!} : n \geq 0\}$



$$a^{m!+k} \notin L$$

**CONTRADICTION!!!**

Therefore: Our assumption that  $L$   
is a regular language is not true

**Conclusion:**  $L$  is not a regular language

END OF PROOF



# Example 2

**Theorem:**  $B = \{0^n 1^n \mid n \geq 0\}$  is not regular.

**Proof.** Assume the contrary that  $B$  is regular.

Let  $p$  be the pumping length. Then,  $s = 0^p 1^p$  can be decomposed as  $s = xyz$  with  $|y| \geq 1$ , and  $xy^n z \in B$  for any  $n \geq 0$ .

There can be three cases  $y = 0^k$ ,  $y = 0^k 1^l$ , and  $y = 1^l$ , for some nonzero  $k, l$ . In each case, we can easily see that  $xy^n z \notin B$ , which leads to contradiction. ( $n \geq 2$ )

# Example 3

**Theorem:**  $C = \{w \mid w \text{ has an equal number of 0s and 1s}\}$   
*is not regular.*

**Proof.** Assume the contrary that  $C$  is regular.

Let  $p$  be the pumping length. Then,  $s = 0^p 1^p$  can be decomposed as  $s = xyz$  with  $|y| \geq 1$  and  $|xy| \leq p$ , and  $xy^n z \in C$  for any  $n \geq 0$ .

Then we must have  $y = 0^k$  for some nonzero  $k$ .

We can easily see that  $xy^n z \notin C$ , which contradicts the assumption.

# Alternative proof of Example 3

- The class of regular languages is closed under the intersection operation. This is easy to prove if we run two DFAs parallelly and accept only string which are accepted by both of the DFAs.
- Now, assume  $C$  is regular.
- Then,  $C \cap 0^*1^* = B$  is also regular.
- This contradicts what we proved in Example 1.38.???

# Example 4

**Theorem:**  $F = \{ww \mid w \in \{0,1\}^*\}$  is not regular.

**Proof.** Assume the contrary that  $F$  is regular.

Let  $p$  be the pumping length and let  $s = 0^p 1 0^p 1 \in F$ . This  $s$  can be split into pieces like  $s = xyz$  with  $|y| \geq 1$  and  $|xy| \leq p$ , and  $xy^n z \in F$  for any  $n \geq 0$ .

Then we must have  $y = 0^k$  for some nonzero  $k$ .

We can easily see that  $xy^n z \notin F$ , which contradicts the assumption.

# Example 5

**Theorem :**  $D = \{1^{n^2} \mid n \geq 0\}$  is not regular.

**Proof.** Assume the contrary that  $D$  is regular.

Let  $p$  be the pumping length and let  $s = 1^{p^2} \in D$ . This  $s$  can be split into pieces like  $s = xyz$  with  $|y| \geq 1$  and  $|xy| \leq p$ , and  $xy^n z \in D$  for any  $n \geq 0$ .

The length of  $xy^n z$  grows linearly with  $n$ , while the lengths of strings in  $D$  grows as  $0, 1, 4, 9, 16, 25, 36, 49, \dots$

These two facts are incompatible as can be easily seen.

# Example 6 (Pumping Down)

**Theorem:**  $E = \{0^i1^j \mid i > j\}$  is not regular.

**Proof.** Assume the contrary that  $E$  is regular.

Let  $p$  be the pumping length and let  $s = 0^{p+1}1^p$ . This  $s$  can be split into  $s = xyz$  with  $|y| \geq 1$  and  $|xy| \leq p$ , and  $xy^n z \in E$  for any  $n \geq 0$ .

Then we must have  $y = 0^k$  for some nonzero  $k$ .

We can easily see that  $xy^0z = xz \notin E$ , which contradicts the assumption.

# Example 6 (Differential Encoding)

**Claim :**  $D = \{w \mid w \text{ contains equal number of occurrences of the substrings } 01 \text{ and } 10\}$  is regular.

