



JAMDA

journal homepage: www.jamda.com

Review

Proinflammatory Cytokines, Aging, and Age-Related Diseases

Martin Michaud MD^{a,*}, Laurent Balardy MD^a, Guillaume Moulis MD^{b,c,d}, Clement Gaudin MD^a, Caroline Peyrot^e, Bruno Vellas MD, PhD^a, Matteo Cesari MD, PhD^a, Fati Nourhashemi MD, PhD^a

^a Department of Geriatric Medicine, G rontop le de Toulouse, Toulouse University Hospital, Toulouse, France

^b Department of Internal Medicine, Toulouse University Hospital, Toulouse, France

^c Toulouse University, UMR1027, Toulouse, France

^d Inserm, UMR1027, Toulouse, France

^e Direction of Research and Innovation, Toulouse University Hospital, Toulouse, France

A B S T R A C T

Keywords:

Inflammation
aging
cytokines
frailty
Alzheimer disease
Interleukin-6
TNF- 

Inflammation is a physiological process that repairs tissues in response to endogenous or exogenous aggressions. Nevertheless, a chronic state of inflammation may have detrimental consequences. Aging is associated with increased levels of circulating cytokines and proinflammatory markers. Aged-related changes in the immune system, known as immunosenescence, and increased secretion of cytokines by adipose tissue, represent the major causes of chronic inflammation. This phenomenon is known as "inflamm-aging." High levels of interleukin (IL)-6, IL-1, tumor necrosis factor- , and C-reactive protein are associated in the older subject with increased risk of morbidity and mortality. In particular, cohort studies have indicated TNF-  and IL-6 levels as markers of frailty. The low-grade inflammation characterizing the aging process notably concurs at the pathophysiological mechanisms underlying sarcopenia. In addition, proinflammatory cytokines (through a variety of mechanisms, such as platelet activation and endothelial activation) may play a major role in the risk of cardiovascular events. Dysregulation of the inflammatory pathway may also affect the central nervous system and be involved in the pathophysiological mechanisms of neurodegenerative disorders (eg, Alzheimer disease). The aim of the present review was to summarize different targets of the activity of proinflammatory cytokines implicated in the risk of pathological aging.

Copyright   2013 - American Medical Directors Association, Inc.

The improvement of knowledge about the mechanisms of aging represents an important goal of current research. It is estimated that the population aged 65 years and older will present a 3.5-fold increase by 2025 to 2030 compared with other age groups.¹ Among the biological mechanisms more likely to affect the distinction between a successful and pathological aging process, inflammation surely is one of the most studied.

Aging is characterized by quantitative and qualitative modifications of the immune system. This phenomenon, known as "immunosenescence," is accompanied by cytokine dysregulation, which is an increase of proinflammatory cytokines and reduction of anti-inflammatory cytokines, leading to a chronic low-grade inflammatory state.¹ The potential clinical consequences of such condition are relevant, including the increased risk of comorbidities and mortality. In other words, inflammation may constitute a biological foundation of the pathophysiological process of frailty.¹ Bone, nutritional, and

muscle metabolisms are all affected by the inflammatory state accompanying aging.^{2–5} The pathophysiology of neurodegenerative disorders (such as Alzheimer disease [AD]) also seems to be partly inflammatory and influenced by these changes in cytokine secretions.⁶ The well-established relationship between systemic inflammation and cardiovascular events is also noteworthy.⁷ On the other hand, the lack of alteration of inflammatory biomarkers appears to be associated with "successful aging."

The aim of the present article was to review the literature on the relationship between proinflammatory cytokines and "pathological aging."

Origin of Low-Grade Inflammation

Immunosenescence

Immunosenescence is a progressive modification of the immune system that leads to greater susceptibility to infections, neoplasias, and autoimmune manifestations. This phenomenon is mainly due to prolonged antigenic stimulation across the life span.⁸ The main alteration responsible for immunosenescence concerns the

The authors declare no conflicts of interest.

* Address correspondence to Martin Michaud, MD, Department of Internal Medicine and Gerontology, 170 Avenue de Casselardit, 31300 Toulouse, France.

E-mail address: martin.michaud85@gmail.com (M. Michaud).

1525-8610/\$ - see front matter Copyright   2013 - American Medical Directors Association, Inc.

<http://dx.doi.org/10.1016/j.jamda.2013.05.009>

functioning of T cells.⁹ Consequences of the thymic involution occurring with aging are decreases of the regulatory T lymphocytes and naïve T lymphocytes (in favor of memory T lymphocytes).¹⁰ Such reorganization of the immune system contributes to modifying the profile of cytokine secretion, causing “inflamm-aging.”¹¹

With aging, the level of type 1 and 2 cytokines increases in T lymphocytes.^{12,13} T-helper (Th) lymphocytes are cells that are central to the immune response and are strongly involved in cytokine synthesis. After activation, Th lymphocytes differentiate into Th1, Th2, Th9, Th17, follicular helper T (Tfh), or memory T lymphocytes.¹⁴ The age-related dysregulation leads to the abnormal distribution of the subtypes Th1 and Th2 (ie, Th1–Th2 commutation), with a consequent relative increase of the number of Th2 lymphocytes compared with Th1.^{15,16} It is noteworthy that the profile of cytokine secretion is different between these 2 subtypes of Th lymphocytes.¹⁷ Plasma levels of proinflammatory Th2 cytokines and their soluble receptors are higher in aged than in young subjects.^{16,18–31}

Other Factors

Although immunosenescence appears to be the principal factor at the origin of low-grade inflammation, the following factors may also play a role.

Adipose tissue, through the intermediary of the macrophages that it contains, produces IL-6, tumor necrosis factor (TNF)- α , and adipokines.³² A large amount of visceral adipose tissue or a high-fat diet is correlated with an increase in plasma levels of C-reactive protein (CRP) and interleukin (IL)-6 in obese individuals, and also in individuals who are not obese.^{33,34} Aging is accompanied by a decline in sex hormones in both sexes. The role of androgens and estrogens in low-grade inflammation has been confirmed *in vitro* by their ability to suppress transcription of the IL-6 gene.³⁵ Chronic periodontitis, frequent in the elderly, contributes to systemic inflammation.³⁶ And, finally, chronic exposure to stress, through the intermediary of increased endogenous secretion of glucocorticoids, has a proinflammatory influence on T lymphocyte function.³⁷

Impacts of Aging on Low-Grade Inflammation

Cytokines, Morbidity, and Mortality

Elevated levels of TNF- α , IL-6, and CRP are correlated with an increased risk of morbidity and mortality (all causes together), not only in a frail population (defined according to Fried et al's criteria³⁸) but also in independent, nonfrail elderly.^{39–49} The effects of cytokine dysregulation are independent of the other usual mortality risk factors (tobacco use, diabetes, arterial hypertension, hypercholesterolemia) and comorbid conditions.^{40,42,45,46,50}

Cytokines and Frailty

Some studies have shown higher levels of IL-6, TNF- α , and CRP in the elderly frail population (as defined by Fried et al³⁸) than in the control population of the same age.^{46,51–57} Elevated levels of IL-6 are associated with slower gait velocity and are predictive of gait speed decline in the medium term in community-dwelling older persons.⁵⁸ Patients with impaired functioning in the activities of daily living (ADLs)⁵⁹ have higher plasma IL-6 levels than patients who are not disabled.^{41,47} Various mechanisms could explain this close link between frailty syndrome and increased proinflammatory cytokines. Current and future research will enable us to study the impact of numerous other mechanisms that have not yet been investigated.

Cytokines and Nutritional Status

Involuntary weight loss is an independent factor of morbidity and mortality.⁶⁰ Proinflammatory cytokines have a direct influence on the development of malnutrition.^{61,62} Increase in levels of TNF- α , IL-1, and IL-6 has metabolic consequences with increased lipolysis, decreased lipid synthesis, decreased lipoprotein lipase activity, decreased protein synthesis, and increased proteolysis.^{3,5,63} In addition, TNF- α acts directly on the digestive system by reducing gastric filling capacity and slowing peristaltic activity.⁶⁴

As well as such direct actions, cytokines induce malnutrition through changes in the hormonal secretions involved in satiety (changes in cholecystokinin, glucagon, and adrenocorticotrophic hormone secretion).^{65–67} The central nervous system also intervenes in the mechanism of malnutrition, in particular the hypothalamic nuclei, which express cytokine receptors that are activated in response to systemic inflammation.^{68–70}

Cytokines and Muscle Mass

Physiologically, muscle mass decreases by 1% to 2% yearly after the fifth decade of life.⁷¹ *In vitro*, exposure of myoblasts to TNF- α results in decreased production of myofilament proteins and in regulation of key transcription factors involved in differentiation of striated muscle cells.⁷² The inflammatory syndrome, and TNF- α in particular, in fact causes increased muscle catabolism.⁷³ In murine models, inhibition of inflammation pathways promotes muscle regeneration.⁷⁴ In addition, the decrease in myofibrillar contractility is partly mediated by the increase in oxidative stress and nitric acid production, both of which are promoted by the increase in TNF- α levels.^{73,75} By contrast, muscle training promotes the decrease of TNF- α levels.^{76,77}

In subjects older than 65 years, elevated plasma levels of IL-6, TNF- α , and/or CRP are associated with lower physical performance, muscle strength, and muscle mass.^{2,77–82} Plasma TNF- α level is moreover predictive of a significant decrease in muscle strength at 4 years in subjects aged 85 years⁸³ and at 5 years in subjects aged 70 to 79 years.⁸⁴ In women in good health aged older than 65 years with femoral neck fracture, the lowest levels of IL-6 were significantly correlated with better functional and muscular recovery in the year after fracture.^{85,86}

Cytokines and Bone Density

Bone remodeling and resorption are in part modulated by proinflammatory cytokines.^{4,87} IL-6 stimulates osteoclastogenesis and osteoclastic activity, while shortening osteoblast survival.^{88–93} Elevated levels of CRP, IL-6, and TNF- α are associated with increased risk of osteoporosis.^{94–97} These markers of inflammation are also predictive of fracture risk in both sexes.^{98–100}

Cytokines and Vascular Risk

Atherosclerosis begins early in the life of an individual with penetration and accumulation of low-density lipoprotein (LDL) cholesterol in the intima. This lipid infiltration is followed by oxidative change of the LDL that is responsible for endothelial dysfunction, causing subendothelial infiltration by inflammatory cells (macrophages, lymphocytes T) attracted by chemotaxis.¹⁰¹ These cells then release proinflammatory cytokines (mainly IL-1, IL-6, TNF- α , and transforming growth factor- β). The result is overexpression of adhesion molecules and procoagulant agents.^{102,103} TNF- α , IL-1, and INF- γ are proapoptotic inducers of smooth muscle cells.¹⁰³ Adding to these effects is the local arterial vasoconstriction induced by TNF- α , IL-6, and IL-10, through their impact on the endothelium and through synthesis of nitric acid.¹⁰⁴ Inflammatory cytokines thus participate in both the growth of the plaque and its fragilization.

In addition, IL-1, IL-2, IL-6, and TNF- α directly influence coagulation pathways. They modify endothelial function, leading to a prothrombotic state with inhibition of fibrinolysis, increased production of platelet activation factors, and activation of the intrinsic and extrinsic coagulation pathways.^{105,106}

The presence of conventional cardiovascular risk factors (tobacco use, arterial hypertension, dyslipidemia, diabetes, obesity) is accompanied by increased plasma levels of TNF- α and CRP.^{107–112}

In the aged population, the relative weight of traditional risk factors in the occurrence of cardiovascular and cerebrovascular events is lower than in the nonelderly population.⁷

The inflammatory syndrome thus seems to have a major influence. In the elderly population, high levels of CRP, fibrinogen, TNF- α , and IL-6 are independent risk factors for cerebrovascular and cardiovascular events.^{42,45,108,110,113–116}

Cytokines and Cognitive Decline

Cytokines are produced within the brain, and in addition, peripheral cytokines have a direct influence on the central nervous system, as reviewed by Rosano et al.¹¹⁷ Peripheral cytokines are capable of inducing central neurological disturbances, as in the course of sickness behavior.¹¹⁸ The endothelial cells of the blood-brain barrier are able to produce cytokines or inflammatory molecules (prostaglandin and nitric acid) in response to an increase in the level of peripheral cytokines.¹¹⁹ Cytokines are also produced within the brain, principally by activated microglia that are capable of synthesizing more than 20 cytokines.¹²⁰

In populations free of cognitive disturbance, high levels of IL-6, CRP, or TNF- α have been found to be markers of decline in cognitive performance in the medium term.^{121–128} Other studies, however, found no significant link.^{129,130}

Cytokines and AD

Inflammation is part of the pathophysiology of AD.¹³¹ Postmortem histopathological examination of the brains of patients with AD showed high levels of proinflammatory cytokines, CRP, and complement proteins in senile plaques and neurofibrillary tangles.^{132,133}

Peripheral cytokines can influence the onset or the cognitive decline of AD.^{6,134} The influence of proinflammatory cytokines has been well argued on the basis of study of genetic polymorphisms. Certain polymorphisms of the TNF- α , IL-1, and IL-6 genes increase the risk of developing AD.^{135,136} High levels of IL-1 and TNF- α have been found to be potential predictive markers of development of AD at 7 years.¹³⁴

In patients with mild or moderate AD, plasma levels of IL1, IL-6, or TNF- α have been found to be higher than in healthy individuals.^{27,137–141} However, other studies did not confirm this relationship.^{142–145}

Higher intrathecal levels of IL-6 have been observed in patients with AD.^{146,147} In addition, hippocampal volume may be inversely correlated with plasma IL-6 level in healthy middle-aged adults.¹⁴⁸ Cognitive decline may be more rapid in patients with high levels of TNF- α .¹⁴⁹

Cytokines and Cognitive Decline of Vascular Origin

Cytokines also play a damaging role in the pathophysiology of vascular dementia. They are atherogenic and prothrombotic factors and can thus directly influence the occurrence of ischemic cerebrovascular events.¹⁵⁰ In addition, an exaggerated cytokine response to ischemia worsens cell damage. Some studies found a correlation between elevated levels of CRP and IL-6 and vascular

Table 1

Proinflammatory Cytokines, Aging, and Age-related Diseases: Principal Cytokines Involved and Pathophysiology

Principal Cytokines Involved	Pathophysiology
Cytokines and Cognitive Decline TNF- α , IL-6, IL-1	Genetic polymorphism and AD risk ^{135,136} Decrease in hippocampal volume ¹⁴⁸ Microglial activation ¹²⁰ Complement activation ^{132,133} Oxidative stress ¹¹⁹
Cytokines and Vascular Aging CRP, TNF- α , IL-6	Endothelial activation ^{105,106} Inhibition of fibrinolysis ^{105,106} Prothrombotic activity ^{105,106} Platelet activation ^{105,106} Activation of coagulation ¹⁰⁶ Atherosclerosis ^{101–103}
Cytokines and Nutritional Status TNF- α , IL-1, IL-6, INF- γ	↓ lipolysis, ↓ lipid synthesis ^{3,5,63} ↓ lipoprotein lipase activity ^{3,5,63} ↓ protein synthesis, ↑ proteolysis ^{3,5,63} ↑ anorexigenic hormones ^{65–67} ↓ gastric filling capacity ⁶⁴ ↓ peristaltic activity ⁶⁴
Cytokines and Bone Metabolism TNF- α , IL-6	↑ bone remodelling ^{4,87} ↑ bone resorption ^{4,87} ↑ osteoclastogenesis ^{4,87} ↓ osteoblast survival ^{88–93}
Cytokines and Muscle Metabolism TNF- α , IL-6	↓ myofibrillar protein synthesis ⁷² ↑ oxidative stress ^{73,75} ↑ muscle catabolism ⁷³

AD, Alzheimer disease; CRP, C-reactive protein; IL, interleukin; INF, interferon; TNF, tumor necrosis factor.

dementia.^{126,127,142} However, this correlation was not found in all studies.¹⁵¹

Conclusion

Aging is accompanied by immune, hormonal, and adipose changes leading to a chronic inflammatory state. Levels of proinflammatory cytokines, principally TNF- α and IL-6, have damaging effects on aging (Table 1). They influence the onset of frailty, cognitive decline, and cardiometabolic, neurological, and vascular events. They are also closely linked to the genesis of cancers, with cardiac remodeling in heart failure, and so forth. The limitations of this review arise from the numerous methodological factors of confusion. In fact, studies are not comparable, as measurement methods and threshold values differ and populations are not homogeneous from one study to another. Also, much work is still required, in particular with regard to the influence of multimodal management on levels of proinflammatory cytokines and on the progressive course of these diseases. Simple measures, such as physical activity and a moderate-fat diet, have always been recognized as having a beneficial effect on aging by reducing, in particular, the degree of low-grade inflammation.

Acknowledgments

The authors thank Dr Christian Michaud and Marie Josée Michaud (Sables d'Olonne, France).

References

1. Miller RA. The aging immune system: Primer and prospectus. *Science* 1996; 273:70–74.
2. Cesari M, Penninx BW, Pahor M, et al. Inflammatory markers and physical performance in older persons: The INCHIANTI study. *J Gerontol A Biol Sci Med Sci* 2004;59:242–248.

3. Grunfeld C, Feingold KR. The metabolic effects of tumor necrosis factor and other cytokines. *Biotherapy* 1991;3:143–158.
4. Manolagas SC, Jilka RL. Bone marrow, cytokines, and bone remodeling. Emerging insights into the pathophysiology of osteoporosis. *N Engl J Med* 1995;332:305–311.
5. Yeh SS, Schuster MW. Geriatric cachexia: The role of cytokines. *Am J Clin Nutr* 1999;70:183–197.
6. Akiyama H, Barger S, Barnum S, et al. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000;21:383–421.
7. Cesari M, Penninx BW, Newman AB, et al. Inflammatory markers and onset of cardiovascular events: Results from the Health ABC study. *Circulation* 2003;108:2317–2322.
8. Gruver AL, Hudson LL, Sempowski GD. Immunosenescence of ageing. *J Pathol* 2007;211:144–156.
9. Linton P, Thoman ML. T cell senescence. *Front Biosci* 2001;6:D248–D261.
10. Weiskopf D, Weinberger B, Grubeck-Loebenstein B. The aging of the immune system. *Transpl Int* 2009;22:1041–1050.
11. Franceschi C, Bonafe M, Valensin S, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* 2000;908:244–254.
12. Zanni F, Vescovini R, Biasini C, et al. Marked increase with age of type 1 cytokines within memory and effector/cytotoxic CD8+ T cells in humans: A contribution to understand the relationship between inflammation and immunosenescence. *Exp Gerontol* 2003;38:981–987.
13. Alberti S, Cevenini E, Ostan R, et al. Age-dependent modifications of Type 1 and Type 2 cytokines within virgin and memory CD4+ T cells in humans. *Mech Ageing Dev* 2006;127:560–566.
14. Hirahara K, Vahedi K, Ghoreschi K, et al. Helper T-cell differentiation and plasticity: Insights from epigenetics. *Immunology* 2011;134:235–245.
15. Shearer GM. Th1/Th2 changes in aging. *Mech Ageing Dev* 1997;94:1–5.
16. Cakman I, Rohwer J, Schutz RM, et al. Dysregulation between TH1 and TH2 T cell subpopulations in the elderly. *Mech Ageing Dev* 1996;87:197–209.
17. Mosmann TR, Coffman RL. TH1 and TH2 cells: Different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 1989;7:145–173.
18. Gerli R, Monti D, Bistoni O, et al. Chemokines, sTNF-Rs and sCD30 serum levels in healthy aged people and centenarians. *Mech Ageing Dev* 2000;121:37–46.
19. Paganelli R, Scala E, Rosso R, et al. A shift to Th0 cytokine production by CD4+ cells in human longevity: Studies on two healthy centenarians. *Eur J Immunol* 1996;26:2030–2034.
20. Wei J, Xu H, Davies JL, Hemmings GP. Increase of plasma IL-6 concentration with age in healthy subjects. *Life Sci* 1992;51:1953–1956.
21. Ershler WB. Interleukin-6: A cytokine for gerontologists. *J Am Geriatr Soc* 1993;41:176–181.
22. Ershler WB, Sun WH, Binkley N, et al. Interleukin-6 and aging: Blood levels and mononuclear cell production increase with advancing age and in vitro production is modifiable by dietary restriction. *Lymphokine Cytokine Res* 1993;12:225–230.
23. Hager K, Machein U, Krieger S, et al. Interleukin-6 and selected plasma proteins in healthy persons of different ages. *Neurobiol Aging* 1994;15:771–772.
24. Forsey RJ, Thompson JM, Ernerudh J, et al. Plasma cytokine profiles in elderly humans. *Mech Ageing Dev* 2003;124:487–493.
25. Young DG, Skibinski G, Mason JI, James K. The influence of age and gender on serum dehydroepiandrosterone sulphate (DHEA-S), IL-6, IL-6 soluble receptor (IL-6 sR) and transforming growth factor beta 1 (TGF-beta1) levels in normal healthy blood donors. *Clin Exp Immunol* 1999;117:476–481.
26. Straub RH, Konecna L, Hrach S, et al. Serum dehydroepiandrosterone (DHEA) and DHEA sulfate are negatively correlated with serum interleukin-6 (IL-6), and DHEA inhibits IL-6 secretion from mononuclear cells in man in vitro: Possible link between endocrinosenescence and immunosenescence. *J Clin Endocrinol Metab* 1998;83:2012–2017.
27. Bruunsgaard H, Andersen-Ranberg K, Jeune B, et al. A high plasma concentration of TNF-alpha is associated with dementia in centenarians. *J Gerontol A Biol Sci Med Sci* 1999;54:M357–M364.
28. Paolisso G, Rizzo MR, Mazziotti G, et al. Advancing age and insulin resistance: Role of plasma tumor necrosis factor-alpha. *Am J Physiol* 1998;275:E294–E299.
29. Hallgren HM, Bergh N, Rodysill KJ, O'Leary JJ. Lymphocyte proliferative response to PHA and anti-CD3/Ti monoclonal antibodies, T cell surface marker expression, and serum IL-2 receptor levels as biomarkers of age and health. *Mech Ageing Dev* 1988;43:175–185.
30. Rabinovich H, Goses Y, Reshef T, Klajman A. Interleukin-2 production and activity in aged humans. *Mech Ageing Dev* 1985;32:213–226.
31. Born J, Uthgenannt D, Dodt C, et al. Cytokine production and lymphocyte subpopulations in aged humans. An assessment during nocturnal sleep. *Mech Ageing Dev* 1995;84:113–126.
32. Trayhurn P, Wood IS. Signalling role of adipose tissue: Adipokines and inflammation in obesity. *Biochem Soc Trans* 2005;33:1078–1081.
33. Pou KM, Massaro JM, Hoffmann U, et al. Visceral and subcutaneous adipose tissue volumes are cross-sectionally related to markers of inflammation and oxidative stress: The Framingham Heart Study. *Circulation* 2007;116:1234–1241.
34. Nappo F, Esposito K, Cioffi M, et al. Postprandial endothelial activation in healthy subjects and in type 2 diabetic patients: Role of fat and carbohydrate meals. *J Am Coll Cardiol* 2002;39:1145–1150.
35. Manolagas SC, Jilka RL, Girasole G, et al. Estrogen, cytokines, and the control of osteoclast formation and bone resorption in vitro and in vivo. *Osteoporos Int* 1993;3:114–116.
36. Paraskevas S, Huizinga JD, Loos BG. A systematic review and meta-analyses on C-reactive protein in relation to periodontitis. *J Clin Periodontol* 2008;35:277–290.
37. Bauer ME. Chronic stress and immunosenescence: A review. *Neuro-immunomodulation* 2008;15:241–250.
38. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: Evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146–M156.
39. Alley DE, Crimmins E, Bandeen-Roche K, et al. Three-year change in inflammatory markers in elderly people and mortality: The Invecchiare in Chianti study. *J Am Geriatr Soc* 2007;55:1801–1807.
40. Bruunsgaard H, Andersen-Ranberg K, Hjelmborg JB, et al. Elevated levels of tumor necrosis factor alpha and mortality in centenarians. *Am J Med* 2003;115:278–283.
41. Ferrucci L, Harris TB, Guralnik JM, et al. Serum IL-6 level and the development of disability in older persons. *J Am Geriatr Soc* 1999;47:639–646.
42. Harris TB, Ferrucci L, Tracy RP, et al. Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. *Am J Med* 1999;106:506–512.
43. Mooradian AD, Reed RL, Osterweil D, Scuderi P. Detectable serum levels of tumor necrosis factor alpha may predict early mortality in elderly institutionalized patients. *J Am Geriatr Soc* 1991;39:891–894.
44. Roubenoff R, Parise H, Payette HA, et al. Cytokines, insulin-like growth factor 1, sarcopenia, and mortality in very old community-dwelling men and women: The Framingham Heart Study. *Am J Med* 2003;115:429–435.
45. Volpato S, Guralnik JM, Ferrucci L, et al. Cardiovascular disease, interleukin-6, and risk of mortality in older women: The women's health and aging study. *Circulation* 2001;103:947–953.
46. Reuben DB, Cheh AI, Harris TB, et al. Peripheral blood markers of inflammation predict mortality and functional decline in high-functioning community-dwelling older persons. *J Am Geriatr Soc* 2002;50:638–644.
47. Cohen HJ, Pieper CF, Harris T, et al. The association of plasma IL-6 levels with functional disability in community-dwelling elderly. *J Gerontol A Biol Sci Med Sci* 1997;52:M201–M208.
48. Newman AB, Sachs MC, Arnold AM, et al. Total and cause-specific mortality in the cardiovascular health study. *J Gerontol A Biol Sci Med Sci* 2009;64:1251–1261.
49. Walston JD, Matteini AM, Nievergelt C, et al. Inflammation and stress-related candidate genes, plasma interleukin-6 levels, and longevity in older adults. *Exp Gerontol* 2009;44:350–355.
50. Bruunsgaard H, Ladelund S, Pedersen AN, et al. Predicting death from tumour necrosis factor-alpha and interleukin-6 in 80-year-old people. *Clin Exp Immunol* 2003;132:24–31.
51. Barzilay JI, Blaum C, Moore T, et al. Insulin resistance and inflammation as precursors of frailty: The Cardiovascular Health Study. *Arch Intern Med* 2007;167:635–641.
52. Cappola AR, Xue QL, Ferrucci L, et al. Insulin-like growth factor I and interleukin-6 contribute synergistically to disability and mortality in older women. *J Clin Endocrinol Metab* 2003;88:2019–2025.
53. Leng SX, Xue QL, Tian J, et al. Inflammation and frailty in older women. *J Am Geriatr Soc* 2007;55:864–871.
54. Walston J, McBurnie MA, Newman A, et al. Frailty and activation of the inflammation and coagulation systems with and without clinical comorbidities: Results from the Cardiovascular Health Study. *Arch Intern Med* 2002;162:2333–2341.
55. Reuben DB, Judd-Hamilton L, Harris TB, Seeman TE. The associations between physical activity and inflammatory markers in high-functioning older persons: MacArthur Studies of Successful Aging. *J Am Geriatr Soc* 2003;51:1125–1130.
56. Ronning B, Wyller TB, Seljeflot I, et al. Frailty measures, inflammatory biomarkers and post-operative complications in older surgical patients. *Age Ageing* 2010;39:758–761.
57. Hubbard RE, O'Mahony MS, Savva GM, et al. Inflammation and frailty measures in older people. *J Cell Mol Med* 2009;13:3103–3109.
58. Verghese J, Holtzer R, Oh-Park M, et al. Inflammatory markers and gait speed decline in older adults. *J Gerontol A Biol Sci Med Sci* 2011;66:1083–1089.
59. Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. *Gerontologist* 1970;10:20–30.
60. Wallace JL, Schwartz RS, LaCroix AZ, et al. Involuntary weight loss in older outpatients: Incidence and clinical significance. *J Am Geriatr Soc* 1995;43:329–337.
61. Liao Z, Tu JH, Small CB, et al. Increased urine interleukin-1 levels in aging. *Gerontology* 1993;39:19–27.
62. Cederholm T, Wretling B, Hellstrom K, et al. Enhanced generation of interleukins 1 beta and 6 may contribute to the cachexia of chronic disease. *Am J Clin Nutr* 1997;65:876–882.
63. Keller U. Pathophysiology of cancer cachexia. *Support Care Cancer* 1993;1:290–294.
64. Bodnar RJ, Pasternak GW, Mann PE, et al. Mediation of anorexia by human recombinant tumor necrosis factor through a peripheral action in the rat. *Cancer Res* 1989;49:6280–6284.
65. Smith GP. Gut hormone hypothesis of postprandial satiety. *Res Publ Assoc Res Nerv Ment Dis* 1984;62:67–75.

66. Moran TH, McHugh PR. Cholecystokinin suppresses food intake by inhibiting gastric emptying. *Am J Physiol* 1982;242:R491–R497.
67. Le Sauter J, Geary N. Hepatic portal glucagon infusion decreases spontaneous meal size in rats. *Am J Physiol* 1991;261:154–161.
68. Turrin NP, Plata-Salaman CR. Cytokine-cytokine interactions and the brain. *Brain Res Bull* 2000;51:3–9.
69. Somm E, Henrichot E, Pernin A, et al. Decreased fat mass in interleukin-1 receptor antagonist-deficient mice: Impact on adipogenesis, food intake, and energy expenditure. *Diabetes* 2005;54:3503–3509.
70. Utsuyama M, Hirokawa K. Differential expression of various cytokine receptors in the brain after stimulation with LPS in young and old mice. *Exp Gerontol* 2002;37:411–420.
71. Sehl ME, Yates FE. Kinetics of human aging: I. Rates of senescence between ages 30 and 70 years in healthy people. *J Gerontol A Biol Sci Med Sci* 2001;56:198–208.
72. Langen RC, Van Der Velden JL, Schols AM, et al. Tumor necrosis factor- α inhibits myogenic differentiation through MyoD protein destabilization. *FASEB J* 2004;18:227–237.
73. Reid MB, Li YP. Tumor necrosis factor- α and muscle wasting: A cellular perspective. *Respir Res* 2001;2:269–272.
74. Mourikioti F, Kratsios P, Luedde T, et al. Targeted ablation of IKK2 improves skeletal muscle strength, maintains mass, and promotes regeneration. *J Clin Invest* 2006;116:2945–2954.
75. Reid MB, Lannergren J, Westerblad H. Respiratory and limb muscle weakness induced by tumor necrosis factor- α : Involvement of muscle myofilaments. *Am J Respir Crit Care Med* 2002;166:479–484.
76. Starkie R, Ostrowski SR, Jauffred S, et al. Exercise and IL-6 infusion inhibit endotoxin-induced TNF- α production in humans. *FASEB J* 2003;17:884–886.
77. Greiwe JS, Cheng B, Rubin DC, et al. Resistance exercise decreases skeletal muscle tumor necrosis factor α in frail elderly humans. *FASEB J* 2001;15:475–482.
78. Visser M, Pahor M, Taaffe DR, et al. Relationship of interleukin-6 and tumor necrosis factor- α with muscle mass and muscle strength in elderly men and women: The Health ABC Study. *J Gerontol A Biol Sci Med Sci* 2002;57:M326–M332.
79. Pedersen M, Bruunsgaard H, Weis N, et al. Circulating levels of TNF- α and IL-6—relation to truncal fat mass and muscle mass in healthy elderly individuals and in patients with type-2 diabetes. *Mech Ageing Dev* 2003;124:495–502.
80. Barbieri M, Ferrucci L, Ragno E, et al. Chronic inflammation and the effect of IGF-I on muscle strength and power in older persons. *Am J Physiol Endocrinol Metab* 2003;284:E481–E487.
81. Ferrucci L, Penninx BW, Volpato S, et al. Change in muscle strength explains accelerated decline of physical function in older women with high interleukin-6 serum levels. *J Am Geriatr Soc* 2002;50:1947–1954.
82. Stenholm S, Maggio M, Lauretani F, et al. Anabolic and catabolic biomarkers as predictors of muscle strength decline: The INCHIANTI study. *Rejuvenation Res* 2010;13:3–11.
83. Taekema DG, Westendorp RG, Frolich M, Gussekloo J. High innate production capacity of tumor necrosis factor- α and decline of handgrip strength in old age. *Mech Ageing Dev* 2007;128:517–521.
84. Schaap LA, Pluijm SM, Deeg DJ, et al. Higher inflammatory marker levels in older persons: Associations with 5-year change in muscle mass and muscle strength. *J Gerontol A Biol Sci Med Sci* 2009;64:1183–1189.
85. Miller RR, Shardell MD, Hicks GE, et al. Association between interleukin-6 and lower extremity function after hip fracture—the role of muscle mass and strength. *J Am Geriatr Soc* 2008;56:1050–1056.
86. Miller RR, Cappola AR, Shardell MD, et al. Persistent changes in interleukin-6 and lower extremity function following hip fracture. *J Gerontol A Biol Sci Med Sci* 2006;61:1053–1058.
87. Ding C, Parameswaran V, Udayan R, et al. Circulating levels of inflammatory markers predict change in bone mineral density and resorption in older adults: A longitudinal study. *J Clin Endocrinol Metab* 2008;93:1952–1958.
88. Clowes JA, Riggs BL, Khosla S. The role of the immune system in the pathophysiology of osteoporosis. *Immunol Rev* 2005;208:207–227.
89. Ishimi Y, Miyaura C, Jin CH, et al. IL-6 is produced by osteoblasts and induces bone resorption. *J Immunol* 1990;145:3297–3303.
90. Manolagas SC. Birth and death of bone cells: Basic regulatory mechanisms and implications for the pathogenesis and treatment of osteoporosis. *Endocr Rev* 2000;21:115–137.
91. Tamura T, Udagawa N, Takahashi N, et al. Soluble interleukin-6 receptor triggers osteoclast formation by interleukin 6. *Proc Natl Acad Sci U S A* 1993;90:11924–11928.
92. Black K, Garrett IR, Mundy GR. Chinese hamster ovarian cells transfected with the murine interleukin-6 gene cause hypercalcemia as well as cachexia, leukocytosis and thrombocytosis in tumor-bearing nude mice. *Endocrinology* 1991;128:2657–2659.
93. Kurihara N, Civin C, Roodman GD. Osteotropic factor responsiveness of highly purified populations of early and late precursors for human multinucleated cells expressing the osteoclast phenotype. *J Bone Miner Res* 1991;6:257–261.
94. Zheng SX, Vrindts Y, Lopez M, et al. Increase in cytokine production (IL-1 β , IL-6, TNF- α but not IFN- γ , GM-CSF, or LIF) by stimulated whole blood cells in postmenopausal osteoporosis. *Maturitas* 1997;26:63–71.
95. Khosla S, Peterson JM, Egan K, et al. Circulating cytokine levels in osteoporotic and normal women. *J Clin Endocrinol Metab* 1994;79:707–711.
96. Abrahamsen B, Bonnevie-Nielsen V, Ebbesen EN, et al. Cytokines and bone loss in a 5-year longitudinal study—hormone replacement therapy suppresses serum soluble interleukin-6 receptor and increases interleukin-1 receptor antagonist: The Danish Osteoporosis Prevention Study. *J Bone Miner Res* 2000;15:1545–1554.
97. Koh JM, Khang YH, Jung CH, et al. Higher circulating hsCRP levels are associated with lower bone mineral density in healthy pre- and postmenopausal women: Evidence for a link between systemic inflammation and osteoporosis. *Osteoporos Int* 2005;16:1263–1271.
98. Barbour KE, Boudreau R, Danielson ME, et al. Inflammatory markers and the risk of hip fracture: The Women's Health Initiative. *J Bone Miner Res* 2012;27:1167–1176.
99. Pasco JA, Kotowicz MA, Henry MJ, et al. High-sensitivity C-reactive protein and fracture risk in elderly women. *JAMA* 2006;296:1353–1355.
100. Cauley JA, Danielson ME, Boudreau RM, et al. Inflammatory markers and incident fracture risk in older men and women: The Health Aging and Body Composition Study. *J Bone Miner Res* 2007;22:1088–1095.
101. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999;340:115–126.
102. Stern DM, Kaiser E, Nawroth PP. Regulation of the coagulation system by vascular endothelial cells. *Haemostasis* 1988;18:202–214.
103. Arvin B, Neville LF, Barone FC, Feuerstein GZ. The role of inflammation and cytokines in brain injury. *Neurosci Biobehav Rev* 1996;20:445–452.
104. Iversen PO, Nicolaysen A, Kvernebo K, et al. Human cytokines modulate arterial vascular tone via endothelial receptors. *Pflugers Arch* 1999;439:93–100.
105. Libby P, Sukhova G, Lee RT, Galis ZS. Cytokines regulate vascular functions related to stability of the atherosclerotic plaque. *J Cardiovasc Pharmacol* 1995;25:S9–S12.
106. Schouten M, Wiersinga WJ, Levi M, van der Poll T. Inflammation, endothelium, and coagulation in sepsis. *J Leukoc Biol* 2008;83:536–545.
107. Mendall MA, Patel P, Ballam L, et al. C reactive protein and its relation to cardiovascular risk factors: A population based cross sectional study. *BMJ* 1996;312:1061–1065.
108. Bruunsgaard H, Skinhoj P, Pedersen AN, et al. Ageing, tumour necrosis factor- α (TNF- α) and atherosclerosis. *Clin Exp Immunol* 2000;121:255–260.
109. Lechleitner M, Koch T, Herold M, et al. Tumour necrosis factor- α plasma level in patients with type 1 diabetes mellitus and its association with glycaemic control and cardiovascular risk factors. *J Intern Med* 2000;248:67–76.
110. Tracy RP, Psaty BM, Macy E, et al. Lifetime smoking exposure affects the association of C-reactive protein with cardiovascular disease risk factors and subclinical disease in healthy elderly subjects. *Arterioscler Thromb Vasc Biol* 1997;17:2167–2176.
111. Spranger J, Kroke A, Mohlig M, et al. Inflammatory cytokines and the risk to develop type 2 diabetes: Results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)—Potsdam Study. *Diabetes* 2003;52:812–817.
112. Pradhan AD, Manson JE, Rifai N, et al. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 2001;286:327–334.
113. Cesari M, Penninx BW, Newman AB, et al. Inflammatory markers and cardiovascular disease (The Health, Aging and Body Composition [Health ABC] Study). *Am J Cardiol* 2003;92:522–528.
114. Gussekloo J, Schaap MC, Frolich M, et al. C-reactive protein is a strong but nonspecific risk factor of fatal stroke in elderly persons. *Arterioscler Thromb Vasc Biol* 2000;20:1047–1051.
115. Tracy RP, Lemaitre RN, Psaty BM, et al. Relationship of C-reactive protein to risk of cardiovascular disease in the elderly. Results from the Cardiovascular Health Study and the Rural Health Promotion Project. *Arterioscler Thromb Vasc Biol* 1997;17:1121–1127.
116. Vasan RS, Sullivan LM, Roubenoff R, et al. Inflammatory markers and risk of heart failure in elderly subjects without prior myocardial infarction: The Framingham Heart Study. *Circulation* 2003;107:1486–1491.
117. Rosano C, Marsland AL, Gianaros PJ. Maintaining brain health by monitoring inflammatory processes: A mechanism to promote successful aging. *Aging Dis* 2012;3:16–33.
118. Dunn AJ, Swiergiel AH. The role of cytokines in infection-related behavior. *Ann N Y Acad Sci* 1998;840:577–585.
119. Serrats J, Schiltz JC, Garcia-Bueno B, et al. Dual roles for perivascular macrophages in immune-to-brain signaling. *Neuron* 2010;65:94–106.
120. Hanisch UK. Microglia as a source and target of cytokines. *Glia* 2002;40:140–155.
121. Dik MG, Jonker C, Hack CE, et al. Serum inflammatory proteins and cognitive decline in older persons. *Neurology* 2005;64:1371–1377.
122. Yaffe K, Lindquist K, Penninx BW, et al. Inflammatory markers and cognition in well-functioning African-American and white elders. *Neurology* 2003;61:76–80.
123. Noble JM, Manly JJ, Schupf N, et al. Association of C-reactive protein with cognitive impairment. *Arch Neurol* 2010;67:87–92.
124. Schram MT, Euser SM, de Craen AJ, et al. Systemic markers of inflammation and cognitive decline in old age. *J Am Geriatr Soc* 2007;55:708–716.
125. Weaver JD, Huang MH, Albert M, et al. Interleukin-6 and risk of cognitive decline: MacArthur studies of successful aging. *Neurology* 2002;59:371–378.

126. Engelhart MJ, Geerlings MI, Meijer J, et al. Inflammatory proteins in plasma and the risk of dementia: The Rotterdam Study. *Arch Neurol* 2004;61:668–672.
127. Schmidt R, Schmidt H, Curb JD, et al. Early inflammation and dementia: A 25-year follow-up of the Honolulu-Asia Aging Study. *Ann Neurol* 2002;52:168–174.
128. Laurin D, David Curb J, Masaki KH, et al. Midlife C-reactive protein and risk of cognitive decline: A 31-year follow-up. *Neurobiol Aging* 2009;30:1724–1727.
129. Wilson CJ, Cohen HJ, Pieper CF. Cross-linked fibrin degradation products (D-dimer), plasma cytokines, and cognitive decline in community-dwelling elderly persons. *J Am Geriatr Soc* 2003;51:1374–1381.
130. Rafnsson SB, Deary IJ, Smith FB, et al. Cognitive decline and markers of inflammation and hemostasis: The Edinburgh Artery Study. *J Am Geriatr Soc* 2007;55:700–707.
131. Griffin WS. Neuroinflammatory cytokine signaling and Alzheimer's disease. *N Engl J Med* 2013;368:770–771.
132. Duong T, Nikolaeva M, Acton PJ. C-reactive protein-like immunoreactivity in the neurofibrillary tangles of Alzheimer's disease. *Brain Res* 1992;749:152–156.
133. Iwamoto N, Nishiyama E, Ohwada J, Arai H. Demonstration of CRP immunoreactivity in brains of Alzheimer's disease: Immunohistochemical study using formic acid pretreatment of tissue sections. *Neurosci Lett* 1994;177:23–26.
134. Tan ZS, Beiser AS, Vasan RS, et al. Inflammatory markers and the risk of Alzheimer disease: The Framingham Study. *Neurology* 2007;68:1902–1908.
135. Collins JS, Perry RT, Watson B Jr, et al. Association of a haplotype for tumor necrosis factor in siblings with late-onset Alzheimer disease: The NIMH Alzheimer Disease Genetics Initiative. *Am J Med Genet* 2000;96:823–830.
136. Licastro F, Pedrini S, Caputo L, et al. Increased plasma levels of interleukin-1, interleukin-6 and alpha-1-antichymotrypsin in patients with Alzheimer's disease: Peripheral inflammation or signals from the brain? *J Neuroimmunol* 2000;103:97–102.
137. Alvarez XA, Franco A, Fernandez-Novoa L, Cacabelos R. Blood levels of histamine, IL-1 beta, and TNF-alpha in patients with mild to moderate Alzheimer disease. *Mol Chem Neuropathol* 1996;29:237–252.
138. Huberman M, Sredni B, Stern L, et al. IL-2 and IL-6 secretion in dementia: Correlation with type and severity of disease. *J Neurol Sci* 1995;130:161–164.
139. Singh VK, Guthikonda P. Circulating cytokines in Alzheimer's disease. *J Psychiatr Res* 1997;31:657–660.
140. Fillit H, Ding WH, Buee L, et al. Elevated circulating tumor necrosis factor levels in Alzheimer's disease. *Neurosci Lett* 1991;129:318–320.
141. Bonaccorso S, Lin A, Song C, et al. Serotonin-immune interactions in elderly volunteers and in patients with Alzheimer's disease (DAT): Lower plasma tryptophan availability to the brain in the elderly and increased serum interleukin-6 in DAT. *Aging (Milano)* 1998;10:316–323.
142. Ravaglia G, Forti P, Maioli F, et al. Blood inflammatory markers and risk of dementia: The Conselice Study of Brain Aging. *Neurobiol Aging* 2007;28:1810–1820.
143. Lombardi VR, Garcia M, Rey L, Cacabelos R. Characterization of cytokine production, screening of lymphocyte subset patterns and in vitro apoptosis in healthy and Alzheimer's Disease (AD) individuals. *J Neuroimmunol* 1999;97:163–171.
144. van Duijn CM, Hofman A, Nagelkerken L. Serum levels of interleukin-6 are not elevated in patients with Alzheimer's disease. *Neurosci Lett* 1990;108:350–354.
145. Angelis P, Scharf S, Mander A, et al. Serum interleukin-6 and interleukin-6 soluble receptor in Alzheimer's disease. *Neurosci Lett* 1998;244:106–108.
146. Bauer J, Ganter U, Strauss S, et al. The participation of interleukin-6 in the pathogenesis of Alzheimer's disease. *Res Immunol* 1992;143:650–657.
147. Strauss S, Bauer J, Ganter U, et al. Detection of interleukin-6 and alpha 2-macroglobulin immunoreactivity in cortex and hippocampus of Alzheimer's disease patients. *Lab Invest* 1992;66:223–230.
148. Marsland AL, Gianaros PJ, Abramowitch SM, et al. Interleukin-6 covaries inversely with hippocampal grey matter volume in middle-aged adults. *Biol Psychiatry* 2008;64:484–490.
149. Holmes C, Cunningham C, Zotova E, et al. Systemic inflammation and disease progression in Alzheimer disease. *Neurology* 2009;73:768–774.
150. Gupta A, Watkins A, Thomas P, et al. Coagulation and inflammatory markers in Alzheimer's and vascular dementia. *Int J Clin Pract* 2005;59:52–57.
151. Comijs HC, Dik MG, Deeg DJ, Jonker C. The course of cognitive decline in older persons: Results from the Longitudinal Aging Study Amsterdam. *Dement Geriatr Cogn Disord* 2004;17:136–142.