

REVIEW



Chemotherapy-induced cognitive impairment: focus on the intersection of oxidative stress and TNF α

Nicole G. Rummel^{1,2} · Luksana Chaiswing^{2,3} · Subbarao Bondada^{2,4} · Daret K. St. Clair^{2,3} · D. Allan Butterfield^{1,2}

Received: 25 January 2021 / Revised: 27 June 2021 / Accepted: 17 August 2021 / Published online: 23 August 2021
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2021

Abstract

Chemotherapy-induced cognitive impairment (CICI) has been observed in a large fraction of cancer survivors. Although many of the chemotherapeutic drugs do not cross the blood–brain barrier, following treatment, the structure and function of the brain are altered and cognitive dysfunction occurs in a significant number of cancer survivors. The means by which CICI occurs is becoming better understood, but there still remain unsolved questions of the mechanisms involved. The hypotheses to explain CICI are numerous. More than 50% of FDA-approved cancer chemotherapy agents are associated with reactive oxygen species (ROS) that lead to oxidative stress and activate a myriad of pathways as well as inhibit pathways necessary for proper brain function. Oxidative stress triggers the activation of different proteins, one in particular is tumor necrosis factor alpha (TNF α). Following treatment with various chemotherapy agents, this pro-inflammatory cytokine binds to its receptors at the blood–brain barrier and translocates to the parenchyma via receptor-mediated endocytosis. Once in brain, TNF α initiates pathways that may eventually lead to neuronal death and ultimately cognitive impairment. TNF α activation of the c-jun N-terminal kinases (JNK) and Janus kinase-signal transducer and activator of transcription (JAK/STAT) pathways may contribute to both memory decline and loss of higher executive functions reported in patients after chemotherapy treatment. Chemotherapy also affects the brain's antioxidant capacity, allowing for accumulation of ROS. This review expands on these topics to provide insights into the possible mechanisms by which the intersection of oxidative stress and TNF α are involved in chemotherapy-induced cognitive impairment.

Keywords Chemotherapy induced cognitive impairment · Oxidative stress; · Pro-inflammatory cytokines · Neuronal death

Introduction

Chemotherapy

Chemotherapy-induced cognitive impairment (CICI), sometimes referred to by patients as “chemobrain” or “chemofog”, was first reported approximately 30 years ago [1].

While cognitive dysfunction might be expected in patients diagnosed with brain-resident tumors, CICI also occurs in patients without primary central nervous system (CNS) tumors [2]. Many patients diagnosed with various types of cancers reported lingering side effects following cessation of their chemotherapy treatments [1]. For example, breast cancer survivors often complained that cognitive impairment is the most troublesome post-treatment symptom, leading to a diminished quality of life characterized by decreased neuropsychological performance with effects mostly in working memory, executive functions, processing speeds, and cognitive processes, i.e., chemobrain or chemofog [3–7]. Chemotherapy has been associated with decreased neurogenesis and decreased cell proliferation along with release of inflammatory cytokines in the periphery [3].

Breast cancer patients were studied: pre-treatment, post-treatment, and 6 months post treatment and were found to have decreased cognitive reserve, increased anxiety, and depression from pre-treatment to 6 months post treatment

✉ D. Allan Butterfield
dabcns@uky.edu

¹ Department of Chemistry, University of Kentucky, Lexington, KY 40506, USA

² Markey Cancer Center, University of Kentucky, Lexington, KY 40536, USA

³ Department of Toxicology and Cancer Biology, University of Kentucky, Lexington, KY 40536, USA

⁴ Department of Microbiology, Immunology and Molecular Genetics, University of Kentucky, Lexington, KY 40536, USA

[6]. Cognitive deficits were described in some patients even prior to treatment with further decline after treatment, which can persist even up to 15 years post treatment [1]. Breast cancer patients with either stage 1, 2, or 3 cancer had lower cognitive performance compared to healthy or stage 0 cancer patients [7]. Different chemotherapy treatments may lead to varying levels of CICI [8].

Through the aid of preclinical animal studies, mechanisms for CICI have been proposed [3]. The effects of CICI were studied in cancer-naïve, female Sprague–Dawley rats showing significant decreased novel object recognition (NOR) performance and decreased [^{18}F]-fluorodeoxyglucose (^{18}F -FDG) uptake. This study suggested long-term impairment of the frontal cortex as a plausible contributor to the mechanism of CICI [9]. Mice were treated with doxorubicin (DOX), a chemotherapeutic agent that produces ROS via redox cycling of its quinone structure, leads to decreased cognitive performance assessed by NOR [10]. Another chemotherapy agent, methotrexate, was shown to lead to cognitive deficits through vascular toxicity and elevated peripheral and central TNF α [11]. The effects of 6 cytotoxic agents were tested on C57BL/6 J mice (cyclophosphamide, docetaxel, doxorubicin, 5-fluorouracil, methotrexate, and topotecan). Mice were sacrificed 3 or 15 weeks and the effects on neurogenesis, blood vessel density, and neuroinflammation were studied. No effect was seen on neurogenesis, increased blood vessel density in the hippocampus and prefrontal cortex, but a decrease in the number of microglial cells was observed. The decrease in microglial cells is consistent with inflammation after treatment [12].

An at-first mystifying aspect of CICI is that most chemotherapy agents do not cross the blood–brain barrier (BBB); however, these agents are associated with significant damage to the brain and CNS and can lead to CICI [13]. The number of possible causes of CICI is extensive, as the exact cause has not been definitively determined. Potential causes of CICI, among others, stem from the cancer itself and cancer treatment, altered brain biochemistry, genetic predisposition, possible tumor metastases to the brain, increased permeability of BBB due to decreased structural integrity, and DNA damage [2, 13]. Neuroinflammation is one of the leading hypotheses in the mechanisms underlying mental health problems, depression, and anxiety associated with CICI [14]. Complete surgical resection of a non-metastatic orthotopic, syngeneic mammary tumors in a rat model reversed the tumor-induced increased levels of circulating cytokines [14].

Adult neurogenesis occurs in niche regions within the brain; however, reduced neurogenesis is reported in chemotherapy-treated patients, increasing the risk of cognitive impairment [15]. Neurons are highly polarized and use spines and dendrites to regulate synaptic plasticity, which plays an important role in learning, memory, and executive

function. Loss of these spines and dendritic arborization is observed post treatment [15]. This process can lead to decreased neurotransmitter release, inflammation, and the breakdown of the BBB [16].

Chemotherapy often causes changes in brain structure

The effects on the brain due to chemotherapy treatments likely contribute to cognitive impairment [1]. Structural changes and inflammation in the brain following chemotherapy were observed by neuroimaging techniques, such as magnetic resonance imaging (MRI) as well as functional MRI (fMRI) [1]. Reductions in white and gray matter one year after treatment were observed as well as decreased activation in prefrontal cortex and hippocampus [1]. Reduced white matter integrity can be observed 10 months to 10 years post treatment [3]. In a study of cerebral white matter integrity, the chemotherapy group performed worse in attention, psychomotor speed, and memory [17]. White matter is necessary for communication between different brain regions with its integrity important for proper brain function.

Cancer survivors diagnosed with CICI tend to have an inverse correlation with intensity of the effects of CICI and the time after treatment, suggesting recovery from chemo-brain [15]. However, in contrast to this suggestion, other studies show diminution of cognition associated with chemotherapy can last at least a decade [15]. Decreased gray matter is seen in several regions immediately following chemotherapy treatments with partial recovery a year later [15]. Consistent with these findings, one month after chemotherapy completion, decreased gray matter is observed in breast cancer patients [18]. This aspect of CICI requires further investigation.


Recent studies suggest that chemotherapy itself can damage brain structure with a connection to cytokine dysregulation that impairs synaptic networks, hippocampal volume, and brain metabolism in the prefrontal and temporal cortex [11]. Chemotherapy-associated release of inflammatory cytokines in the periphery can lead to immune dysregulation. Induction of central inflammatory cytokines can cause activation of microglia [3] (Fig. 1).

Chemotherapy-induced oxidative stress results in the activation of the TNF α cascade

Cytokines and Inflammation

Microglia are considered the immune cells of the CNS and respond to most kinds of pathology [19]. When microglial

Fig. 1 Summary of cytokine-related mechanism of CICI discussed in this current review



cells are fully activated, they can be neurotoxic, following release of ROS, inducible nitric oxide synthase (iNOS) and TNF α . Cytokines play an important role in cell signaling by regulating immune and inflammatory responses [16]. Under normal conditions, inflammatory response and cytokine release are regulated [20]. Importantly, cytokines reportedly play a role in cancer progression and development [21]. Pro-inflammatory cytokine production has been suggested as a candidate mechanism for cognitive dysfunction [13, 22]. There are multiple conditions and stressors that may contribute to increased production of cytokines, which, among others, include cancer, acute and chronic stressors, physical and psychological stressors, and chemotherapy [21]. However, it is the increased complaints of cognitive dysfunction from patients following chemotherapy treatment that suggest these treatments are correlated to brain dysfunction [16].

Circulating cytokines can reach the brain via several pathways leading to stimulation of microglial cells [20, 23]. Communication between the peripheral cytokines and those in the CNS occurs through active transport across the BBB, passive transport through leaky regions of the BBB, receptor–ligand endocytosis, or stimulation through local inflammatory networks [21]. One cytokine, in particular, has been observed to play a major role in CICI: TNF α .

TNF α is a key player in pro-inflammatory cytokine cascades. TNF α is upstream of IL-1 and elicits the latter's production within the brain [24]. IL-1 β is pivotal for hippocampal learning in memory, however, high levels interfere with long-term potentiation and synaptic plasticity [24]. In mice following surgery, TNF α is detectable within the first 30 min and is the first cytokine to be released, while others are not detectable until 6 h post-surgery [24]. However, pre-operation administration of anti-TNF antibody reduced systemic IL-1 β . Trace fear conditioning also was performed in mice, and post-operation hippocampal-dependent memory impairment was observed. Pretreatment with anti-TNF antibody ameliorated cognitive decline [24].

In addition to in animal studies, plasma TNF α levels have been correlated with cognitive impairment in breast cancer patients. In two groups of patients, before chemotherapy (BCG) and after chemotherapy (ACG), plasma cytokine levels were evaluated (Am J TRANSL RES). The BCGs were further categorized into two groups, cognitive impaired

group (CIG) and cognitive normal group (CNG). Higher levels of IL-1 β , TNF α , and IL-4 were seen in ACG compared to BCG. The CIG also showed higher levels of the three cytokines compared to the CNG [25]. Two correlations were seen in the BCG, a direct correlation between cognition and quality of life and an inverse correlation between cognitive function and cytokine levels [25].

Post-chemotherapy patients reportedly demonstrated an increase in the soluble TNF receptor II (sTNF-RII). Higher baselines for the receptor were seen in patients with memory complaints, consistent with the notion that a post-chemotherapy increase in TNF α may play a role in the decline of cognition [26]. Blood cytokine levels of early-stage breast cancer patients were measured pre-surgery, post-surgery (prior to chemotherapy), and 6 months post-chemotherapy. At 6 months, the women reported a decline in quality of life and an increase in fatigue. An increase in sTNFRII, interleukin-6, and interleukin-10 was observed only in the post-chemotherapy group [27]. An increase in TNFRI and TNFRII is correlated to an increase in TNF α . The majority of biological effects of TNF α are mediated through its interaction with TNFRI [28]. Interleukin-10 is essential for exercise-induced analgesia; however, interleukin-10 has been suggested to resolve neuropathic pain induced by chemotherapeutics such as cisplatin [29]. Complete surgical resection of a non-metastatic orthotopic, syngeneic mammary tumor reversed the tumor-induced increases of circulating cytokines [14].

Activation of TNF α via ROS

Different chemotherapeutics induce damage to cancer cells via varying mechanisms. Alkylating agents such as cyclophosphamide cause DNA damage, antimetabolites, methotrexate and 5FU, disturb function and biosynthesis of DNA or RNA leading to cell death, and anti-microtubule agents, paclitaxel and docetaxel, interfere with cell division and proliferation [3]. In this review, we will focus on doxorubicin (DOX) for its cytotoxic effects through the production of ROS. DOX, used for the treatment of solid tumors and lymphoma, generates intracellular ROS evidenced by the observation that the intraperitoneal administration of DOX generates an increase in TNF α and oxidative damage in plasma and brain in mice [30]. The quinone structure

in DOX undergoes redox cycling between the quinone and semi-quinone radical, producing superoxide free radical in the presence of oxygen [31]. Patients treated with DOX had an increase in plasma oxidative stress and protein carbonylation as well as an increase in TNF α 6 h after treatment [30].

DOX has been shown to damage plasma proteins and increase TNF α levels in the periphery and in the brain [30, 31]. To gain insights into these observations, J774A.1 macrophage cultures were incubated with potassium superoxide, KO₂, which when dissolved in aqueous solution releases superoxide radicals; increased TNF α levels were observed, consistent with the notion that the superoxide free radical produced from DOX under redox cycling is connected to the increase in TNF α [32]. Lipid peroxidation in brain from mice occurs after injection intraperitoneally with DOX [33]. Although DOX does not cross the BBB, this chemotherapeutic agent induces a strong TNF α immunoreactivity in the cortex and hippocampus of DOX-treated mice. In addition, DOX increases pro-apoptotic factors, such as p53 and Bax, in brain mitochondria and increased release of cytochrome c and caspase-3 was noted following DOX treatment, consistent with apoptotic brain cell death [34].

To solidify this concept that TNF α is involved with oxidative damage in brain following treatment with BBB-impermeable DOX, mice with the gene for TNF α knocked out were investigated. TNF knockout mice had significant amelioration of oxidative stress in the brain as well as preservation of brain mitochondria function following DOX treatment [35]. In plasma, apolipoprotein A1 (ApoA1) inhibits the production of TNF α , however ROS in plasma causes oxidation of ApoA1 and promotes TNF α to interact with receptors on the BBB that lead to endocytic transfer to brain parenchyma [16]. DOX also activates microglia and astrocytes through TNF receptor 1 (TNFR1), which triggers iNOS release and therefore an increase in ROS and reactive nitrogen species (RNS) [36]. Supportive of the role of TNFA in CICI in the studies noted above, rats treated for four weeks with interferon- β -1a or Infliximab, both TNF α inhibitors, protected against DOX-induced chemo-brain [37].

Induction of the JAK-STAT pathway amplifies the signal from TNF α

Signal transducer and activator of transcription (STAT) mediates a vast array of processes that are required for homeostasis and development [38]. STAT activation induces cytokines and growth factors. Janus kinases (JAKs) phosphorylate and activate STAT monomers in the cytoplasm leading to homo- or heterodimerization [38]. JAK-STAT pathways relay signals from cellular membrane receptors to the nucleus and are involved in the pathways of cytokines, as the JAK-STAT enzymes are bound to the cytoplasmic

regions of cytokine receptors [39]. Oxidative stress activates these pathways evidenced by STAT1 and STAT3 activation in response to H₂O₂ [40].

TNF α and IL-6 were elevated in bone marrow aspirates of multiple myeloma patients, and the JAK-STAT pathway was directly involved in TNF α signaling pathways [41]. In murine samples, TNFA induced tyrosine phosphorylation and activation of JAKs, which was associated with tyrosine phosphorylation of STAT [42]. Activation of JAK-STAT pathways via TNF α leads to other cytokine activation. The majority of immune responses initiated by cytokines are dependent on STAT activation [38]. JAK-STAT induction of cytokine production along with TNF α initiation of other cytokines can lead to an amplification of effects of pro-inflammatory cytokines.

Activation of MAPK pathways follow TNF α activation of the JAK-STAT pathway

Mitogen-activated protein kinases (MAPK) are important signaling molecules and are classified in four groups: (1) extracellular signal-related kinases (ERK), (2) c-jun N-terminal (JNK) or stress-activated protein kinases (JNK/SAPK), (3) ERK5/big MAP kinase 1 (BMK1), and (4) the group of p38 MAPKs [43]. JNKs respond to cellular stresses, ERKs are important for signal transduction, and the p38 MAPK group is involved in inflammation, cellular growth, cell cycle, and cell death [43]. There are three isoforms of JNKs with JNK3 being expressed primarily in neuronal and cardiac cells [44]. These enzymes phosphorylate and regulate activity of transcription factors and are activated by MAPK modules. JNKs are pro-apoptotic kinases that play a role in apoptosis of neurons, especially TNF α -induced apoptosis as JNK is a downstream protein of the TNF α pathway [44]. The group of p38 MAPKs is activated by environmental stressors, growth factors, and MAP kinase kinase kinase (MAP3K) plays a role in the activation of both JNK and p38. Inflammation is a major activator of p38, which then plays a role in the production of cytokines, such as interleukin-1 β , interleukin-6, and TNF α [43]. ERKs are phosphorylated in response to mitogens and are involved in signaling for cell proliferation [45].

When DOX and cisplatin were administered together, activation of the JNK/p38 MAPK pathway and an increase in cytochrome c release from mitochondria were observed [46]. This observation is consistent with the notion that enhanced activation of JNK/p38 may lead to mitochondrial intrinsic apoptotic pathways [46]. As noted above, pro-inflammatory cytokines enter the BBB through receptor-mediated transport inducing the JNK/p38 MAPK pathway, which has been shown to be triggered by DOX [16]. Carmustine, an alkylating chemotherapeutic agent used to treat brain tumors, caused an accumulation of ROS via glutathione reductase

inhibition triggering a concentration- and time-dependent cell death of neuronally differentiated cells in association with increased caspase-3 activation and induction of phosphorylation of MAPKs, such as ERK, JNK, and p38. Inhibitors of these proteins prevented cell death via carmustine suggesting carmustine may induce cell death through these pathways [47].


DOX treatment led to increased mitochondrial apoptosis pathways evidenced by increased Bax to Bcl-2 ratios. MAPKs, ERK and p38 expressions also were altered after DOX administration [36]. Rats dosed with DOX led to increased pERK and p-p38 MAPK in sensory neurons and cortical neurons, blocking long-term synaptic facilitation (LTF). LTF was rescued in the presence of a p38 MAPK inhibitor, suggesting long-term memory impairment via the p38 MAPK pathway [48]. Ovariectomized rodents were given a drug cocktail of cyclophosphamide and DOX (AC-chemo), and the levels of pro-inflammatory cytokines and oxidative stress response gene markers were evaluated. ERK1/2 and JNK/SAPK signaling activities were elevated in the hippocampus [49] (Fig. 2).

Neuronal death in chemobrain is seen in a decrease in BDNF

Brain-derived neurotrophic factor (BDNF) plays a role in the survival and differentiation of neurons [50]. New neurons involved in memory and learning require BDNF to nurture the processes of adult neurogenesis. 5-fluorouracil (5-FU) disrupts cell proliferation and readily crosses the BBB. The effect of 5-FU on spatial working memory in mice was determined. 5-FU significantly reduced hippocampal BDNF and doublecortin protein levels associated with marginal disruption in spatial working memory. Reduced levels of BDNF and doublecortin in the hippocampus are indicative of alterations in neurotrophin levels and neurogenesis [51].

Exercise, which is reported to elevate brain levels of BDNF [52], has been known to positively affect brain function. Wistar rats, separated into four groups: control, control + exercise, chemobrain, and chemobrain + exercise were studied. Cognitive dysfunction was induced by DOX administration for four weeks. Exercise was completed via low-intensity treadmill for 6 days a week for the four weeks. The chemobrain group showed a decline in cognitive function, neurogenesis, and BDNF in the hippocampus. However, exercise attenuated these impairments in cognitive function and neuroplasticity for those with chemobrain [53].

Fig. 2 Anti-cancer treatments trigger ROS production and oxidative stress in the periphery inducing an increase in TNF α . TNF α interacts with its receptors in the bilayer activating the JAK/STAT pathway. Increased TNF α within the cytoplasm results in oxidative stress and impairment of antioxidants. Mitochondrial dysfunction leads to mPTP opening and Cyt. C release followed by neurodegeneration and CICI. JNK/ERK/p38 activation through TNF α also results in CICI.



Chemotherapy reduces ability for cellular defenses to combat effects of cognitive impairment

Although most chemotherapies do not cross the BBB, with some exceptions, they have been associated with cell death and decreased cell division as well as decreased antioxidant capabilities [13]. Glutathione (GSH), a major intracellular antioxidant found ubiquitously in the brain, under the action of glutathione peroxidase (GPx), scavenges H_2O_2 , which is converted to H_2O and O_2 via oxidation of GSH to glutathione disulfide (GSSG) [35]. GSSG is then converted back to GSH via glutathione reductase (GR). In DOX-treated mice, however, an increase in peripheral TNF α was observed [34, 35]. Decreased GSH levels in brain also were observed as well as decreased GSH/GSSG ratios [35], a marker of elevated oxidative stress upon Dox treatment. An increase in GSSG or a lack of activity of GR means that the powerful oxidative stress protection of GSH is lost after DOX treatment, which can result in oxidative stress-mediated neuronal dysfunction. Other antioxidants are also seen to be altered in brain after chemotherapy treatment. After intraperitoneal treatment of mice with DOX, a reduction in activity of the antioxidant enzyme manganese superoxide dismutase (MnSOD) in brain was observed, likely resulting from nitration of key Tyr residues in this protein and leading to more oxidative and nitrosative stress [54]. This is because MnSOD is a highly efficient enzyme for scavenging superoxide free radicals, particularly since this enzyme is mitochondrially resident, and mitochondria are a large source of superoxide free radical due to inefficiency of Complex I [31, 61].

Co-administration of protective agents with DOX may help to ameliorate the effects of DOX on the brain. For example, astaxanthin, a carotenoid antioxidant, anti-inflammatory and anti-apoptotic agent, reportedly protects against DOX-induced memory impairment. Administration of astaxanthin was able to restore hippocampal histopathological architecture and halted DOX-induced oxidative damage [55]. All three astaxanthin isomers were shown to reduce TNF α -induced secretion of interleukin 8 by 22–27% by inhibiting TNF-induced Nf-kB activation [56]. Another antioxidant, N-acetyl-cysteine (NAC) raises brain levels of GSH [57]. NAC is an approved FDA drug which provides the necessary cysteine, which is the rate-limiting substrate for the synthesis of GSH. NAC itself can act as an antioxidant scavenging for ROS as well as raise cysteine levels for GSH production [57]. In clinical studies, 40 children newly diagnosed with acute lymphoblastic leukemia were evaluated for the effects of NAC coupled with vitamin E. Twenty children took the combination vitamin E and NAC supplements, while the other 20 took a placebo. Glutathione peroxidase, malondialdehyde and TNF α levels were measured. Compared to the placebo group, a reduction in chemotherapy- and

radiotherapy-mediated toxicity was observed as indicated by an increase in glutathione peroxidase levels [58].

Sodium-2-mercapthoethane sulfonate (MESNA), due to its negative charge does not enter cancer cells and therefore does not interfere with cancer chemotherapy [59]. The -SH moiety on MESNA efficiently binds the lipid peroxidation product, HNE, preventing its toxic effects [20]. Treatment of mice with MESNA significantly modulated oxidative stress in brain and prevented cognitive loss assessed by NOR [8]. MESNA was shown to have poor cellular uptake by most cell types allowing it to act as an antioxidant in urine and blood and not interfere with the effects of the chemotherapeutic drug [59]. Co-administration of MESNA with DOX reduced post-treatment levels of TNF-related cytokines and TNFR1/2 [59]. Patients not treated with MESNA showed a statistically significant increase in proteins adducted by 4-hydroxynonenal (HNE) compared to those treated with MESNA [30]. Mulmina, a formulation of natural compounds known to help improve function and prevent cognitive decline, was given to normal rats also injected with a combination of cyclophosphamide, methotrexate, and 5-FU (CMF). Those injected with mulmina showed significant improvements in cognitive function post chemotherapy [60].

Antioxidant therapy for cancer patients has to be carefully considered. For example, elevation of brain levels of GSH by i.p. injection in mice of γ -glutamylcysteine ethyl ester (GCEE) led to protection against elevated oxidative stress associated with DOX treatment, indicating that free radicals are involved in brain effects induced by DOX-induced, TNF α -mediated oxidative damage in brain [33]. However, cancer cell-permeable antioxidants have the risk of lowering the redox state of cancer cells and allowing GSH-mediated, glutathione-S-transferase-facilitated binding to chemotherapeutic agents. Such GSH-chemotherapy agent complexes are excellent substrates for the multi-drug resistance protein-1 (MRP-1), which effluxes these complexes out of the cancer cell. That is, the therapeutic efficacy of chemotherapy agents is potentially drastically reduced with such agents [33]. In contrast, MESNA mentioned above, does not enter cancer cells and thereby eliminates these concerns. Moreover, selected redox-sensitive antioxidants have the property of raising the already higher redox state of cancer cells to an even higher state, exceeding the apoptosis threshold, leading to cancer cell death without affecting non-cancer cells appreciably [61].

Conclusion

CICI is often observed in cancer patients without CNS tumors, but the underlying mechanisms by which this quality-of-life diminution is caused have not been definitively

elucidated. Numerous hypotheses have been formed for the cause to CICI, with one being the production of ROS from chemotherapy treatments leading to the activation of pro-inflammatory cytokines such as TNF α that cross the BBB and lead to neuronal death. Although many patients seem to recover slowly over time after chemotherapy treatments end, understanding the mechanisms by which CICI occurs and how to combat this impairment will allow for better quality of life outcomes for the ever-growing number of cancer survivors as a result of improvements in cancer treatment.

Acknowledgements This work was supported in part by a multiple PI R01 NIH grant [CA217934] to D.K.S.C., S.B., and D.A.B.

Authors' contributions NGR and DA Butterfield wrote the article, while LC, SB, and DC each edited the article and offered suggestions for clarity of expression. All authors agreed to publish this manuscript.

Funding This work was supported in part by a multiple PI R01 NIH Grant [CA217934] to D.K.S.C., S.B., and D.A.B.

Availability of data and material Not applicable for this review paper.

Code availability Not applicable for this review paper.

Declarations

Conflicts of interest/Competing interests The authors declare that they have no conflict of interest.

Ethics approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

References

- Orchard TS et al (2017) Clearing the fog: a review of the effects of dietary omega-3 fatty acids and added sugars on chemotherapy-induced cognitive deficits. *Breast Cancer Res Treat* 161(3):391–398
- Jean-Pierre P, Johnson-Greene D, Burish TG (2014) Neuropsychological care and rehabilitation of cancer patients with chemobrain: strategies for evaluation and intervention development. *Support Care Cancer* 22(8):2251–2260
- Mounier NM et al (2020) Chemotherapy-induced cognitive impairment (CICI): An overview of etiology and pathogenesis. *Life Sci* 258:118071
- Boykoff N, Moieni M, Subramanian SK (2009) Confronting chemobrain: an in-depth look at survivors' reports of impact on work, social networks, and health care response. *J Cancer Surviv* 3(4):223–232
- McDonald BC et al (2010) Gray matter reduction associated with systemic chemotherapy for breast cancer: a prospective MRI study. *Breast Cancer Res Treat* 123(3):819–828
- Janelins MC et al (2017) Cognitive complaints in survivors of breast cancer after chemotherapy compared with age-matched controls: an analysis from a nationwide, multicenter Prospective Longitudinal Study. *J Clin Oncol* 35(5):506–514
- Ahles TA et al (2008) Cognitive function in breast cancer patients prior to adjuvant treatment. *Breast Cancer Res Treat* 110(1):143–152
- Cheung YT et al (2013) Cytokines as mediators of chemotherapy-associated cognitive changes: current evidence, limitations and directions for future research. *PLoS One* 8(12):e81234
- Barry RL et al (2018) In vivo neuroimaging and behavioral correlates in a rat model of chemotherapy-induced cognitive dysfunction. *Brain Imaging Behav* 12(1):87–95
- Keeney JTR et al (2018) Doxorubicin-induced elevated oxidative stress and neurochemical alterations in brain and cognitive decline: protection by MESNA and insights into mechanisms of chemotherapy-induced cognitive impairment ("chemobrain"). *Oncotarget* 9(54):30324–30339
- Das A et al (2020) An overview on chemotherapy-induced cognitive impairment and potential role of antidepressants. *Curr Neuropharmacol* 18(9):838–851
- Seigers R et al (2016) Neurobiological changes by cytotoxic agents in mice. *Behav Brain Res* 299:19–26
- Ahles TA, Saykin AJ (2007) Candidate mechanisms for chemotherapy-induced cognitive changes. *Nat Rev Cancer* 7(3):192–201
- Pyter LM et al (2017) Novel rodent model of breast cancer survival with persistent anxiety-like behavior and inflammation. *Behav Brain Res* 330:108–117
- Nguyen LD, Ehrlich BE (2020) Cellular mechanisms and treatments for chemobrain: insight from aging and neurodegenerative diseases. *EMBO Mol Med* 12(6):12075
- Ren X, St Clair DK, Butterfield DA (2017) Dysregulation of cytokine mediated chemotherapy induced cognitive impairment. *Pharmacol Res* 117:267–273
- Deprez S et al (2012) Longitudinal assessment of chemotherapy-induced structural changes in cerebral white matter and its correlation with impaired cognitive functioning. *J Clin Oncol* 30(3):274–281
- McDonald BC et al (2013) Frontal gray matter reduction after breast cancer chemotherapy and association with executive symptoms: a replication and extension study. *Brain Behav Immun* 30(Suppl):S117–S125
- Hanisch UK, Kettenmann H (2007) Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 10(11):1387–1394
- Seruga B et al (2008) Cytokines and their relationship to the symptoms and outcome of cancer. *Nat Rev Cancer* 8(11):887–899
- Wang XM et al (2015) Chemobrain: a critical review and causal hypothesis of link between cytokines and epigenetic reprogramming associated with chemotherapy. *Cytokine* 72(1):86–96
- Wardill HR et al (2016) Cytokine-mediated blood brain barrier disruption as a conduit for cancer/chemotherapy-associated neurotoxicity and cognitive dysfunction. *Int J Cancer* 139(12):2635–2645
- Schrepf A, Lutgendorf SK, Pyter LM (2015) Pre-treatment effects of peripheral tumors on brain and behavior: neuroinflammatory mechanisms in humans and rodents. *Brain Behav Immun* 49:1–17
- Terrando N et al (2010) Tumor necrosis factor- α triggers a cytokine cascade yielding postoperative cognitive decline. *Proc Natl Acad Sci USA* 107(47):20518–20522
- Zhao J et al (2020) Changes in plasma IL-1 β , TNF- α and IL-4 levels are involved in chemotherapy-related cognitive impairment in early-stage breast cancer patients. *Am J Transl Res* 12(6):3046–3056
- Ganz PA et al (2013) Does tumor necrosis factor- α (TNF- α) play a role in post-chemotherapy cerebral dysfunction? *Brain Behav Immun* 30(Suppl):S99–108

27. Jenkins V et al (2016) A feasibility study exploring the role of pre-operative assessment when examining the mechanism of “chemobrain” in breast cancer patients. *Springerplus* 5:390
28. Tian T, Wang M, Ma D (2014) TNF- α , a good or bad factor in hematological diseases? *Stem Cell Investig* 1:12
29. Laumet G et al (2020) Interleukin-10 resolves pain hypersensitivity induced by cisplatin by reversing sensory neuron hyperexcitability. *Pain* 161(10):2344–2352
30. Aluise CD et al (2011) 2-Mercaptoethane sulfonate prevents doxorubicin-induced plasma protein oxidation and TNF- α release: implications for the reactive oxygen species-mediated mechanisms of chemobrain. *Free Radic Biol Med* 50(11):1630–1638
31. Butterfield DA (2014) The 2013 SFRBM discovery award: selected discoveries from the butterfield laboratory of oxidative stress and its sequela in brain in cognitive disorders exemplified by Alzheimer disease and chemotherapy induced cognitive impairment. *Free Radic Biol Med* 74:157–174
32. Keeney JT et al (2015) Superoxide induces protein oxidation in plasma and TNF- α elevation in macrophage culture: Insights into mechanisms of neurotoxicity following doxorubicin chemotherapy. *Cancer Lett* 367(2):157–161
33. Joshi G et al (2007) Glutathione elevation by gamma-glutamyl cysteine ethyl ester as a potential therapeutic strategy for preventing oxidative stress in brain mediated by in vivo administration of adriamycin: Implication for chemobrain. *J Neurosci Res* 85(3):497–503
34. Tangpong J et al (2006) Adriamycin-induced, TNF- α -mediated central nervous system toxicity. *Neurobiol Dis* 23(1):127–139
35. Joshi G et al (2010) Alterations in brain antioxidant enzymes and redox proteomic identification of oxidized brain proteins induced by the anti-cancer drug adriamycin: implications for oxidative stress-mediated chemobrain. *Neuroscience* 166(3):796–807
36. Ongnok B, Chattipakorn N, Chattipakorn SC (2020) Doxorubicin and cisplatin induced cognitive impairment: the possible mechanisms and interventions. *Exp Neurol* 324:113118
37. Wahdan SA et al (2020) Abrogating doxorubicin-induced chemobrain by immunomodulators IFN- β 1a or infliximab: Insights to neuroimmune mechanistic hallmarks. *Neurochem Int* 138:104777
38. Owen KL, Brockwell NK, Parker BS (2019) JAK-STAT Signaling: A Double-Edged Sword of Immune Regulation and Cancer Progression. *Cancers (Basel)* 11(12):2002
39. Seif F et al (2017) The role of JAK-STAT signaling pathway and its regulators in the fate of T helper cells. *Cell Commun Signal* 15(1):23
40. Simon AR et al (1998) Activation of the JAK-STAT pathway by reactive oxygen species. *Am J Physiol* 275(6):C1640–C1652
41. Lee C et al (2013) TNF α mediated IL-6 secretion is regulated by JAK/STAT pathway but not by MEK phosphorylation and AKT phosphorylation in U266 multiple myeloma cells. *Biomed Res Int* 2013:580135
42. Guo D et al (1998) Induction of Jak/STAT signaling by activation of the type 1 TNF receptor. *J Immunol* 160(6):2742–2750
43. Ono K, Han J (2000) The p38 signal transduction pathway: activation and function. *Cell Signal* 12(1):1–13
44. Liu J, Lin A (2005) Role of JNK activation in apoptosis: a double-edged sword. *Cell Res* 15(1):36–42
45. Busca R, Pouyssegur J, Lenormand P (2016) ERK1 and ERK2 Map kinases: specific roles or functional redundancy? *Front Cell Dev Biol* 4:53
46. Ohtsuka T et al (2003) Synergistic induction of tumor cell apoptosis by death receptor antibody and chemotherapy agent through JNK/p38 and mitochondrial death pathway. *Oncogene* 22(13):2034–2044
47. An JM et al (2011) Carmustine induces ERK- and JNK-dependent cell death of neuronally-differentiated PC12 cells via generation of reactive oxygen species. *Toxicol In Vitro* 25(7):1359–1365
48. Liu RY et al (2014) Doxorubicin attenuates serotonin-induced long-term synaptic facilitation by phosphorylation of p38 mitogen-activated protein kinase. *J Neurosci* 34(40):13289–13300
49. Bagnall-Moreau C et al (2019) Chemotherapy-induced cognitive impairment is associated with increased inflammation and oxidative damage in the hippocampus. *Mol Neurobiol* 56(10):7159–7172
50. Miranda M et al (2019) Brain-derived neurotrophic factor: a key molecule for memory in the healthy and the pathological brain. *Front Cell Neurosci* 13:363
51. Mustafa S et al (2008) 5-Fluorouracil chemotherapy affects spatial working memory and newborn neurons in the adult rat hippocampus. *Eur J Neurosci* 28(2):323–330
52. Cotman CW, Berchtold NC (2002) Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci* 25(6):295–301
53. Park HS et al (2018) Physical exercise prevents cognitive impairment by enhancing hippocampal neuroplasticity and mitochondrial function in doxorubicin-induced chemobrain. *Neuropharmacology* 133:451–461
54. Tangpong J et al (2007) Adriamycin-mediated nitration of manganese superoxide dismutase in the central nervous system: insight into the mechanism of chemobrain. *J Neurochem* 100(1):191–201
55. El-Agamy SE et al (2018) Astaxanthin ameliorates doxorubicin-induced cognitive impairment (chemobrain) in experimental rat model: impact on oxidative, inflammatory, and apoptotic machineries. *Mol Neurobiol* 55(7):5727–5740
56. Yang C et al (2019) Anti-inflammatory effects of different astaxanthin isomers and the roles of lipid transporters in the cellular transport of astaxanthin isomers in Caco-2 cell monolayers. *J Agric Food Chem* 67(22):6222–6231
57. Huang Q et al (2010) Potential in vivo amelioration by N-acetyl-L-cysteine of oxidative stress in brain in human double mutant APP/PS-1 knock-in mice: toward therapeutic modulation of mild cognitive impairment. *J Neurosci Res* 88(12):2618–2629
58. Al-Tonbary Y et al (2009) Vitamin e and N-acetylcysteine as antioxidant adjuvant therapy in children with acute lymphoblastic leukemia. *Adv Hematol* 2009:689639
59. Hayslip J et al (2015) Plasma TNF- α and soluble TNF receptor levels after doxorubicin with or without co-administration of Mesna-A randomized Cross-Over Clinical Study. *PLoS ONE* 10(4):e0124988
60. Kinra M et al (2021) Neuroprotective effect of Mulmina against chemotherapy-induced cognitive decline in normal rats. *Biomed Rep* 14(1):1
61. Miriyala S et al (2012) Manganese superoxide dismutase, MnSOD and its mimics. *Biochim Biophys Acta* 1822(5):794–814
62. Figures Created with Biorender.com

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.