

MASTER EQUATION CODEX

Volume II

The Emergence of Life

Chemistry · Biochemistry · Cell Biology · Genetics · Physiology

Plants · Fungi · Insects · Fish · Reptiles · Birds · Mammals

Pharmacology · Ecology · Evolution

Keith I. Blaze · DTC / Rosetta Stone Protocol Research Programme · June 2026

Volume I established the chain from primitive distinction to spacetime, matter, and forces. Volume II extends the chain downward into complexity: from atomic bonds to molecules, from molecules to self-replicating chemistry, from replication to the cell, from the cell to the organism, from the organism to the ecosystem. Every level is a persistence problem — the same formal question asked at a new scale. The equations are the grammar. The living world is the text.

The spectral cascade of Vol. I ends with atoms: nuclei bound by the strong force, electrons occupying orbitals defined by the Schrödinger equation. Chemistry begins where quantum mechanics hands off — at the scale where electrons are shared between nuclei to form bonds.

0.1

$$\hat{H}\Psi = E\Psi$$

Atomic orbital: electron wavefunction in nuclear potential

0.2

$$E^n = -m_e e^4 / (8\epsilon_0^2 h^2 n^2)$$

Bohr energy levels: discrete electron shells

0.3

$$\Psi_a^b = c_1\phi_1 + c_2\phi_2$$

Molecular orbital (LCAO): bond from wavefunction overlap

Spec(L) → m_n, α_k → atomic orbitals ϕ_i [spectral → chemistry handoff]

Chemical bonds are persistent constraint structures — atoms minimise free energy by sharing electrons. Reaction dynamics are governed by thermodynamic potentials and kinetic barriers. The same persistence logic from Vol. I operates here at the molecular scale.

1.1

$$\Delta G = \Delta H - T\Delta S$$

Gibbs free energy: thermodynamic driver of all chemical change

1.2

$$\Delta G^\circ = -RT \ln K_{eq}$$

Equilibrium constant: bond stability as free-energy minimum

1.3

$$k = Ae^{-E_a/RT}$$

Arrhenius equation: reaction rate over energy barrier E_a

1.4

$$d[A]/dt = -k[A]$$

First-order kinetics: concentration decay rate

1.5

$$\text{pH} = -\log[H^+]$$

Acid-base balance: proton activity governs molecular ionisation

► Persistence criterion: a bond persists if $\Delta G < 0$. Every covalent, ionic, or hydrogen bond is a local free-energy minimum — a spectral mode below the dissociation threshold $\lambda \mathfrak{Z}$.

2 BIOCHEMISTRY — THE MOLECULES OF LIFE

Biochemistry

Four molecular classes — nucleic acids, proteins, lipids, carbohydrates — are the persistent distinction structures of the living world. Their behaviour is governed by the same thermodynamic and kinetic laws, now applied to polymer chains and catalytic surfaces.

2.1

$$v = V_{\max}[S] / (K^M + [S])$$


Michaelis-Menten: enzyme catalysis rate as a function of substrate

2.2

$$\Delta G = \Delta G^\circ + RT \ln([P]/[R])$$

Reaction free energy: drives metabolic flux through pathways

2.3



ATP hydrolysis: universal energy currency of the cell

2.4

$$K^D = [RL] / ([R][L])$$

Binding affinity: receptor-ligand dissociation constant

ATP hydrolysis → protein conformation change [free energy transduction]

► Enzymes are persistence amplifiers: they lower E_a without changing ΔG , dramatically accelerating the rate at which the system finds its free-energy minimum. Enzyme specificity is encoded in protein structure, which is encoded in the genome.

3 GENETICS — THE CODE OF PERSISTENCE

Molecular Biology / Genetics

The central dogma is the information chain of life: DNA is copied faithfully, transcribed into RNA, and translated into protein. This is the cell's mechanism for propagating its distinction structure across time — the biological realisation of the persistence chain $\Delta \rightarrow \tau \rightarrow \kappa \rightarrow \Pi$.

3.1

DNA → mRNA → Protein

Central dogma: information flow from genome to function

3.2

$$dN/dt = \mu N(1 - N/K)$$

Logistic replication: population growth with carrying capacity K

3.3	$P(\text{mutation}) = \mu \text{ per base per replication}$	Mutation rate: stochastic variation in the information copy
3.4	$H(X) = -\sum p_i \log_2 p_i$	Sequence entropy: information content of the genetic code
3.5	$F_{ij} = \langle \psi_i \hat{H} \psi_j \rangle$	Overlap integral: molecular recognition energy (gene expression)

► A genome is a compressed description of a persistence-maximising organism. The fidelity of DNA polymerase (error rate $\approx 10^{-9}$) is the physical mechanism of constraint κ in the biological domain.

4	THE CELL — MINIMAL PERSISTENT SYSTEM	Cell Biology
---	--------------------------------------	--------------

The cell is the minimal unit of life: a lipid membrane that enforces organisational closure (Φ), an energy budget that must stay solvent ($dE/dt \geq 0$), and a genome that replicates the whole structure. Every multicellular organism, every tissue, every organ, is an elaboration of this one level.

4.1	$\Phi = \{f_i : \Sigma f_i \rightarrow f_i\}$	Organisational closure: each cellular process produces the next
4.2	$dE/dt = G^{cee} - W^c \eta^c$	Energy balance: ATP generation minus maintenance cost
4.3	$V^m = dV/dt = k_p^{oo} - k^{Ded}$	Cell volume regulation: osmotic balance
4.4	$J = -D \nabla c$	Fick's first law: molecular diffusion across membranes
4.5	$V^M = (RT/zF) \ln([X]_o/[X]_i)$	Nernst equation: ion equilibrium potential across membrane

Genome → Proteins → Organelles → Φ [gene expression → closure]

► Prokaryotic cells (≈ 13.5 billion years old) solved the minimal persistence problem. Eukaryotic cells added membrane-bound organelles, massively expanding the constraint architecture — enabling all multicellular life.

Natural selection is the universe's algorithm for solving the persistence maximisation problem at the population level. Heritable variation plus differential reproduction equals adaptive evolution. The fitness function is the biological proxy for $A^* = \text{argmax}(\Pi/S)$.

5.1

$$\bar{\Delta z} = h^2 S$$

Breeder's equation: response to selection = heritability × differential

5.2

$$\bar{w} = \sum p_i w_i$$

Mean fitness: population-averaged persistence rate

5.3

$$\Delta p = p(1-p)(w_1 - w_s) / \bar{w}$$

Hardy-Weinberg change: allele frequency shift under selection

5.4

$$dS/dt \propto \text{variance}(w)$$

Fisher's fundamental theorem: fitness increases at rate of its variance

5.5

$$K = T_{tt}^d / T_{r_i^n}^{d \circ L \circ}$$

Substitution rate: neutral vs. selected mutation accumulation

mutation μ + selection w → adapted phenotype [evolutionary time]

► Evolution is persistence under noise. Mutation generates new distinctions; selection retains those that maximise Π/S in the current environment. Genetic drift (random walk in allele space) operates when selection coefficients are smaller than $1/2N_e$.

Plants solved the persistence problem by tapping the largest available free-energy source: electromagnetic radiation from the Sun. Photosynthesis couples light energy to carbon fixation, building structural biomass from atmospheric CO_2 .

6.1



Net photosynthesis: light energy stored as glucose

6.2

$$A = g_s(C_a - C_i)$$

Fick diffusion of CO_2 : stomatal conductance controls fixation rate

6.3	$A = V_{\text{max}} (C_i - \Gamma^*) / (K_m + C_i)$	Farquhar-von Caemmerer-Berry: RuBisCO-limited photosynthesis
6.4	$GP^P = NPP + R_a$	Gross primary production = net production + autotrophic respiration
6.5	$dB/dt = \mu B(1 - B/K) - mB$	Plant biomass dynamics: growth vs. mortality

► Vascular plants (tracheophytes, ≈470 Mya) added xylem and phloem — the first long-range transport network in a multicellular organism — enabling persistence at the scale of forests.

7	FUNGI — CONSTRAINT DISSOLUTION AND MYCORRHIZAL NETWORKS	<i>Mycology</i>
---	--	-----------------

Fungi are heterotrophs that dissolve organic constraint structures — breaking down dead biomass and cycling nutrients. Their hyphal networks are the internet of the forest floor, redistributing carbon and minerals across kilometres.

7.1	$dC/dt = k^d C$	Decomposition rate: first-order carbon mineralisation
7.2	$\Phi_{\text{myc}} = \alpha N^{\text{np}} / (K + C_p)$	Mycorrhizal flux: nutrient transfer to plant root per hyphal length
7.3	$J^c = -D_e^{\text{oo}} \nabla C^c$	Effective diffusivity: nutrient flow through hyphal network

Dead biomass → Mineral nutrients (N, P, K) [fungal decomposition]

► Mycorrhizal symbiosis (≈90% of plant species) is a constraint-sharing arrangement: plant provides photosynthate (glucose), fungus provides phosphorus and water. Both parties persist longer by coupling their Π/S ratios.

Animals solved persistence differently: instead of waiting for resources to diffuse, they evolved motility and nervous systems to actively seek, evaluate, and acquire them. The action potential is the animal's fundamental computation unit.

8.1

$$C^M dV/dt = -\Sigma I_i + I_a^{pp}$$

Hodgkin-Huxley membrane equation: voltage across neural membrane

8.2

$$I^{ia} = g^{ia} m^3 h (V^M - E^{ia})$$

Na⁺ current: voltage-gated channel activation drives action potential

8.3

$$I^K = g^K n^4 (V^M - E^K)$$

K⁺ current: repolarisation restores resting state

8.4

$$V(t) = V_s^{Re} + (V_0 - V_s^{Re})e^{-t/\tau}$$

Passive membrane: exponential approach to rest potential

Sensory input Δ_n → action potential spike *[neural encoding of distinction]*

► The nervous system is a real-time distinction-processing network. Sensory organs transduce physical signals into spike trains; motor neurons convert computed decisions into muscular force. The brain is a Laplacian operator on the graph of sensory-motor experience.

Fish were the first vertebrates, appearing ≈530 Mya. They solved persistence in aquatic environments by mastering buoyancy, hydrodynamic drag reduction, and osmoregulation — maintaining internal ionic homeostasis against the concentration gradients of ocean or freshwater.

9.1

$$F^D = \frac{1}{2} C^D \rho A v^2$$

Drag force: hydrodynamic resistance scaled by velocity squared

9.2

$$B = \rho^{Lood} V g$$

Buoyancy: displaced fluid weight sustains neutral position

9.3

$$J_i^n = P_n^{Lo} \alpha ([X]_o - [X]_i)$$

Ion flux: gill epithelium maintains osmotic homeostasis

► The swim bladder (gas-filled sac) is a biological solution to the free-energy cost of vertical positioning: by adjusting gas volume, fish maintain neutral buoyancy at any depth with near-zero metabolic expenditure — a spectral optimisation.

10	REPTILES — ECTOTHERMY AND ENVIRONMENTAL COUPLING	Herpetology / Thermal Biology
----	---	-------------------------------

Reptiles are ectotherms: their body temperature tracks the environment. This strategy minimises metabolic overhead — the energetic constraint $W\propto\eta$ is reduced by outsourcing thermoregulation to solar radiation, allowing far greater body mass per unit food consumed.

10.1	$T^b = T_a + Q_s^{OM}/h_a$	Operative temperature: body temp from solar load and convective loss
10.2	$T_s^{me} = T^b + Q_s^{me}/h_s^p$	Metabolic temperature correction: endogenous heat in large reptiles
10.3	$SMR = aM^b$	Standard metabolic rate: allometric scaling with body mass M

Solar radiation $h\nu \rightarrow T^b \rightarrow$ enzyme activity $v(T)$ [environmental free energy $\rightarrow$ biochemistry]

► The amniotic egg (appearing ≈ 312 Mya) was the key constraint innovation of reptiles: it internalised the aquatic environment required for embryo development, decoupling vertebrate reproduction from standing water.

11	BIRDS — ENDOTHERMY, FLIGHT, AND CONSTRAINT MASTERY	Ornithology / Aerodynamics
----	---	----------------------------

Birds solved the persistence problem at the cost of high metabolic overhead: endothermy (self-heating) plus powered flight. The payoff is access to resources across vast spatial scales and independence from ambient temperature — a high-cost, high-persistence strategy.

11.1	$L = \frac{1}{2}C_D\rho Av^2$	Aerodynamic lift: wing area and velocity sustain weight
11.2	$p^{ReO} = mgv^{ReO}$	Minimum power speed: optimal cruising flight efficiency

11.3	$BMR \propto M^{0.73}$	Basal metabolic rate: Kleiber's law — scales as M to the $\frac{3}{4}$ power
------	------------------------	--

11.4	$T^b = T_a + Q_n^d / (h A_s)$	Fourier heat loss: insulation (feathers) limits thermal flux
------	-------------------------------	--

► Kleiber's law ($BMR \propto M^{0.75}$) is one of ecology's deepest empirical regularities, holding across 18 orders of magnitude of body mass. Its origin lies in the fractal geometry of vascular networks — a spectral argument in disguise.

12	MAMMALS — HOMEOSTASIS, MILK, AND MIND	Mammalogy / Physiology
----	--	------------------------

Mammals pushed endothermy and parental investment to their logical extreme: live birth, lactation, and extended learning periods. The neocortex is the mammalian solution to navigating high-dimensional environments — a distinction graph instantiated in neural tissue.

12.1	$dT^b/dt = Q_n^d - Q_{oss} / (mC^p)$	Thermoregulation: core body temperature dynamics
------	--------------------------------------	--

12.2	$C = \beta R_i^\infty (1 - e^{-K/\tau_i})$	Cardiac output: volume per beat $\times$ heart rate
------	--	---

12.3	$HRV \propto 1/f \text{ noise}$	Heart rate variability: fractal autonomic regulation
------	---------------------------------	--

12.4	$H = \text{ArgMax}[\Pi(\Delta, \tau, \kappa, L, R, V)]$	Human observer: self-referential persistence maximiser
------	---	--

► The human brain (Vol. I, Eq. 10.4) is the mammalian neocortex taken to its current recursion depth: a distinction graph that models itself, models others, and models models. Language is the compression algorithm; culture is the distributed constraint network.

13	INSECTS — SWARMS, EXOSKELETONS, AND METAMORPHOSIS	Entomology
----	--	------------

Insects are the most species-rich animal class on Earth (≈ 10 million species, $>75\%$ of all animal species). Their success rests on three constraint innovations: the exoskeleton (rigid external constraint structure), complete metamorphosis (constraint restructuring across life stages), and eusocial colonies (distributed persistence across individuals).

13.1	$\sigma_p^e = F/A$	Exoskeletal stress: load-bearing capacity of chitin cuticle
------	--------------------	---

13.2	$dN/dt = rN - \alpha NP$	Lotka-Volterra predator term: insect population under predation
13.3	$p_{\text{in}}^{\text{colony}} = e^{-K_c/k_B T}$	Pupation probability: metamorphosis threshold from thermal accumulation
13.4	$C_n^{\text{colony}} = \gamma N^{\alpha} e^{-\delta d}$	Pheromone concentration: colony signalling decays with distance d

► Eusocial insect colonies (ants, bees, termites) are superorganisms: the colony is the persistence unit, not the individual. Queen reproductive monopoly + worker specialisation = maximal division of labour under constraint.

14	PHARMACOLOGY — DIRECTED CONSTRAINT MODIFICATION	Pharmacology / Medicine
----	--	-------------------------

Pharmacology is the science of targeted molecular intervention: drugs modify the constraint landscape of biological systems by binding to receptors, enzymes, or ion channels, shifting equilibria and changing the rate of specific reactions. Every drug is a designed distinction.

14.1	$E = E_{\text{max}} [D]^n / (ED_{50}^n + [D]^n)$	Hill equation: dose-response curve with cooperativity n
14.2	$dC/dt = -k_{\text{el}} C + F \cdot D/V$	PK equation: drug concentration (elimination – absorption)
14.3	$t_{1/2} = 0.693/k_{\text{el}}$	Half-life: time for drug concentration to halve
14.4	$K^D = k_{\text{off}} / k_{\text{on}}$	Binding affinity: thermodynamic measure of drug-target specificity
14.5	$LD_{50} / ED_{50} = \text{Therapeutic Index}$	Safety ratio: toxic vs. effective dose separation

$$K^D (\text{drug-receptor}) \rightarrow \Delta G_{\text{in}}^{\text{d}} = -RT \ln K^D \quad [\text{binding thermodynamics}]$$

► A drug succeeds when it shifts the system's persistence landscape: reducing the energy barrier for a beneficial process, or raising it for a pathological one. Drug discovery is constraint engineering at the molecular scale.

An ecosystem is a network of energy and matter flows in which every species is both a constraint on and a constraint for every other species. Stability is not a static property but a dynamic balance between disturbance and recovery — the biological analogue of $dE/dt = G - W\Xi\eta$.


15.1	$dN_i/dt = r_i N_i (1 - \sum_j \alpha_{ij} N_j / K_i)$	Lotka-Volterra competition: multi-species population dynamics
15.2	$E = \text{NPP} / R_a^o p$	Ecological efficiency: energy transfer between trophic levels
15.3	$S = cA_r$	Species-area relationship: biodiversity scales with habitat area
15.4	$H' = -\sum (p_i \ln p_i)$	Shannon diversity index: information content of species abundances
15.5	$V_{\text{eco}} = \text{NPP} / \delta_a^{\text{oo}}$	Ecosystem viability: primary production per unit disturbance rate

Photosynthesis (Level 6) $\rightarrow$ NPP $\rightarrow$ Herbivore $\rightarrow$ Predator [trophic energy chain]

► The $\approx 10\%$ ecological efficiency rule means each trophic level retains $\approx 10\%$ of the energy of the level below. A food chain of length L requires 10^L times the basal plant productivity to sustain a top predator — a direct spectral constraint on food web architecture.

THE LIFE EMERGENCE CASCADE

One persistence problem. Infinite solutions. One chain.



LIFE MASTER EQUATION

$$d\Pi/dt = f(\Delta G, \Phi, w, \mu) - \delta\Pi$$

Persistence Π grows through free-energy capture, organisational closure, fitness, and heritable information — and decays through disturbance δ .

UNIFIED BRIDGE: PHYSICS $\Rightarrow$ LIFE



$\Omega = \lim e^{-n\beta} L U$ <i>Spectral attractor</i>		$d\Pi/dt = f(\Delta G, \Phi, w, \mu) - \delta\Pi$ <i>Persistence dynamics</i>		$M \rightarrow M(M)$ <i>Self-reference</i>
--	--	--	--	---