

Homework 3

Instructor: Lijun Zhang

Name: 高辰潇, StudentId: 181220014

Notice

- The submission email is: **njuoptfall2019@163.com**.
- Please use the provided L^AT_EX file as a template. If you are not familiar with L^AT_EX, you can also use Word to generate a PDF file.

Problem 1: Equality Constrained Least-squares

Consider the equality constrained least-squares problem

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|Ax - b\|_2^2 \\ & \text{subject to} && Gx = h \end{aligned}$$

where $A \in \mathbf{R}^{m \times n}$ with $\text{rank } A = n$, and $G \in \mathbf{R}^{p \times n}$ with $\text{rank } G = p$.

- Derive the Lagrange dual problem with Lagrange multiplier vector v .
- Derive expressions for the primal solution x^* and the dual solution v^* .

Solution.

a) 解:

该问题的 Lagrange 函数为 $L(x, v) = \frac{1}{2}(Ax - b)^T(Ax - b) + v^T(Gx - h) = \frac{1}{2}x^T A^T A x + (v^T G - b^T A)x + \frac{1}{2}b^T b - v^T h$, $\nabla_x L(x, v) = A^T A x + G^T v - A^T b$.

得到拉格朗日函数的极小值点为 $x = (A^T A)^{-1}(A^T b - G^T v)$.

因此 $g(v) = \inf_{x \in D} L(x, v) = \frac{1}{2}(v^T G - b^T A)(A^T A)^{-1}(A^T b - G^T v) + \frac{1}{2}b^T b - v^T h$.

所以对偶问题为

$$\text{maximize} \quad \frac{1}{2}(v^T G - b^T A)(A^T A)^{-1}(A^T b - G^T v) + \frac{1}{2}b^T b - v^T h \quad (1)$$

b) 解: 由于对偶问题是关于变量 v 的无约束优化问题, 故 $g(v)$ 对 v 求导得

$$\nabla_v g(v) = G(A^T A)^{-1}A^T b - h - (G(A^T A)^{-1}G^T)v$$

因此, $v^* = (G(A^T A)^{-1}G^T)^{-1}(G(A^T A)^{-1}A^T b - h)$.

又由 Slater 条件, 原问题满足强对偶性。进一步由 KKT 条件, 有 $A^T A x^* + G^T v^* - A^T b = 0$.

代入 v^* 得到 $x^* = (A^T A)^{-1}[A^T b - G^T(G(A^T A)^{-1}G^T)^{-1}(G(A^T A)^{-1}A^T b - h)]$

□

Problem 2: Support Vector Machines

Consider the following optimization problem

$$\text{minimize} \quad \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b)) + \frac{\lambda}{2} \|w\|_2^2$$

where $x_i \in \mathbf{R}^d, y_i \in \mathbf{R}, i = 1, \dots, n$ are given, and $w \in \mathbf{R}^d, b \in \mathbf{R}$ are the variables.

a) Derive an equivalent problem by introducing new variables $u_i, i = 1, \dots, n$ and equality constraints

$$u_i = y_i(w^T x_i + b), i = 1, \dots, n.$$

b) Derive the Lagrange dual problem of the above equivalent problem.

c) Give the Karush-Kuhn-Tucker conditions.

Hint: Let $\ell(x) = \max(0, 1 - x)$. Its conjugate function $\ell^(y) = \sup_x (yx - \ell(x)) = \begin{cases} y, & -1 \leq y \leq 0 \\ \infty, & \text{otherwise} \end{cases}$*

Solution.

a) 解：等价问题为

$$\begin{aligned} &\text{minimize} \quad \sum_{i=1}^n \max(0, 1 - u_i) + \frac{\lambda}{2} \|w\|_2^2 \\ &\text{subject to} \quad u_i = y_i(w^T x_i + b), i = 1, \dots, n. \end{aligned}$$

b) 解：拉格朗日函数为 $L(v, u_i, w) = \sum_{i=1}^n \max(0, 1 - u_i) + \frac{\lambda}{2} \|w\|_2^2 + \sum_{i=1}^n v_i[u_i - y_i(w^T x_i + b)]$.

$$g(v) = \inf_w \{ \sum_{i=1}^n \max(0, 1 - u_i) + \sum_{i=1}^n v_i u_i \} + \inf_w \{ \frac{\lambda}{2} \|w\|_2^2 - \sum_{i=1}^n v_i y_i w^T x_i \} + \inf_b \{ -v_i y_i b \}$$

$$\nabla_w L = \lambda w - \sum_{i=1}^n (v_i y_i x_i), \text{ 故极值点为 } w = \frac{1}{\lambda} \sum_{i=1}^n v_i y_i x_i$$

$$\text{由 Hint 知 } \inf_{u_i} \{ \max(0, 1 - u_i) + y_i u_i \} = -\sup_{u_i} \{ -v_i u_i - \max(0, 1 - u_i) \} = \begin{cases} v_i, & 0 \leq v_i \leq 1 \\ -\infty, & \text{otherwise} \end{cases}$$

$$\text{而 } \inf_b \{ -v_i y_i b \} = \begin{cases} 0, & v_i y_i = 0 \\ -\infty & \text{otherwise} \end{cases}$$

因此 $g(v) = \sum_{i=1}^n v_i + \frac{1}{2\lambda} (\sum_{i=1}^n v_i y_i x_i)^T (\sum_{i=1}^n v_i y_i x_i) - \sum_{i=1}^n \frac{1}{\lambda} v_i^2 y_i^2 x_i^T x_i$, 其中 $v_i \in [0, 1], v_i y_i = 0, i \in \{1, \dots, n\}$

故对偶问题为：

$$\begin{aligned} &\text{maximize} \quad g(v) = \sum_{i=1}^n v_i + \frac{1}{2\lambda} \left(\sum_{i=1}^n v_i y_i x_i \right)^T \left(\sum_{i=1}^n v_i y_i x_i \right) - \sum_{i=1}^n \frac{1}{\lambda} v_i^2 y_i^2 x_i^T x_i \\ &\text{subject to} \quad 0 \leq v_i \leq 1 \\ &\quad \quad \quad y_i v_i = 0 \end{aligned} \tag{2}$$

c) 解：KKT 条件为：

$$1) \quad u_i^* = y_i((w^*)^T x_i + b), i \in \{1, \dots, n\}$$

2)

□

Problem 3: Euclidean Projection onto the Simplex

Consider the following optimization problem

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|y - x\|_2^2 \\ & \text{subject to} && \mathbf{1}^T y = r \\ & && y \succeq 0 \end{aligned}$$

where $r > 0$, $x \in \mathbb{R}^n$ is given, and $y \in \mathbf{R}^n$ is the variable. Give an algorithm to solve this problem and prove the correctness of your algorithm.

Hint: Derive the Lagrangian of this problem and apply the Karush-Kuhn-Tucker conditions. If you need more hints, please read the following paper [1]

Algorithm 1 Solution

Input: $r > 0, x \in \mathbb{R}^n = (x_1, x_2, \dots, x_n)$

- 1: Sort x into u : $u_1 \geq u_2 \geq \dots \geq u_n$
- 2: $j = 1$
- 3: **while** $u_j + \frac{1}{j}(1 - \sum_{i=1}^j u_i) > 0$ **do**
- 4: $j = j + 1$
- 5: **end while**
- 6: $\rho = j - 1$
- 7: Define $v = -\frac{1}{\rho}(1 - \sum_{i=1}^{\rho} u_i)$

Output: optimal y^* s.t. $y_i = \max\{x_i - v, 0\}, i = 1, \dots, n$.

Solution. 解：首先给出算法。(见 Solution)

下面证明算法正确性。

该问题的拉格朗日函数为 $L(y, \lambda, v) = \frac{1}{2} \|y - x\|_2^2 - \lambda^T y + v(1^T y - r)$.

故对偶函数 $g(\lambda, v) = \inf_{y \in D} L(y, \lambda, v) = -\frac{1}{2}(x^T + \lambda^T - v \cdot \mathbf{1}^T)(x + \lambda - v \cdot \mathbf{1})$

故原问题的对偶问题为

$$\begin{aligned} & \text{maximize} && g(\lambda, v) \\ & \text{subject to} && \lambda \succeq 0 \end{aligned} \tag{3}$$

由于原问题为凸问题，且满足 Slater 条件，因此原问题的最优解 y^* 和对偶问题的最优解 (λ^*, v^*) 应满足 KKT 条件：

$$y^* \succeq 0$$

$$\mathbf{1}^T y^* = r$$

$$\lambda^* \succeq 0$$

$$\lambda^* y^* = 0$$

$$y^* - x - \lambda^* + v^* \mathbf{1} = 0$$

考虑最优解 y^* 的各分量 y_i ，若 $y_i = 0$ ，则由条件 4 知 $\lambda_i \geq 0$ ，故 $v^* - x_i \geq 0$ ；若 $y_i > 0$ ，则 $\lambda_i = 0$ ， $v^* - x_i = -y_i < 0$ 。

不失一般性，将给定的向量 x 按照从大到小的顺序重新排列各分量，设排列后得到新向量 $u = (u_1, u_2, \dots, u_n)$ 。

按照原本各维度的对应关系相应地排序最优解，设最优解为 $y^* = y_1, \dots, y_n$ 。由上述分析知，在 $\lambda_i \geq 0$ 的情况下数值较小的 u_i 一定对应着较小的 y_i 。因此有 $y_1 \geq y_2 \geq \dots \geq y_\rho > y_{\rho+1} = \dots = y_n = 0$ 。

又由于 $1^T y^* = y_1 + y_2 + \dots + y_\rho$, 且对于 y_1, \dots, y_m 有 $y_i = x_i - v^*$, 因此 $\sum_{i=1}^\rho (u_i - v^*) = 1$,
 $v^* = -\frac{1}{\rho}(1 - \sum_{i=1}^\rho u_i)$

为了确定 v^* , 还需要确定 ρ 的大小。对于 $j \in \{1, 2, \dots, n\}$ 分三种情况考虑:

a) 若 $j < \rho$, 则

$$\begin{aligned} & \frac{1}{j}(1 - \sum_{i=1}^j u_i) + u_j \\ &= \frac{1}{j}(1 - \sum_{i=1}^\rho u_i + \sum_{j+1}^\rho u_i + ju_j) \\ &= \frac{1}{j}(-\rho v^* + \sum_{i=j+1}^\rho u_i + ju_j) \\ &= \frac{1}{j}(j(u_j - v^*) + \sum_{i=j+1}^\rho (u_i - v^*)) \end{aligned} \quad (4)$$

由于对于 $i \in \{1, 2, \dots, \rho\}$ 和 j 都有 $u_i - v^* = y_i > 0$, 故 $\frac{1}{j}(1 - \sum_{i=1}^j u_i) + u_j > 0$.

b) 若 $j > \rho$, 则

$$\begin{aligned} & \frac{1}{j}(1 - \sum_{i=1}^j u_i) + u_j \\ &= \frac{1}{j}(1 - \sum_{i=1}^\rho u_i - \sum_{\rho+1}^j u_i + ju_j) \\ &= \frac{1}{j}(-\rho v^* - \sum_{i=\rho+1}^j u_i + ju_j) \\ &= \frac{1}{j}(j(u_j - v^*) - \sum_{i=\rho+1}^j (u_i - v^*)) \end{aligned} \quad (5)$$

由于对于 $i \in \{\rho+1, \dots, n\}$ 和 $j > \rho$ 都有 $u_i - v^* \leq 0$, 故 $\frac{1}{j}(1 - \sum_{i=1}^j u_i) + u_j < 0$.

c) 若 $j = \rho$, 则有

$$\frac{1}{\rho}(1 - \sum_{i=1}^\rho u_i) + u_\rho = \frac{1}{\rho}(\rho(u_\rho - v^*)) > 0.$$

因此, 算法第 3-6 行通过判断 $\frac{1}{j}(1 - \sum_{i=1}^j u_i) + u_j$ 的正负性, 找到使该式为正值的最大的下标, 该下标即为 ρ . 再由 $v^* = -\frac{1}{\rho}(1 - \sum_{i=1}^\rho u_i)$ 可进一步计算出 v^* (算法的第 7 行)。

而对于原问题最优解 y^* , 由 KKT 条件知它的非零分量 $y_i (i \leq \rho)$ 均满足 $y_i = x_i - \rho$, 且对于它的为零的分量 $y_i (i > \rho)$ 有 $x_i - v^* \leq 0$. 因此, y^* 的各分量 y_i 可统一表示为 $y_i = \max\{x_i - v^*, 0\}$.

故算法的输出即为原问题的最优解。正确性得证。

□

Problem 4: Optimality Conditions

Consider the problem

$$\begin{aligned} & \text{minimize} && \text{tr}(2X) - \log \det(3X) \\ & \text{subject to} && 2Xs = y \end{aligned}$$

with variable $X \in \mathbf{S}^n$ and domain \mathbf{S}_{++}^n . Here, $y \in \mathbf{R}^n$ and $s \in \mathbf{R}^n$ are given, with $s^T y = 1$.

- a) Give the Lagrange and then derive the Karush-Kuhn-Tucker conditions.
 b) Verify that the optimal solution is given by

$$X^* = \frac{1}{2} \left(I + yy^T - \frac{ss^T}{s^Ts} \right).$$

Solution.

- a) 解: Lagrange 函数为 $L(X, v) = \text{tr}(2X) - \log \det(3X) + v^T(2Xs - y)$, $\nabla_X L = 2I - X^{-1} + vs^T + sv^T$.

KKT 条件为:

- 1) $2Xs = y$
- 2) $2I - X^{-1} + vs^T + sv^T = 0$

其中 $X \succ 0$

- b) 解:

由 $2Xs = y$ 知 $s = \frac{1}{2}X^{-1}y$

代入 $X^{-1} = 2I + vs^T + sv^T$ 得到 $s = \frac{1}{2}(2I + vs^T + sv^T)y = y + \frac{1}{2}(v + (v^Ty)s)$.

$s^Ty = y^Ty + \frac{1}{2}v^Ty + \frac{1}{2}v^Ty = 1$, 所以有 $y^Ty + v^Ty = 1$.

代入 $s = y + \frac{1}{2}(v + (v^Ty)s)$ 可得到 $v = (1 + y^Ty)s - 2y$.

下面证明, 题中所给出的 X^* 满足 KKT 条件。

- 1) $2Xs = 2 \cdot \frac{1}{2}(I + yy^T - \frac{ss^T}{s^Ts})s = s + y - s = y$.
- 2) 等价于证明 $X^{-1}X = (2I + v^Ts + s^Tv)X = I$

$$X^{-1}X = (2I + (v^*)^Ts + s^Tv)X = I$$

$$= (2I + 2(1 + y^Ty)ss^T - 2ys^T - 2sy^T)(\frac{1}{2}(I + yy^T - \frac{ss^T}{s^Ts}))$$

$$= I + (1 + y^Ty)ss^T - ys^T - sy^T + yy^T + (1 + yy^T)sy^T - yy^T - sy^Tyy^T - \frac{ss^T}{s^Ts} - (1 + y^Ty)ss^T + ys^T + \frac{ss^T}{s^Ts} \quad (6)$$

因此题目所给定的 X 满足 KKT 条件。又由于 $X \succ 0$, 故 X 是原问题的最优解。

□

参考文献

- [1] gugu Weiran Wang, and Miguel Á. Carreira-Peroiñán. Projection onto the probability simplex: An efficient algorithm with a simple proof, and an application. *arXiv:1309.1541*, 2013.