

1* Regarding pharynx anatomy which of the following statements is incorrect:

- a) It extends from base of skull to inferior border of cricoid cartilage anteriorly and inferior border of C6 posteriorly
- b) Widest portion (5cm) is at hyoid level
- c) Narrowest portion (1.5cm) is at its caudal end
- d) The pharynx is divided into 3 parts: nasopharynx, oropharynx, and laryngo(hypo)pharynx
- ☒ The pharynx has more than 20 cm. in length

2*. The structures of pharyngeal wall are the following, with one exception. Indicate the exception:

- a) Mucous membrane
- b) Pharyngeal aponeurosis (pharyngobasilar fascia)
- c) Muscular fascia
- d) Buccopharyngeal fascia
- ☒ Periosteum

3. Waldeyer's ring is composed on the following elements:

- ☒ Nasopharyngeal tonsil or the adenoids
- ☒ Palatine tonsils or simply the tonsils
- c) Gastric tonsils
- d) Mastoid tonsils
- ☒ Tubal tonsil (in fossa of Rosenmüller)

4*. Waldeyer's ring is composed on the following elements, with one exception. Indicate the exception:

- ☒ Submaxillary glands
- b) Nasopharyngeal tonsil or the adenoids
- c) Palatine tonsils or simply the tonsils
- d) Tubal tonsil (in fossa of Rosenmüller)
- e) Nodules (in posterior pharyngeal wall)

5*. Nasopharyngeal functions are the following, with one exception. Indicate the exception:

- a) Conduit for air (warmed and humidified)
- b) Ventilates the middle ear through Eustachian tube
- c) Resonating chamber during voice production
- ☒ Passage for foods and liquids
- e) Drainage channel for the mucus secreted by nasal and nasopharyngeal glands

6*. Regarding oropharyngeal functions the following answer is incorrect:

- a) Conduit for passage of air or food
- b) Helps in pharyngeal phase of deglutition
- c) Provide local defense and immunity against harmful intruders into the air and food passages
- d) Forms part of vocal tract for certain speech sounds
- ☒ Helps in appreciation of the smell

7. The hypopharynx is clinically subdivided into the following regions:

- ☒ Pyriform sinus (fossa): lies on the either side of the larynx, expands from pharyngoepiglottic fold to the upper end of esophagus
- b) Vocal cord
- ☒ Post-cricoid region: the part of the anterior wall of laryngopharynx between the upper and lower borders of cricoid lamina
- d) Ventricular fold
- ☒ Posterior pharyngeal wall: extends from the level of hyoid bone to the level of cricoarytenoid joint

8. Adenoiditis aural symptoms are:

- ☒ Tubal obstruction
- b) Sinusitis
- ☒ Recurrent attacks of acute otitis media
- d) Nasal discharge
- ☒ Serous otitis media

9. Regarding Juvenile Nasopharyngeal Angiofibroma the following statements are correct:

- ☒ The exact cause is unknown
- b) Age and sex (exclusively in males in the age group of 40-60 years)
- ☒ Appears progressive nasal obstruction and nasal speech
- ☒ Biopsy of the tumor is associated with profuse bleeding and is, therefore, avoided
- e) The treatment is always surgical

10. Regarding Nasopharyngeal Cancer the following statements are correct:

- ☒ Age of onset: 5th to 7th decades, but may involve younger age groups
- ☒ Cervical nodal metastases may be the only manifestation
- ☒ Otologic symptoms and signs are represented by the conductive hearing loss, serous and suppurative otitis media
- d) Ophtho-neurologic symptoms are always present at debut
- e) Distant metastases are into the Eustachian tube

11. Which of the following are symptoms of acute tonsillitis:

- ☒ Sore throat
- ☒ Difficulty in swallowing
- c) Oral breathing
- ☒ Fever
- ☒ Earache

12. Which of the following are acute tonsillitis complications:

- a) Convergent strabismus
- ☒ Rheumatic fever
- ☒ Chronic tonsillitis
- ☒ Acute glomerulonephritis
- ☒ Subacute bacterial endocarditis

13. Differential diagnosis of acute tonsillitis is performed with:

- ☒ Membranous tonsillitis
- ☒ Diphtheria
- c) Rhinitis
- ☒ Vincent's angina
- ☒ Infectious mononucleosis

14. Differential diagnosis of acute tonsillitis is performed with:

- a) Sinusitis
- ☒ Bucopharyngeal manifestations in Agranulocytosis
- ☒ Bucopharyngeal manifestations of Leukemia
- ☒ Aphthous ulcers
- ☒ Malignant tumor of the tonsil

15. Regarding hypopharyngeal malignant tumors, which of the following statements are correct?

- ☒ Clinical features are represented by progressive dysphagia, malnutrition and weight loss, change of voice and disphonia
- ☒ Mirror examination may not be useful
- c) The most frequent site of origin is subglottic region
- ☒ Prognosis is poor
- ☒ Surgery, Radiotherapy, and combined treatment

16. *The nasal septum divides the nose into:

- a) superior and inferior nasal cavity
- ☒ right and left nasal cavity
- c) lateral and medial cavity
- d) four cavities 2 superior and 2 inferior
- e) the nasal septum is virtual

17. *The olfactory nerves, (12-20) nerves which pass through the cribriform plate and end in the:

- a) frontal lobe
- b) temporal lobe
- c) olfactory lobe
- d) parietal lobe
- ☒ olfactory bulb

18. *The rhinophyma is a:

- a) benign, slow growing mass
- b) slow growing malign tumor
- ☒ appears due to the hypertrophy of the sebaceous glands of the tip of the nose
- ☒ is treated with surgical excision or with laser CO₂
- e) appears due to the hypertrophy of the perspiratory glands of the tip of the nose

19. *The maxillary sinus:

- ☒ opens into the posterior part of ethmoidal infundibulum into the middle meatus
- b) opens into the superior meatus
- c) its innervations is from the IX-th cranial nerve
- d) it is the smallest sinus
- e) has a 5 ml capacity

20. The choanal atresia:

- ☒ can be: uni-, bilateral, complete, incomplete, bony (90%) or membranous (10%)
- b) it's always bilateral, complete, bony or membranous
- ☒ it is an emergency in bilateral choanal atresia
- ☒ one of the diagnostic signs is the absence of air bubbles in the nasal discharge
- ☒ an important diagnostic sign is the failure to pass a catheter from nose to pharynx

21. *A dermoid cyst is:

- a) a malign tumor
- ☒ a congenital tumor
- c) an infection
- d) a blood collection
- e) a sign of consumption

22. * Into the middle meatus open the following sinuses:

- ☒ maxillary, frontal and anterior ethmoidal
- b) posterior ethmoidal sinuses
- c) sphenoid sinus
- d) cavernous sinus
- e) sinus Tympani

23. The septal haematoma:

- ☒ is a collection of blood under the perichondrium or periosteum of the nasal septum
- ☒ it can be caused by nasal trauma or septal surgery
- ☒ the treatment consists in aspiration with a wide bore sterile needle
- ☒ incision and drainage in larger ones with bilateral nasal packing
- e) is a collection of connective tissue

24. About the nasal skin, which affirmations are correct?

- ☒ thin skin and freely mobile, that covers the nasal bones
- b) thin skin with many sebaceous glands that covers the nasal bones and the lower lateral cartilages
- ☒ thick, adherent and rich in sebaceous glands skin, that covers the alar cartilages
- d) thin and adherent skin, covering the alar cartilages
- ☒ thick skin rich in sebaceous glands that covers the nasal bones and the upper lateral cartilages

25. The rhinoscleroma:

- ☒ its etiology is gram negative bacillus - Klebsiella Rhinoscleromatis
- b) its etiology is gram + Staphylococcus Aureus
- ☒ its clinical stages are: atrophic, granulomatous, cicatricial
- ☒ its certain diagnosis is given by biopsy
- ☒ it is treated with streptomycin + tetracycline

26. About the treatment in allergic rhinitis, the following are correct:

- ☒ a) avoidance of allergen
- ☒ b) supraexposure to allergen
- ☒ c) antihistaminics
- ☒ d) corticosteroids (oral and topical)
- ☒ e) radiotherapy

27. * The first step in the management of anterior epistaxis is:

- ☒ a) posterior nasal packing
- ☒ b) cauterization
- ☒ c) pinching the nose with thumb and index finger for 5 minutes
- ☒ d) ligation of vessels
- ☒ e) anterior nasal packing

28. The symptoms of vasomotor rhinitis are:

- ☒ a) wheezing
- ☒ b) excessive rhinorrhea
- ☒ c) nasal obstruction
- ☒ d) postnasal drip
- ☒ e) cough

29. The local causes of epistaxis are:

- ☒ a) trauma/foreign bodies
- ☒ b) infections
- ☒ c) neoplasms of nose and paranasal sinuses
- ☒ d) heat
- ☒ e) cold

30. The posterior epistaxis:

- ☒ a) originates mostly from posterosuperior part of nasal cavity, often difficult to localize the bleeding point
- ☒ b) occurs spontaneous often due to hypertension or arteriosclerosis
- ☒ c) the most common cause is trauma
- ☒ d) mostly occurs in children or young adults
- ☒ e) is usually mild and easy to control

31. The complications of frontal sinusitis are:

- ☒ a) orbital cellulites
- ☒ b) osteomyelitis of frontal bone
- ☒ c) meningitis, extradural abscess, or frontal lobe abscess
- ☒ d) maxillary sinus necrosis
- ☒ e) acute conjunctivitis

32. Choose the correct affirmations about laryngomalacia:

- ☒ a) Most common congenital laryngeal anomaly (50-75%)
- ☒ b) Most frequent cause of stridor in adults
- ☒ c) Male predominance 2:1
- ☒ d) Flaccidity of supraglottic laryngeal tissues
- ☒ e) Characterized by inward collapse of pharynx structures

33. Subglottic haemangiomas are:

- ☒ a) Asymptomatic till 12 months
- ☒ b) 50% have associated cutaneous haemangiomas
- ☒ c) Stridor, but a normal cry
- ☒ d) Reddish-blue mass below the vocal cords
- ☒ e) Biopsy - strongly indicated

34. * The main indication for tracheotomy is:

- ☒ a) To administrate anesthesia
- ☒ b) To bypass upper airway obstruction
- ☒ c) Nasal obstruction
- ☒ d) to treat malignancy
- ☒ e) to treat hearing loss

35. * The epiglottis is formed by:

- ☒ a) Muscle
- ☒ b) Bone
- ☒ c) Fibrous cartilage
- ☒ d) Adipose tissue
- ☒ e) Skin

36. * The most common cause of bilateral recurrent laryngeal nerve paralysis is:

- ☒ a) Laryngeal trauma
- ☒ b) Surgical iatrogenic damage in case of total thyroidectomy due to thyroid gland carcinoma
- ☒ c) A tumor mass expanding in upper part of mediastinum
- ☒ d) Toxic polyneuropathy of recurrent laryngeal nerves
- ☒ e) Acute subglottic laryngitis

37. Larynx functions are:

- ☒ a) phonation
- ☒ b) respiration
- ☒ c) inferior respiratory airway protection
- ☒ d) thoracic fixation
- ☒ e) expectoration

38. Chronic non specific laryngitis can be:

- ☒ a) catarrhal
- ☒ b) caused by tuberculosis
- ☒ c) hypertrophic-red and white type
- ☒ d) atrophic
- ☒ e) caused by syphilis

39. Vocal nodules:

- ☒ a) are considered inflammatory tissue reactions
- ☒ b) are favored by voice overload
- ☒ c) re surgically treated
- ☒ d) voice rest is recommended
- ☒ e) more frequent in male patients

40. Complications of laryngotracheal trauma are represented by:

- ☒ a) Laryngeal stenosis
- ☒ b) Loss of taste
- ☒ c) Perichondritis / Laryngeal abscess
- ☒ d) Vocal cord paralysis
- ☐ e) Hiposmia

41. Congenital lesions of the larynx are:

- ☒ a) Laryngeal Web
- ☒ b) Laryngomalacia
- ☐ c) Choanal atresia
- ☐ d) Cleft palate
- ☒ e) Laryngocele

42. Larynx benign tumors are:

- ☐ a) Adenocarcinoma
- ☒ b) Chondroma
- ☒ c) Haemangioma
- ☐ d) Tracheostomy
- ☐ e) Squamous cell carcinoma

43. Laryngotracheal trauma can be represented by:

- ☒ a) Haematoma and edema of the larynx
- ☐ b) Othematoma
- ☒ c) Dislocation of cricoarytenoid joint
- ☒ d) Fractures of hyoid bones, thyroid and cricoid cartilages
- ☒ e) Fractures of upper tracheal rings

44. Which of the following statements are correct:

- ☐ a) In Weber test the tuning fork is placed on the mastoid process
- ☒ b) In Weber test a patient with normal hearing localizes the tone either in the center of the head or equally in both ears
- ☒ c) In Weber test a patient with unilateral conductive hearing loss localizes the tone in the affected ear
- ☐ d) In Weber test a patient with neurosensory hearing loss localizes the tone in the affected ear
- ☒ e) In Weber test a patient with neurosensory hearing loss localizes the tone in the healthy ear

45. Following are intracranial complications of otitis media

- ☒ a) extradural abscess
- ☐ b) petrositis
- ☒ c) meningitis
- ☒ d) labyrinthitis
- ☐ e) brain abscess

46. Following are intratemporal complications of otitis media

- ☒ a) mastoiditis
- ☒ b) petrositis
- ☒ c) subdural abscess
- ☐ d) facial paralysis
- ☐ e) meningitis

47. Following are signs of acute mastoiditis

- ☒ a) mastoid tenderness
- ☒ b) purulent ear discharge
- ☒ c) sagging of posterosuperior meatal wall and conductive hearing loss
- ☐ d) blood counts show monocytes and lymphocyte leucocytosis
- ☐ e) neurosensory hearing loss

48. Acute mastoiditis

- ☒ a) Treatment consists of antibiotics
- ☒ b) Treatment consists of myringotomy
- ☐ c) Treatment consists of stapedotomy
- ☒ d) Treatment consists of cortical mastoidectomy
- ☐ e) Treatment consists of reconstructive surgery with hearing restoration by tympanoplasty

49. Abscesses in relation to mastoid infection are:

- ☒ a) postauricular abscess
- ☒ b) zygomatic abscess
- ☒ c) Bezold abscess
- ☐ d) petrositis
- ☐ e) orbital cellulitis

50. Otosclerosis is characterized by:

- ☒ a) conductive hearing loss
- ☒ b) paracusis Willisii
- ☒ c) tinnitus
- ☐ d) Eustachian tube function is impaired
- ☐ e) ossification and calcification of subepithelial connective tissue, white chalky deposits in the promontory and ossicles