

## Variant 2

UN SINGUR RASPUNS:

Single Answer

1. Which of the following etiologies is the most frequent in kidney cancer
  - a) transitional carcinoma
  - b) papillary adenocarcinoma
  - c) **clear cell adenocarcinoma**
  - d) chromophobe adenocarcinoma
  - e) embryonic carcinoma
  
2. The most frequent neoplasm of prostate metastasis:
  - a) hepatic
  - b) cerebral
  - c) **bone**
  - d) lung
  - e) skin
  
3. Which of the following is not a negative prognostic element for a patient with bronchopulmonary adenocarcinoma
  - a) Unintentional weight loss of 10 kg in the last 6 months
  - b) **EGFR mutation present**
  - c) Performance index ECOG 3
  - d) Contralateral lung nodule
  - e) Hepatic metastasis
  
4. What is the frequency of hormone dependent breast cancers
  - a) Less than 5%
  - b) 20-30%
  - c) 40-50%
  - d) **60-70%**
  - e) Over 90%
  
5. Which is the principal risk factor in cervical cancer
  - a) Prolonged exposure to estrogen
  - b) Prolonged exposure to sun rays
  - c) **Unprotected sexual contact**
  - d) Infection with HP
  - e) Diet rich in animal proteins and fats
  
6. The most frequent type of hepatic cancer is
  - a) **Hepatocarcinoma**
  - b) Cholangiocarcinoma
  - c) Hepatoblastoma

- d) Angiosarcoma
- e) Lymphoma

7. Paraneoplastic syndromes that can be found in gastric cancer are
- a) acanthosis nigricans
  - b) dermatomyositis
  - c) Trousseau Syndrome
  - d) Nephrotic Syndrome
  - e) All of the above
8. Which of the following affirmations about biochemical resistance to chemotherapy is not true
- a) The tumor can be incapable of converting the antineoplastic agent to its active form
  - b) Pleiotropic resistance is given by a type of membrane pump cellular structure
  - c) the cell may increase the level of target enzymes that the cytotoxic agent intends to inactivate
  - d) resistance is due to deficit in irrigation of malignant tissue
  - e) the cell may increase the expression of DNA repair enzymes
9. Which is the most frequent clinical sign at debut of breast cancer
- a) tumoral breast nodule
  - b) axillary adenopathy
  - c) nipple leakage
  - d) carcinomatous mastitis
  - e) Peau D'Orange
10. Which of the following medications are antitumoral antibiotics
- a) mustard nitrogen analogues
  - b) nitrosureas
  - c) purine analogs
  - d) anthracyclines
  - e) taxanes
11. Which of the following antiemetic medications is a neurokinin 1 antagonist (NK-1)
- a) Ondansetron
  - b) Aprepitant
  - c) Metoclopramide
  - d) Palonosetron
  - e) Prochlorperazine
12. Which of the following medications is a cytostatic
- a) Tamoxifen
  - b) Paclitaxel
  - c) Trastuzumab
  - d) Imatinib
  - e) Exemestane

13. Bortezomib is
  - a) Inhibitor of mTOR kinase
  - b) Monoclonal antibody
  - c) Tyrosine kinase inhibitor
  - d) A cytokine
  - e) **Proteasome inhibitor**
  
14. CA 19-9 levels rise most frequently in
  - a) Lung cancer
  - b) Ovarian cancer
  - c) Breast cancer
  - d) Thyroid cancer
  - e) **Pancreatic cancer**
  
15. Aromatase inhibitors
  - a) **Reduce the level of estrogen through the blockage of synthesis**
  - b) Reduce the level of estrogen through the binding of estrogen receptors
  - c) Reduce the level of estrogen through the binding of estrogen-progesterone receptors
  - d) Reduce the level of estrogen through chemical castration
  - e) Reduce the level of estrogen through chemical castration and binding of estrogen receptors

**MULTIPLE RESPONSES:**

16. Carcinoembryonic antigens have frequent values ... raised in individuals
  - a) **with colorectal cancer**
  - b) **with lung cancer**
  - c) **with hepatic cirrhosis**
  - d) non-smokers
  - e) with bacterial meningitis
  
17. The following affirmations are true about MRI
  - a) Are based on the usage of x rays
  - b) **Are based on the property of hydrogen atoms in a magnetic field**
  - c) **Is contraindicated in the presence of metallic deposits and pacemakers**
  - d) **Is more effective than CT in evaluating brain or pelvic lesions**
  - e) The time of an MRI is very short
  
18. A series of criteria that microscopically suggests malignancy
  - a) **Cellular pleomorphism**
  - b) Decreased mitotic activity
  - c) **Nuclear pleomorphism**
  - d) Conservation of tissue architecture
  - e) **Nuclear hyperchromic**
  
19. Immunohistochemistry
  - a) Is utilized only for testicular tumors...
  - b) **Permits the detection of certain antigens expressed at the cytoplasmic or nuclear**

- membrane level
  - c) Permits the phenotyping of the tumor and its classification into a certain tumor type
  - d) Is a complementary exam useful in tumor anaplasia with a non-precise point of exit
  - e) Requires an initial scintigraphic exam
20. Grading of malignant tumors:
- a) Appreciates cellular anaplasia
  - b) Appreciates the number of mitoses
  - c) Appreciates the size of the primary tumor
  - d) Appreciates the nuclear pleomorphism
  - e) Appreciates the existence of distant metastasis
21. The following are examples of analytical epidemiological studies
- a) Incidence studies
  - b) descriptive studies
  - c) cohort studies
  - d) case-control studies
  - e) mortality studies
22. Which of the following statements are true about dietary risk factors:
- a) Consumption of vegetables and fruits increases the risk of digestive cancers
  - b) Vegetable glucan may increase the immunity of the intestinal wall
  - c) Phytoestrogens have a protective effect for breast and prostate cancer
  - d) Olive oil is an oncological risk factor
  - e) Fungal aflatoxins are the main risk factor for lung cancer
23. Post-therapeutic follow-up
- a) is also called screening
  - b) is also called follow-up
  - c) is also called primary prevention
  - d) is also called secondary prevention
  - e) is also called tertiary prevention
24. The role of surgery in the diagnosis and staging of malignant tumors
- a) is lately increasingly important as new staging techniques develop
  - b) is performed as the initial procedure in all digestive neoplasms
  - c) allows surgical biopsy
  - d) allows the subsequent tumorous excision to be performed in the same anesthetic session
  - e) is the procedure that sets the stage of TNM in AML
25. Reconstructive surgery in the oncological patient
- a) improves survival
  - b) repairs the defects
  - c) Improves quality of life
  - d) Restores functionality of affected organs or segments
  - e) Reduces tumor mass

26. Which of the following statements are true about brachytherapy:
- a) In brachytherapy, radiation sources are in intimate contact with tumor tissue
  - b) Endocavitary brachytherapy is especially used in prostate cancer
  - c) Contact brachytherapy is used in cervical cancer
  - d) Afterloading techniques protect medical staff from irradiation
  - e) Interstitial brachytherapy is used in oral cavity cancer
27. The benefits of neoadjuvant chemotherapy are:
- a) reducing the likelihood of subsequent systemic recurrence
  - b) is done after removal of the primary tumor and allows for obtaining microscopic information about the tumor
  - c) allows for evaluation of treatment efficiency by measuring the primary tumor size
  - d) reducing the mass of the primary tumor and facilitating surgical progression
  - e) is done at low doses that do not cause immediate or late complications
28. The chemotherapy combinations have the following principles:
- a) substances that generate mechanisms differing in resistance
  - b) substances that act through the same mechanism
  - c) substances with different adverse effects
  - d) substances that act at different times in the cell cycle
  - e) substances that act through different mechanisms
29. Thrombocytopenia during chemotherapy:
- a) is rarely severe or life-threatening
  - b) Determine treatment if the platelets are  $<100,000 / \text{mm}^3$
  - c) Determine treatment if the platelets are  $<10,000 / \text{mm}^3$
  - d) Determine treatment if the platelets are  $<50,000 / \text{mm}^3$
  - e) Can be corrected with platelets
30. Which of the following are risk factors for breast cancer:
- a) menstrual cycle with late onset and early menopause
  - b) Oral contraceptives
  - c) Obesity
  - d) Alcohol
  - e) coming from a poor country
31. Trastuzumab:
- a) is an anti-HER2 monoclonal antibody
  - b) is an anti-CD20 monoclonal antibody
  - c) is an anti-VEGF monoclonal antibody
  - d) may be associated with chemotherapy
  - e) is indicated in metastatic colorectal cancer
32. Which of the following statements about the prognosis of esophageal cancer are true:
- a) Esophageal cancer has a good prognosis, survival at 5 years being  $> 40\%$
  - b) TNM status is the most important prognostic factor
  - c) Decreasing of  $> 10\%$  of body weight is associated with a worse prognosis
  - d) The presence of liver metastases does not influence prognosis

- e) The location of the tumor at the lower and middle third level has a worse prognosis
33. Prognostic factors in colorectal cancer are:
- a) tumor grade, G1 and G2 tumors with a worse prognosis than G3 and G4
  - b) histopathological type, carcinoma with Signet ring cell (?) and mucinous carcinoma having a more reserved prognosis
  - c) the stage of TNM is the most important prognostic element
  - d) lymphatic or venous invasion
  - e) residual tumor
34. The risk factors for pancreatic cancer are:
- a) Smoking
  - b) hyperlipidic diet
  - c) partial gastrectomy
  - d) chronic pancreatitis
  - e) Excessive exposure to sunlight
35. Symptoms seen in gastric cancer are as follows:
- a) epigastric pain
  - b) anorexia
  - c) early satiety
  - d) food vomiting
  - e) hematuria
36. The negative prognostic factors in pancreatic cancer are:
- a) old age
  - b) location at head of pancreas
  - c) the advanced stage of the disease
  - d) poorly differentiated tumor
  - e) well-differentiated tumor
37. Which of the following is a risk factor in lung cancer
- a) liver cirrhosis
  - b) exposure to asbestos
  - c) exposure to radon
  - d) HTA
  - e) DZ
38. Which of the following statements are true of radiotherapy in lung cancer
- a) Cranial prophylactic CT is performed only in non-small cell cancer patients
  - b) Radical radiotherapy radiates the primary tumor, hilar ganglions and mediastinum
  - c) Palliative RT can be used in upper vena cava syndrome
  - d) Concomitant chemo-radiotherapy brings a benefit to sequential therapy
  - e) Radical RT is performed up to a maximum of 12 GY
39. Chemotherapy in prostate cancer

- a) is usually a solution used late after hormone therapy
  - b) is initial treatment in metastatic prostate cancer
  - c) Docetaxel and Prednisone are used in the first line of chemotherapy
  - d) Mitoxantrone can be used in patients who can not tolerate taxane therapy
  - e) uses combinations of 3-4 cytostatics that give more effective responses
40. Which of the following histologies is characteristic of a testicular cancer
- a) cell carcinoma with Signet ring cell
  - b) Yolk sac tumor
  - c) Coriocarcinoma
  - d) transitional carcinoma
  - e) Embryonic carcinoma
41. Which of the following drugs are tyrosine kinase inhibitors used in kidney neoplasm?
- a) Sunitinib
  - b) Temsirolimus
  - c) Bevacizumab
  - d) everolimus
  - e) sorafenib
42. Which of the following statements are true about endometroid adenocarcinoma:
- a) occurs in 20% of cases of endometrial cancer
  - b) has a better prognosis than clear cell adenocarcinoma
  - c) cells have increased levels of progesterone receptors
  - d) affects more often older people
  - e) associated with obesity and diabetes
43. Which of the following statements about systemic therapy in endometrial cancer are true:
- a) Systemic chemotherapy is indicated in advanced or recurrent forms
  - b) hormone therapy is particularly indicated in well differentiated tumors with positive hormone receptors
  - c) hormone therapy is indicated in the adjuvant treatment of endometrial cancer
  - d) hormone therapy is indicated in metastatic or recurrent forms
  - e) hormone therapy uses bicalutamide or GnRH analogues
44. Which of the following statements about ovarian cancer are true:
- a) Most ovarian cancers are germ cell tumors
  - b) Most ovarian cancers are stromal tumors
  - c) Most ovarian cancers are of epithelial origin
  - d) are usually chemosensitive tumors
  - e) usually detected in advanced stages
45. Which of the following statements about ovarian cancer are true:
- a) The most frequent dissemination is peritoneal seeding
  - b) the most frequent dissemination is haematogene
  - c) Pulmonary and cerebral metastases rarely occur
  - d) Ovarian cancer is most often detected in early stages, limited to the ovary

- e) the bone is the most common place of metastasis
46. What surgical procedures can be used in cervical cancer:
- a) Conization
  - b) Lumpectomy
  - c) radical hysterectomy with bilateral pelvic lymphadenectomy
  - d) sectorectomy
  - e) pelvic exenteration
47. Tumor suppression genes:
- a) may give rise to protooncogenes by mutations
  - b) control the cell cycle
  - c) are also called protooncogenes
  - d) play a role in carcinogenesis
  - e) The most well-known p53 with the role of genomic guardian
48. Anemia in cancer may be due:
- a) most often an autoimmune mechanism
  - b) direct inhibition of erythropoiesis by the tumor by cytokine production
  - c) an iron deficiency mechanism through occult hemorrhages
  - d) Edema (?)
  - e) direct MOH (?) infiltration
49. In hepatic metastases, changes may occur in:
- a) Transaminases
  - b) Bilirubin
  - c) PSA
  - d) serum albumin
  - e) clotting tests
50. The following statements regarding the treatment of colorectal cancer are true
- a) Chemotherapy is only useful in patients with unresectable tumors or metastatic disease
  - b) Radiotherapy is useful in the treatment of rectal cancer
  - c) the association of monoclonal antibodies with chemotherapy leads to prolonging the survival of patients with metastatic colorectal cancer
  - d) Chemotherapy usually uses 5-FU or capecitabine-based combinations
  - e) radiotherapy is useful in the treatment of cancer and proximal colon cancer