

- functions are the following, with one exception:
- a. Conduit for air (warmed and humidified)
 - b. Ventilates the middle ear through Eustachian tube
 - c. Resonating chamber during voice production
 - d. Passage for foods and liquids**
 - e. Drainage channel for the mucus secreted by nasal and nasopharyngeal glands

Nasopharynx, Epipharynx

10. Regarding oropharyngeal functions the following answer is incorrect:

- a. Conduit for passage of air or food
- b. Helps in pharyngeal phase of deglutition
- c. Provide local defense and immunity against harmful intruders into the air and food passages
- d. Forms part of vocal tract for certain speech sounds
- e. Helps in appreciation of the smell**

taste

11. The most common cause of bilateral recurrent laryngeal nerve paralysis is:

- a. laryngeal trauma
- b. surgical iatrogenic damage in case of total thyroidectomy**
- c. a tumor mass expanding in upper part of mediastinum
- d. toxic polyneuropathy of recurrent laryngeal nerves
- e. acute subglottic laryngitis

right in lecture!

12. The most important symptom in otosclerosis is:

- a. vertigo
- b. conductive hearing loss**
- c. sensorineural hearing loss
- d. aural fullness
- e. tinnitus

13. The election treatment in cholesteatoma is:

- a. antibiotherapy
- b. chemotherapy
- c. radiotherapy
- d. surgery**
- e. wait and scan policy

14. The nasal septum divides the nose into:

- a. superior and inferior nasal cavity
- b. right and left nasal cavity**
- c. lateral and medial cavity
- d. four cavities 2 superior and 2 inferior
- e. four cavities 2 medial and 2 lateral

15. The maxillary sinus:

- a. opens into the middle meatus**
- b. opens into the superior meatus
- c. its innervations is from the IX-th cranial nerve
- d. it is the smallest sinus
- e. has a 5 ml capacity

16. Relations of external acoustic meatus are:

- a. superiorly - middle cranial fossa**
- b. posteriorly - mastoid air cells and facial nerve**
- c. inferiorly - parotid gland**
- d. anteriorly - temporo-mandibular joint**
- e. anteriorly - mastoid air cells and facial nerve**

17. The Eustachian (PHARYNGOTYMPANIC) Tube:

- a. Connects the tympanic cavity with the nasopharynx
- b. Bony part - lateral 1/3
- c. Fibrocartilaginous part - medial 2/3
- d. Torus tubaris at the pharyngeal end
- e. Zygomatic cells in the middle part

18. Ossicles of the middle ear are:

- a. The malleus
- b. Tensor tympani
- c. The incus
- d. The stapes
- e. Stapedius

19. Predisposing factors for acute suppurative otitis media are:

- a. Recurrent attacks of common cold, upper respiratory tract infections, and exanthematous
- b. Infections of tonsils and adenoids
- c. Chronic Tympanosclerosis
- d. Nasal allergy
- e. Tumours of nasopharynx, packing of nose or nasopharynx for epistaxis

20. Signs of acute mastoiditis are:

- a. Mastoid tenderness
- b. Ear discharge - mucopurulent or purulent discharge
- c. Sagging of inferior meatal wall → superior !
- d. Swelling over the mastoid
- e. Hearing loss - conductive type

21. Otosclerosis symptoms are:

- a. Hearing loss - bilateral neurosensorial type → conductive !
- b. Paracusis willisii - patient hears better in noisy than quiet surroundings
- c. Tinnitus
- d. Vertigo
- e. Monotonous, well modulated soft speech

22. Differential diagnosis of Otosclerosis:

- a. Serous otitis media
- b. Tympanosclerosis
- c. Articular fixation of malleus head
- d. Acoustic neuroma
- e. Ossicular discontinuity

23. Meniere's disease cardinal symptoms are:

- a. Episodic vertigo
- b. Paracusis willisii - patient hears better in noisy than quiet
- c. Fluctuating hearing loss
- d. Tinnitus

e. Sense of fullness in involved ear.

24) Malignant tumors of external auditory canal are:

- a. squamous cell carcinoma
- b. basal cell carcinoma
- c. melanoma
- d. acoustic neuroma
- e. eustachian tube

- b. Cricoid
- c. Epiglottis
- d. Arytenoid
- e. Corniculate

Yes

27. Nerve supply of the larynx:

- a. Motor: recurrent laryngeal nerve
- b. Sensory: internal laryngeal nerve (above the vocal cords)
- c. Motor: recurrent laryngeal nerve (below the vocal cords)
- d. Motor: internal laryngeal nerve (above the vocal cords)
- e. Sensory: recurrent laryngeal nerve (below the vocal cords)

28. Chronic hyperaemic laryngitis treatment:

- a. Elimination of the infections of upper or lower respiratory tract
- b. Avoidance of irritating factors
- c. Surgery
- d. Voice rest and speech therapy
- e. Expectorants

29. Congenital lesions of the larynx are:

- a. Laryngomalacia
- b. Subglottic Haemangioma
- c. Laryngoesophageal cleft
- d. Pectus excavatum
- e. Laryngeal Web

→ nahi sein erst! :D

30. Treatment of polypoid corditis (reinke's edema)

- a. Smoking cessation
- b. Surgery: Cordectomy
- c. Antireflux medication
- d. Preoperative vocal therapy
- e. Surgery: vocal cord stripping

31. Benign tumors of larynx are:

- a. Squamous cell carcinoma
- b. Chondroma
- c. Haemangioma
- d. Granular cell tumor
- e. Glandular tumor

32. Etiology of larynx cancer:

- a. Tobacco, alcohol
- b. Neck trauma
- c. Previous radiation
- d. Genetic factors
- e. Occupational exposure

33. Glottic cancer symptoms are:

- a. hoarseness is an early symptom,
- b. hoarseness is an late symptom
- c. stridor and laryngeal obstruction in late stages
- d. stridor and laryngeal obstruction in early stages
- e. throat pain

34. Types of cordectomies:

c. Type 4: extended cordectomy, include total resection of unilateral vocal cord
anterior 1/3 of contralateral vocal cord

→ Type 1:

35. Indications of tracheotomy are:

- a. Respiratory obstruction
- b. Retained secretions
- c. Respiratory insufficiency
- d. Puberphonia
- e. Hoarseness

36. Types of tracheotomy are:

- a. High tracheotomy: above the level of thyroid isthmus ✓
- b. High tracheotomy: through the II and III rings ✓
- c. Mid tracheotomy: through the II and III rings ✓
- d. Mid tracheotomy: below the level of thyroid isthmus
- e. Low tracheotomy: below the level of thyroid isthmus ✓

37. The frontal sinus drains:

- a. in the superior meatus
- b. into frontal recess (55%)
- c. above but not into infundibulum (30%)
- d. into the infundibulum (15%)
- e. above the bulla ethmoidalis

38. Functions of paranasal sinuses are:

- a. Air conditioning of the inspired air: warming and humidifying
- b. Voice resonance
- c. Thermal insulators - protection of delicate structures in the orbit and cranium from variation in intranasal temperature
- d. Nasal drainage
- e. Lighten the skull bones

39. Clinical Features of acute maxillary sinusitis are:

- a. Constitutional symptoms: fever, general malaise
- b. Headache
- c. Pain - over upper jaw, occasionally ipsilateral supraorbital region ✓
- d. Tenderness (pressure over the anterior wall of antrum produces pain) ✓
- e. Post nasal discharge (pus on the hard palate at anterior rhinoscopy)

40. Complications of acute maxillary sinusitis are:

- a. Deep neck infection
- b. Subacute, chronic sinusitis
- c. Osteitis or osteomyelitis
- d. Orbital cellulitis or abscess, optic neuritis
- e. Cavernous sinus thrombosis

41. Clinical Features of acute ethmoid sinusitis are:

- a. Pain - over the bridge of the nose, medial and deep to the eye; aggravated by movement
- b. Edema of lids
- c. Hoarseness
- d. Nasal discharge
- e. Swelling of the middle turbinate

42. Waldeyer's ring consists of:

a. Nasopharyngeal tonsil or the adenoids

b. Palatine tonsils

c. Lingual tonsils

d. Tubal tonsils (in Fossa of Rosenmüller)

Zu 34 o

type I - a subepithelial cordectomy, which is resection of the epithelium;

type II - a subligamental cordectomy, which is a resection of the epithelium, Reinke's space and vocal ligament;

type III - transmuscular cordectomy, which proceeds through the vocalis muscle;

type IV - total cordectomy;

type Va - extended cordectomy, which encompasses the contralateral vocal fold and the anterior commissure;

type Vb - extended cordectomy, which includes the arytenoid;

type Vc - extended cordectomy, which encompasses the subglottis;

type Vd - extended cordectomy, which includes the ventricle.

anterior 1/3 of contralateral vocal cord

e. Type 4: extended cordectomy, include total resection of unilateral vocal cord, anterior commissure and anterior 1/3 of contralateral vocal cord

35. Indications of tracheotomy are:

- a. Respiratory obstruction
- b. Retained secretions
- c. Respiratory insufficiency
- d. Puberphonia
- e. Hoarseness

36. Types of tracheotomy are:

- a. High tracheotomy: above the level of thyroid isthmus
- b. High tracheotomy: through the II and III rings
- c. Mid tracheotomy: through the II and III rings
- d. Mid tracheotomy: below the level of thyroid isthmus
- e. Low tracheotomy: below the level of thyroid isthmus

37. The frontal sinus drains:

- a. in the superior meatus
- b. into frontal recess (55%)
- c. above but not into infundibulum (30%)
- d. into the infundibulum (15%)
- e. above the bulla ethmoidalis

38. Functions of paranasal sinuses are:

- a. Air conditioning of the inspired air: warming and humidifying
- b. Voice resonance
- c. Thermal insulators - protection of delicate structures in the orbit and cranium from variation of intranasal temperature
- d. Nasal drainage
- e. Lighten the skull bones

39. Clinical Features of acute maxillary sinusitis are:

- a. Constitutional symptoms: fever, general malaise
- b. Headache
- c. Pain - over upper jaw, occasionally ipsilateral supraorbital region
- d. Tenderness (pressure over the anterior wall of antrum produces pain)
- e. Post nasal discharge (pus on the hard palate at anterior rhinoscopy)

40. Complications of acute maxillary sinusitis are:

- a. Deep neck infection
- b. Subacute, chronic sinusitis
- c. Osteitis or osteomyelitis
- d. Orbital cellulitis or abscess, optic neuritis
- e. Cavernous sinus thrombosis

41. Clinical Features of acute ethmoid sinusitis are:

- a. Pain - over the bridge of the nose, medial and deep to the eye; aggravated by movement
- b. Edema of lids
- c. Hoarseness
- d. Nasal discharge
- e. Swelling of the middle turbinate

Waldeyer's ring consists of:

Uvula, tonsils, adenoids

Same page!

Functions of nasopharynx, oropharynx

(a) warm, humid.

(b) protects ear

→ ungedrehte Stenographie!

(c) cleanses air

② mucous secreted by nasal + nasopharynx

14. The nasopharynx is the part of the pharynx that is located behind the oral cavity and above the larynx.

- a. The nasopharynx is the part of the pharynx that is located behind the oral cavity and above the larynx.
- b. The nasopharynx is the part of the pharynx that is located behind the oral cavity and above the larynx.
- c. The nasopharynx is the part of the pharynx that is located behind the oral cavity and above the larynx.
- d. The nasopharynx is the part of the pharynx that is located behind the oral cavity and above the larynx.
- e. The nasopharynx is the part of the pharynx that is located behind the oral cavity and above the larynx.

15. The Hypopharynx clinically subdivides into three regions:

- a. Pyriform sinus (fossa): lies on the either side of the larynx, extends from pharyngoepiglottic fold to the upper end of esophagus
- b. Post-cricoid region: the part of the anterior wall of laryngopharynx between the upper and lower borders of cricoid lamina
- c. Posterior pharyngeal wall: extends from the level of hyoid bone to the level of cricoarytenoid
- d. Anterior pharyngeal wall: extends from the level of hyoid bone to the level of cricoarytenoid
- e. Post-cricoid region: the part of the posterior wall of laryngopharynx between the upper and lower borders of cricoid lamina

16. Functions of the Hypopharynx are:

- a. Conduit for passage of air or food
- b. Breathing
- c. Helps in pharyngeal phase of deglutition
- d. Air conditioning of the inspired air: warming and humidifying
- e. Forms part of vocal tract for certain speech sounds

17. Peritonsillar abscess (quinsy) complications are:

- a. Parapharyngeal abscess
- b. Larynx edema
- c. Septicemia
- d. Pneumonitis or lung abscess
- e. Spontaneous haemorrhage from carotid vein or jugular artery

18. On the lateral wall of the nasal cavities we can find:

- a. The middle meatus
- b. The nasal septum
- c. The inferior meatus
- d. The maxillary sinus ostium
- e. The adenoids

19. Nasal septum deviation can lead to:

- a. Sinusitis
- b. Oral respiration
- c. Hipo- Anosmia

✓ epistaxis

✓ external deformity

✓ middle ear infection

HA